

# CYCLOPHOSPHAMIDE

Cyclophosphamide was considered by previous IARC Working Groups in 1980 and 1987 ([IARC, 1981](#), [1987a](#)). Since that time, new data have become available, these have been incorporated into the *Monograph*, and taken into consideration in the present evaluation.

## 1. Exposure Data

### 1.1 Identification of the agent

*Chem. Abstr. Serv. Reg. No.:* 50-18-0


*Chem. Abstr. Name:* 2H-1,3,2-

Oxazaphosphorin-2-amine, *N,N*-bis(2-chloroethyl)tetrahydro-, 2-oxide

*IUPAC Systematic Name:* *N,N*-Bis(2-chloroethyl)-1-oxo-6-oxa-2-aza-1λ<sup>5</sup>-phosphacyclohexan-1-amine

*Synonyms:* 2-[Bis(2-chloroethyl)amino] tetrahydro-2*H*-1,3,2-oxazaphosphorin 2-oxide; bis(2-chloroethyl)phosphoramidate cyclic propanolamide ester; *N,N*-bis(β-chloroethyl)-*N',O*-trimethylenephosphoric acid ester diamide; *N,N*-bis(2-chloroethyl)-*N',O*-propylenephosphoric acid ester diamide; Cytoxan; Endoxan; Neosar  
*Description:* Crystalline solid [anhydrous form] ([O'Neil, 2006](#))

#### 1.1.1 Structural and molecular formulae, and relative molecular mass



$C_7H_{15}Cl_2N_2O_2P$

Relative molecular mass: 261.1

### 1.2 Use of the agent

Cyclophosphamide is an antineoplastic agent metabolized to active alkylating metabolites with properties similar to those of chlormethine. It also possesses marked immunosuppressant properties. It is widely used, often in combination with other agents, in the treatment of several malignant diseases. Information for Section 1.2 is taken from [McEvoy, \(2007\)](#), [Royal Pharmaceutical Society of Great Britain \(2007\)](#), and [Sweetman \(2008\)](#).

### 1.2.1 Indications

Cyclophosphamide is used in the treatment of chronic lymphocytic leukaemia, lymphomas, soft tissue and osteogenic sarcoma, and solid tumours. It is given orally or intravenously. Cyclophosphamide is inactive until metabolized by the liver.

#### (a) *Hodgkin lymphoma*

Cyclophosphamide is used in combination regimens (e.g. bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, and prednisone [known as BEACOPP]) for the treatment of Hodgkin lymphoma.

#### (b) *Non-Hodgkin lymphoma*

Cyclophosphamide is used in combination therapy for the treatment of non-Hodgkin lymphoma, including high-grade lymphomas, such as Burkitt lymphoma and lymphoblastic lymphoma, as well as intermediate- and low-grade lymphomas. Cyclophosphamide is commonly used with doxorubicin (hydroxydaunorubicin), vincristine (oncovin), and prednisone (known as the CHOP regimen), with or without other agents, in the treatment of various types of intermediate-grade non-Hodgkin lymphoma. Cyclophosphamide has also been used as a single agent in the treatment of low-grade lymphomas.

#### (c) *Multiple myeloma*

Cyclophosphamide is used in combination with prednisone, or as a component of combination chemotherapy (i.e. vincristine, carmustine, melphalan, cyclophosphamide, and prednisone [VBMCP]) for the treatment of multiple myeloma.

#### (d) *Leukaemia*

Cyclophosphamide is used commonly for the treatment of chronic lymphocytic (lymphoblastic) leukaemia. Cyclophosphamide is used in combination with busulfan as a conditioning regimen before allogeneic haematopoietic

progenitor cell transplantation in patients with chronic myelogenous leukaemia.

Cyclophosphamide is used in the treatment of acute lymphoblastic leukaemia, especially in children. In the treatment of acute myeloid (myelogenous, non-lymphocytic) leukaemia, cyclophosphamide has been used as an additional drug for induction or post-induction therapy.

#### (e) *Cutaneous T-cell lymphoma*

Cyclophosphamide is used alone or in combination regimens for the treatment of advanced mycosis fungoides, a form of cutaneous T-cell lymphoma.

#### (f) *Neuroblastoma*

Cyclophosphamide alone is used in the treatment of disseminated neuroblastoma. Combination chemotherapy that includes cyclophosphamide is also used for this neoplasm.

#### (g) *Cancer of the ovary*

Cyclophosphamide is used in combination chemotherapy (vincristine, actinomycin D, and cyclophosphamide [VAC]) as an alternative regimen for the treatment of ovarian germ-cell tumours.

Cyclophosphamide has been used in combination with a platinum-containing agent for the treatment of advanced (Stage III or IV) epithelial ovarian cancer.

#### (h) *Retinoblastoma*

Cyclophosphamide is used in combination therapy for the treatment of retinoblastoma.

#### (i) *Cancer of the breast*

Cyclophosphamide is used alone and also in combination therapy for the treatment of breast cancer.

Combination chemotherapy with cyclophosphamide is used as an adjunct to surgery in premenopausal and postmenopausal women

with node-negative or -positive early (TNM Stage I or II) breast cancer. Adjuvant combination chemotherapy that includes cyclophosphamide, methotrexate, and fluorouracil has been used extensively.

Adjuvant combination chemotherapy (e.g. cyclophosphamide, methotrexate, and fluorouracil; cyclophosphamide, adriamycin, and fluorouracil; cyclophosphamide and adriamycin with or without tamoxifen) is used in patients with node-positive early breast cancer (Stage II) in both premenopausal and postmenopausal patients once treatment to control local disease (surgery, with or without radiation therapy) has been undertaken.

In Stage III (locally advanced) breast cancer, combination chemotherapy (with or without hormonal therapy) is used sequentially following surgery and radiation therapy for operable disease or following biopsy and radiation therapy for inoperable disease; commonly employed effective regimens include cyclophosphamide, methotrexate, and fluorouracil; cyclophosphamide, doxorubicin, and fluorouracil; and cyclophosphamide, methotrexate, fluorouracil, and prednisone. These and other regimens also have been used in the treatment of more advanced (Stage IV) and recurrent disease.

#### (j) *Small cell cancer of the lung*

Cyclophosphamide is used in combination chemotherapy regimens (e.g. cyclophosphamide, adriamycin, and vincristine [CAV]; cyclophosphamide, adriamycin, and etoposide [CAE]) for the treatment of extensive-stage small cell lung cancer.

#### (k) *Sarcoma*

Cyclophosphamide has been used in combination regimens (usually with dactinomycin and vincristine) and as an adjunct to surgery and radiation therapy in the treatment of rhabdomyosarcoma and Ewing sarcoma.

### 1.2.2 *Dosage*

Cyclophosphamide is administered orally or by intravenous injection or infusion. Less frequently, the drug has been administered intramuscularly and by intracavitary (e.g. intrapleural, intraperitoneal) injection and direct perfusion.

In patients with no haematological deficiencies receiving cyclophosphamide monotherapy, induction therapy in adults and children is usually initiated with an intravenous cyclophosphamide loading dose of 40–50 mg/kg administered in divided doses over 2–5 days. Other regimens for intravenous administration include 10–15 mg/kg every 7–10 days or 3–5 mg/kg twice weekly.

When cyclophosphamide is administered orally, the usual dose for induction or maintenance therapy is 1–5 mg/kg daily, depending on the tolerance of the patient.

A daily oral dose of 2–3 mg/kg for 60–90 days has been used in children with nephrotic syndrome, and in whom corticosteroids have been unsuccessful. In patients who are to undergo stem-cell transplantation, very high doses of cyclophosphamide such as 60 mg/kg daily for 2 days may be given as part of the conditioning regimen.

Various cyclophosphamide-containing combination chemotherapy regimens have been used in the treatment of breast cancer. One commonly employed regimen for the treatment of early breast cancer includes a cyclophosphamide dosage of 100 mg/m<sup>2</sup> orally on Days 1 through 14 of each cycle combined with intravenous methotrexate at 40 mg/m<sup>2</sup> on Days 1 and 8 of each cycle, and intravenous fluorouracil at 600 mg/m<sup>2</sup> on Days 1 and 8 of each cycle. In patients older than 60 years of age, the initial intravenous methotrexate dosage is reduced to 30 mg/m<sup>2</sup> and the initial intravenous fluorouracil dosage is reduced to 400 mg/m<sup>2</sup>. Dosage is also reduced if myelosuppression develops. Cycles

are generally repeated monthly (i.e. allowing a 2-week rest period between cycles) for a total of 6–12 cycles (i.e. 6–12 months of therapy).

Cyclophosphamide is available as 25 and 50 mg tablets for oral administration, and as 200 mg, 500 mg, 1 g, or 2 g vials of powder for reconstitution for parenteral administration.

### 1.2.3 Trends in use

No information was available to the Working Group.

## 2. Cancer in Humans

The carcinogenicity of cyclophosphamide in humans was established initially on the basis of a large number of case reports, as well as several epidemiological studies (IARC 1981, 1987a). The interpretation of the epidemiological studies was limited by the small numbers of cases, the difficulty in separating the role of cyclophosphamide from other agents, or both factors.

The most substantial evidence available to previous Working Groups was a Danish study of 602 patients treated “mainly with cyclophosphamide” for non-Hodgkin lymphoma, in which nine cases of acute myeloid leukaemia were observed compared to 0.12 expected (Pedersen-Bjergaard *et al.*, 1985), and a case–control study of leukaemia following ovarian cancer in the former German Democratic Republic where a strong dose–response relationship was observed (Haas *et al.*, 1987). All other studies reported at most three cases of leukaemia or bladder cancer in people who had received cyclophosphamide as the only potentially carcinogenic agent (IARC, 1981; Kinlen, 1985; Greene *et al.*, 1986).

Subsequently, further studies have been published that have provided more detailed information on the carcinogenicity of cyclophosphamide. This review is restricted to epidemiological studies that have used appropriate comparison

groups to investigate the role of cyclophosphamide as the cause of specific types of cancer.

There have been several reported cohort studies in which patients treated with cyclophosphamide were followed up, and the occurrence of second cancers investigated. Valagussa *et al.* (1994) followed 2465 women who had received treatment with cyclophosphamide, methotrexate and fluorouracil, a combination in which only cyclophosphamide is considered to have carcinogenic potential in humans. There were three cases of acute myeloid leukaemia observed compared to 1.3 expected, and five cases of bladder cancer compared to 2.1 expected. Statistical significance was not reported but was calculated by the Working Group to be greater than 0.05 for both types of cancer. Smith *et al.* (2003) followed 8563 women who had received cyclophosphamide and doxorubicin as adjuvant therapy for breast cancer and observed 43 cases of acute myeloid leukaemia or myelodysplastic syndromes (AML/MDS). The incidence of AML/MDS overall was seven times higher than expected rates in the general population, and was increased 3-fold in regimens that contained double the cumulative dose of cyclophosphamide.

Several case–control studies have also been reported. For leukaemia, Kaldor *et al.* (1990) investigated 114 cases of a cohort of ovarian cancer patients. The relative risks were, respectively, 2.2 and 4.1 in two increasing dose categories of cyclophosphamide. Neither increase was reported as statistically significant. Travis *et al.* (1994) carried out a study involving 35 cases of leukaemia following non-hodgkin lymphoma, and found that prior treatment with cyclophosphamide was associated with a relative risk of 1.8 that was not statistically significant when comparison was made to treatment with radiotherapy alone. In an investigation by Nandakumar *et al.* (1991) of 97 cases of myeloid leukaemia as second primary cancers, patients receiving cyclophosphamide had a relative risk of 12.6 compared to those treated surgically, and

was substantially higher when prednisone was co-administered with cyclophosphamide. [Curtis et al. \(1992\)](#) compared 90 women who developed acute myeloid leukaemia following breast cancer to controls, and found that the risk of leukaemia was 2.6 times greater in those who had received cyclophosphamide, compared to women who had been treated by surgery only.

There have also been two case-control studies of bladder cancer in relation to cyclophosphamide. [Kaldor et al. \(1995\)](#) investigated 63 cases of bladder cancer following ovarian cancer, and found that in comparison to surgery alone, the relative risk associated with chemotherapy containing cyclophosphamide as the only potential bladder-cancer-causing agent was 4.2 ( $P = 0.025$ ) in the absence of radiotherapy, and 3.2 ( $P = 0.08$ ) with radiotherapy. [Travis et al. \(1995\)](#) studied 31 cases of bladder cancer and 17 cases of kidney cancer as well as matched controls within a cohort of 2-year survivors of non-Hodgkin lymphoma. The relative risk associated with cyclophosphamide treatment was 4.5 ( $P < 0.05$ ) for bladder cancer, and 1.3 for kidney cancer.

## 2.1 Synthesis

The studies summarized above provide a comprehensive epidemiological basis for identifying cyclophosphamide as an independent cause of acute myeloid leukaemia and bladder cancer, that fully supports the conclusions drawn from earlier case reports, and more limited studies. Several studies have assessed the risk of all second primary cancers following cyclophosphamide treatment, and some have found rates of occurrence that appear to be elevated, but have not provided evidence for risk of other specific cancer types.

## 3. Cancer in Experimental Animals

Cyclophosphamide has been tested for carcinogenicity by oral administration to mice and rats, by subcutaneous injection to mice, by topical application to mice, by intravenous injection to rats, by intraperitoneal injection to mice and rats, and by perinatal exposure to mice.

Oral administration of cyclophosphamide resulted in skin tumours in transgenic mice ([Yamamoto et al., 1996](#); [Eastin et al., 2001](#)), and in urinary bladder carcinoma, leukaemia, and nervous system tumours in rats ([Schmähl & Habs, 1979](#); [Habs & Schmähl, 1983](#)). Subcutaneous injection of cyclophosphamide to mice caused a variety of neoplasms, including mammary gland carcinoma and leukaemia ([Schmähl & Osswald, 1970](#); [Walker & Bole, 1971, 1973](#); [Walker & Anver, 1979, 1983](#); [Petru et al., 1989](#)).

Intravenous injection of cyclophosphamide to rats caused both benign and malignant neoplasms ([Schmähl, 1967, 1974](#); [Schmähl & Osswald, 1970](#)).

Intraperitoneal administration of cyclophosphamide increased the incidences of lung adenoma and adenocarcinoma, bladder papilloma, and leukaemia in mice ([Shimkin et al., 1966](#); [Weisburger et al., 1975](#); [Mahgoub et al., 1999](#)), and mammary gland adenoma and carcinoma in rats ([Weisburger et al., 1975](#)).

Administration of cyclophosphamide to newborn mice caused lung and liver adenoma and carcinoma, and Harderian gland adenoma ([Kelly et al., 1974](#); [McClain et al., 2001](#)).

See [Table 3.1](#).

Table 3.1 Studies of cancer in experimental animals exposed to cyclophosphamide

Route Species, strain (sex), age Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Significance	Comments
<b>Oral administration</b>				
Mouse, Tg ras H2/CB6F1 & B6C6F1 (M), 9 wk 26 wk <a href="#">Yamamoto et al. (1996)</a>	0, 10, 30 mg/kg bw by gavage (in water, volume NR), twice/wk for 25 wk Initial number/group NR	Tg ras H2/CB6F1: Lung (adenomas)– 0/9, 3/16, 3/27 Multiplicity– 0, 0.19, 0.11 tumours/mouse CB6F1: Lung (adenomas)– 0/6, 2/18, 2/20 Multiplicity– 0, 0.11, 0.10 tumours/mouse	[NS] <sup>a</sup>	Pharmaceutical grade
Mouse, Tg.AC (M, F), 8–9 wk 27 wk <a href="#">Eastin et al. (2001)</a>	0, 10, 30, 60 mg/kg bw by gavage (in water 50% ethanol, volume NR); twice/wk for 26 wk 15/sex/group	Skin tumours (at all sites; histologically confirmed): 5/15, 1/2, 5/5, 5/15 (M); 2/15, 5/11, 11/11, 14/15 (F) Skin tumours (squamous cell papillomas of vulva): 2/15, 4/11, 10/11, 12/15 (F) Leukaemia (erythrocytic): 0/15, 0/15, 4/15, 1/15 (F)	[P < 0.0001 for 30 and 60 mg/kg bw doses in female mice] <sup>a</sup> [P ≤ 0.0003 for 30 and 60 mg/kg bw doses in female mice] <sup>a</sup> P < 0.05, for 30 mg/kg bw group	Purity NR; Tg.AC mice are transgenic mice that carry a v-Ha-ras oncogene
Rat, Sprague-Dawley (M, F) Lifetime <a href="#">Schmähl &amp; Habs (1979)</a>	0, 0.31, 0.63, 1.25, 2.5 mg/kg bw in drinking-water, 5 ×/wk for life 40/sex/group	Malignant tumours: 4/38, 11/34, 14/36, 15/35, 13/31 (M); 5/34, 11/37, 13/37, 11/33, 9/27 (F) Urinary bladder (carcinomas): 0/38, 2/34, 2/36, 5/35, 7/31 (M); 0/34, 0/37, 0/37, 0/33, 1/27 (F) Lymphoid and haematopoietic tissue (leukaemia): 0/72, 3/71, 6/73, 6/68, 4/58 (M, F) Nervous system (sarcomas): 1/72, 7/71, 5/73, 6/68, 1/58 (M, F)	[P < 0.05, for 3 highest doses] [P ≤ 0.02 for 2 highest doses in males] <sup>a</sup> [P ≤ 0.04 for 3 highest doses for combined males and females] <sup>a</sup> [P ≤ 0.05 for 0.31 and 1.25 mg/kg doses for combined males and females] <sup>a</sup>	Purity NR

Table 3.1 (continued)

Route Species, strain (sex), age Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Significance	Comments
Rat, Sprague-Dawley (M), 100 d 20 mo <a href="#">Habs &amp; Schmähel (1983)</a>	0, 2.5 mg/kg bw in drinking-water, 5 x/wk for 20 mo 100/group	Urinary bladder (papillomas or transitional-cell carcinomas): 0/63, 24/80 Nervous system tumours: 1/63, 11/80	[ <i>P</i> < 0.0001] <sup>a</sup>  [ <i>P</i> < 0.0076] <sup>a</sup>	Reported as “chemically pure”
Rat, Sprague-Dawley (M), 100 <sup>a</sup> Lifetime <a href="#">Schmähel &amp; Habs (1983)</a>	0, 2.5 mg/kg bw in drinking-water, 5 times/wk for life 100/group	Urinary bladder (papillomas): 0/100, 15/100 Urinary bladder (transitional- cell carcinomas): 0/100, 17/100	[ <i>P</i> < 0.0001] <sup>a</sup>  [ <i>P</i> < 0.0001] <sup>a</sup>	Purity NR; only data on bladder tumours reported
<b>Subcutaneous injection</b>				
Mouse, NMRI (F) 52 wk <a href="#">Schmähel &amp; Osswald (1970)</a>	0, 26 mg/kg bw/wk (in solvent NR), for 5 wk 50/group	Malignant tumours (primarily mammary carcinomas): 3/46, 28/46	[ <i>P</i> < 0.001] <sup>b</sup>	Purity > 98%
Mouse, New Zealand Black/New Zealand White (F) 64 wk <a href="#">Walker &amp; Bole (1971)</a>	0, 8 mg/kg bw (in saline; volume NR), daily for 64 wk 16, 10	Neoplasms (mainly lymphomas): 0/16, 6/10	<i>P</i> = 0.00002	Purity NR
Mouse, New Zealand Black/New Zealand White (M, F) 93 wk <a href="#">Walker &amp; Bole (1973)</a>	0, 1, 8 mg/kg bw (in 100 µL saline), daily for 93 wk 20, 10, 10 per sex	Neoplasms (mainly lymphomas): 2/16, 3/9, 8/9 (M); 1/20, 1/10, 9/9 (F)	<i>P</i> = 0.003 for 8 mg/kg bw males; <i>P</i> < 0.0001 for 8 mg/kg bw females	Purity NR
Mouse, New Zealand Black/New Zealand White (F) Lifetime <a href="#">Walker &amp; Anver (1979)</a>	0, 5.7, 16 mg/kg bw (in 100 µL saline), daily for life 15, 17, 21	Neoplasms (mainly mammary carcinomas): 0/13, 15/15, 17/19 Mammary carcinomas: 0/13, 5/15, 16/19	[ <i>P</i> < 0.0001 for 5.7 and 16 mg/kg bw groups] <sup>a</sup>  [ <i>P</i> ≤ 0.03 for 5.7 and 16 mg/kg bw groups] <sup>a</sup>	Purity NR; treatment groups not started simultaneously

Table 3.1 (continued)

Route Species, strain (sex), age	Dosing regimen Animals/group at start	Incidence of tumours	Significance	Comments
Mouse, New Zealand Black/New Zealand White (F), 6 wk Lifetime <a href="#">Walker &amp; Anver (1983)</a>	0, 56 mg/kg bw (in 100 µL saline), weekly for life 15, 22	Neoplasms: 0/13, 17/19	[ <i>P</i> < 0.0001] <sup>a</sup>	Purity NR; groups not started simultaneously; Neoplasms were mainly mammary gland carcinomas, lung adenomas and lymphomas
Mouse, NMRI & AKR (F), 7 wk, Lifetime <a href="#">Petru et al. (1989)</a>	0, 13, 26 mg/kg bw (in saline, volume NR), weekly for life 30/group	Leukaemia (NMRI mice): 2/30, 16/30, 10/30 Leukaemia (AKR mice): 30/30, 25/30, 19/30	<i>P</i> ≤ 0.027 for 13 & 26 mg/kg bw groups <i>P</i> ≤ 0.006 for 13 & 26 mg/kg bw groups	Purity NR [negative trend in AKR mice]
<b>Skin application</b>				
Mouse, Tg.AC (M, F), 8–9 wk 27 wk <a href="#">Eastin et al. (2001)</a>	0, 10, 30, 90 mg/kg bw (in 50% ethanol, 3.3 mL/kg bw), 2 ×/wk for 26 wk 15/sex/group	Skin tumours (at site of application): 1/15, 0/15, 2/15, 3/15 (M); 1/15, 0/15, 0/15, 2/15 (F) Skin tumours (at all skin sites): 1/15, 2/15, 3/15, 3/15 (M); 4/15, 3/15, 9/15, 14/15 (F)	[NS] <sup>a</sup>	Purity NR; Tg.AC mice are transgenic mice that carry a v-Ha- <i>ras</i> oncogene
<b>Intravenous administration</b>				
Rat, BR 46 (M) 23 mo <a href="#">Schmähel (1967)</a>	0, 15 mg/kg bw (vehicle and volume NR), weekly (750 mg/kg bw total dose) 50, 40	Neoplasms (benign and malignant combined): 1/50, 14/26	[ <i>P</i> < 0.001] <sup>b</sup>	Purity > 98%
Rat, BR 46 (M) 23 mo <a href="#">Schmähel &amp; Osswald (1970)</a>	0, 13 mg/kg bw (vehicle and volume NR), weekly for 52 wk 89, 48	Neoplasms: 3/65, 4/36 (benign); 4/65, 6/36 (malignant)	[NS] <sup>b</sup>	Purity > 98%
Rat, BR 46 (M) 23 mo <a href="#">Schmähel &amp; Osswald (1970)</a>	0, 33 mg/kg bw (vehicle and volume NR), 5 times every 2 wk 89, 96	Neoplasms: 3/65, 5/66 (benign); 4/65, 16/66 (malignant)	[ <i>P</i> < 0.01, malignant tumours] <sup>b</sup>	Purity > 98%
Rat, Sprague-Dawley (M) 700 d <a href="#">Schmähel (1974)</a>	0, 13 mg/kg bw (vehicle and volume NR), weekly (670 mg/kg bw total dose) 52, 32	Neoplasms (malignant): 6/52, 14/32	[ <i>P</i> < 0.001] <sup>b</sup>	Purity > 98%

Table 3.1 (continued)

Route Species, strain (sex), age Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Significance	Comments
<b>Intraperitoneal administration</b>				
Mouse, dd (M, F) 48 wk <a href="#">Tokuoka (1965)</a>	0 or 5 mg/kg bw (in saline 5 mL/ kg), 2 injections/wk for 15 wk 20, 29	Lung (adenomas or carcinomas): 1/20, 3/29 Liver (adenomas): 0/20, 2/29 Testis (interstitial cell tumours): 0/20, 4/29 Mammary gland (carcinomas): 1/20, 3/29	NS NS NS NS	Purity NR
Mouse, A (M, F) 48 wk <a href="#">Tokuoka (1965)</a>	0 or 5 mg/kg bw (in saline 10 mL/ kg), 2 injections/wk for 15 wk 16, 25	Lung (adenomas or carcinomas): 2/16, 6/25 Testis (interstitial cell tumours): 0/16, 3/25	NS NS	Purity NR
Mouse, A/I (M, F, equally split) 39 wk <a href="#">Shimkin et al. (1966)</a>	0, 32.2, 129, 516, 1609 µmol/kg bw (total dose; in 200 µL water), 3 injections/wk for 4 wk 360, 30, 30, 30, 30	Lung (adenomas or adenocarcinomas): 107/339, 12/30, 11/26, 20/27, 2/4 (incidence); 0.38, 0.4, 0.6, 1.3, 2.5 (tumours per mouse)	[p < 0.001 (for 516 µmol/ kg bw dose, incidence)] <sup>b</sup>	Purity NR
Mouse, Swiss-Webster- derived (M, F) 18 mo <a href="#">Weisburger et al. (1975)</a>	0, 12, 25 mg/kg bw (vehicle and volume NR), 3 injections/wk for 6 mo 101, 25, 25 (M) 153, 25, 25 (F)	Lung (adenomas or adenocarcinomas): 10/101, 7/30 (M); 21/153, 10/35 (F) Bladder (papillomas or carcinomas): 3/101 & 4/30 (M)	P = 0.031 (M) and P = 0.027 (F) (combined 12 & 25 mg/kg bw vs control) P = 0.048 (combined 12 & 25 mg/kg bw vs control)	Purity NR; not all control mice were treated with the vehicle

Table 3.1 (continued)

Route Species, strain (sex), age	Dosing regimen Animals/group at start	Incidence of tumours	Significance	Comments
Mouse, 129/Sv & 129/ Sv X C57BL/6 <i>Nf1<sup>+/+</sup></i> & <i>Nf1<sup>+/-</sup></i> (sex NR), 6–10 wk 15 mo <a href="#">Mahgoub et al. (1999)</a>	0 or 100 mg/kg bw/wk (solvent and volume NR) for 6 wk 129/Sv <i>Nf1<sup>+/+</sup></i> ; 31 & 5 mice 129/Sv <i>Nf1<sup>+/-</sup></i> ; 46 & 12 mice 129/Sv X C57BL/6 <i>Nf1<sup>+/+</sup></i> ; 14 & 15 mice 129/Sv X C57BL/6 <i>Nf1<sup>+/-</sup></i> ; 412 & 25 mice	Leukaemia (129/Sv <i>Nf1<sup>+/+</sup></i> ): 2/31, 0/5 Leukaemia (129/Sv <i>Nf1<sup>+/-</sup></i> ): 8/46, 7/12 Leukaemia (129/Sv X C57BL/6 <i>Nf1<sup>+/+</sup></i> ): 0/14, 2/25 Leukaemia (129/Sv X C57BL/6 <i>Nf1<sup>+/-</sup></i> ): 0/12, 7/25	$P = 0.004$	Purity NR
Rat, Sprague-Dawley (M, F) 18 mo <a href="#">Weisburger et al. (1975)</a>	0, 5, 10 mg/kg bw (vehicle and volume NR), 3 injections/wk for 6 mo 179, 25, 25 (M) 181, 25, 28 (F)	Mammary gland (adenomas): 2/105 & 24/53 (F; combined 5 & 10 mg/kg bw) Mammary gland (carcinomas): 13/105 & 9/53 (F; combined 5 & 10 mg/kg bw)	$P = 0.028$  $P = 0.035$	Purity NR; not all control rats were treated with the vehicle
<b>Perinatal exposure</b>				
Mouse, CD-1 (M, F) 79 wk <a href="#">Kelly et al. (1974)</a>	i.p. injection 0, 0.8, 4.0, 20.0 mg/kg bw (in 10 $\mu$ L/ kg saline), on postnatal Days 1, 3, 6 30/sex/group	Lung (adenomas): 0/28, 2/29, 4/27, 0/21 (M); 1/25, 2/27, 2/28, 3/21 (F)	$P < 0.05$ for 4 mg/kg bw males (life-table analysis)	Purity NR; the 20 mg/kg dose caused marked bw changes and nearly 100% mortality

Table 3.1 (continued)

Route Species, strain (sex), age Duration Reference	Dosing regimen Animals/group at start	Incidence of tumours	Significance	Comments
Mouse, CD-1 (M, F) 1 yr <a href="#">McClellan et al. (2001)</a>	Oral 0, 10, 20, 40, 60 mg/kg bw by gavage (100 µL and 200 µL) on postnatal Days 8 & 15 [solvent NR] 48 (control), 24/sex	Liver (adenomas): 2/48, 2/24, 4/24, 6/24, 5/24 (M)  Liver (carcinomas): 0/48, 0/24, 1/24, 6/24, 1/24 (M)	[P < 0.04 for 40 & 60 mg/kg bw] <sup>a</sup>  [P = 0.0009 for 40 mg/ kg bw] <sup>a</sup>	Purity NR
		Lung (adenomas): 3/48, 0/24, 8/24, 12/24, 13/24 (M); 7/48, 3/24, 6/24, 16/24, 13/24 (F)  Lung (carcinomas): 0/48, 1/24, 0/24, 6/24, 3/24 (M); 0/48, 1/24, 3/24, 3/24, 0/24 (F)  Harderian gland (adenomas): 2/48, 1/24, 1/24, 1/24, 5/24 (F)	[P < 0.005 for 20, 40, & 60 mg/kg bw (M); 40 & 60 mg/kg bw (F)] <sup>a</sup>  [P < 0.03 for 40 & 60 mg/kg bw (M); 20 & 40 mg/kg bw (F)] <sup>a</sup>  [P < 0.04 for 60 mg/kg bw] <sup>a</sup>	
<b>Pre and postnatal exposure</b>				
Mouse, BR 46 (M, F) 24 mo <a href="#">Roschlau &amp; Justus (1971)</a>	i.p injection 25 mg/kg bw on gestation Day 14 [solvent and volume NR]. Male and female offspring treated every 2 wk for a total of 30 times Initial number NR	Lung (adenomas): male offspring 4/16, 2/16; female offspring 5/12 & 1/18  Lung (carcinomas): male offspring 0/16, 3/16; female offspring 0/12, 4/18	NS  NS	Purity NR

<sup>a</sup> Current Working Group analysis (Fisher Exact test)<sup>b</sup> Previous Working Group analysis

bw, body weight; d, day or days; F, female; i.p., intraperitoneal; M, male; mo, month or months; NR, not reported; NS, not significant; vs, versus; wk, week or weeks; yr, year or years

## 4. Other Relevant Data

### 4.1 Absorption, distribution, metabolism, and excretion

In most species, cyclophosphamide is rapidly absorbed, metabolized, and excreted. Its metabolic pathway has been studied in several species including mice, rats, hamsters, rabbit, dogs, sheep, and monkeys. Cyclophosphamide is not cytotoxic *per se*, because it requires metabolic activation before it can act as an alkylating agent. Activation takes place predominantly in the liver, although this may occur in other tissues ([IARC, 1981](#)).

Cyclophosphamide undergoes metabolism to several intermediates with alkylating activity. The principal metabolites identified are phosphoramidate mustard, and acrolein. Phosphoramidate mustard can undergo dephosphoramidation to yield nornitrogen mustard, which also has alkylating activity. Metabolites of cyclophosphamide can interact with DNA and proteins, resulting in the formation of adducts. The metabolism of cyclophosphamide and DNA adducts formation are summarized in Fig. 4.1.

A minor pathway results in dechloroethylation and the formation of 2-dechloroethylcyclophosphamide and another alkylating agent, chloroacetaldehyde ([Balu et al., 2002](#)).

The other compounds such as 4-ketocyclophosphamide and propionic acid derivative are relatively non-toxic, and are the major urinary metabolites of cyclophosphamide in several species ([IARC, 1981](#)).

### 4.2 Genetic and related effects

#### 4.2.1 Interaction with DNA

Using 4-hydroperoxycyclophosphamide as an activated form of cyclophosphamide, [Mirkes et al. \(1992\)](#) identified by mass spectrometric analysis the formation of the monofunctional


adduct *N*-(2-chloroethyl)-*N*-[2-(7-guaninyl)ethyl]amine (nor-G) and the bifunctional adduct *N,N*-bis[2-(7-guaninyl)ethyl]amine (G-nor-G) in rat embryos in in-vitro culture. The monofunctional adduct *N*-(2-hydroxyethyl)-*N*-[2-(7-guaninyl)ethyl]amine (nor-G-OH) was detected in bladder tissue of rats injected with [<sup>3</sup>H] cyclophosphamide ([Benson et al., 1988](#)). Using <sup>32</sup>P-postlabelling analysis, a phosphotriester was shown to be formed: (1) when phosphoramidate mustard was reacted with deoxyguanosine 5'-monophosphate, (2) when cyclophosphamide was incubated with calf thymus DNA in the presence of reconstituted cytochrome P450 (CYP) metabolizing system, and (3) in liver DNA from mice injected intraperitoneally with cyclophosphamide ([Maccubbin et al., 1991](#)).

Nornitrogen mustard reacts with guanosine and with guanine bases in DNA to form nor-G initially, but this is converted to a hydroxylated derivative (nor-G-OH), and to a cross-linked (between guanines) adduct (G-nor-G) ([Hemminki, 1987](#)). Both monofunctional adducts, but not the cross-linked adduct, were also detected when phosphoramidate mustard was reacted with DNA ([Cushnir et al., 1990](#)). Acrolein reacts with DNA to form *O*<sup>6</sup>-(*n*-propanalyl)guanine, and the product of chloroacetaldehyde reaction with DNA is *O*<sup>6</sup>-(ethanalyl)guanine ([Balu et al., 2002](#)). Acrolein can produce exocyclic adducts in DNA, including 1,*N*<sup>2</sup>-hydroxypropanodeoxyguanosine and 1,*N*<sup>6</sup>-hydroxypropanodeoxyadenosine ([Chung et al., 1984](#); [Foiles et al., 1990](#); [Smith et al., 1990](#)). The former was detected in acrolein-treated human fibroblasts and in peripheral blood lymphocytes of a dog treated with cyclophosphamide ([Wilson et al., 1991](#)).

Nornitrogen mustard also reacts covalently with proteins, and a method for the detection of cysteine-34 residue adducts in human serum albumin has been described ([Noort et al., 2002](#)).

The single-cell gel comet assay is used to detect single-strand breaks and other alkali-labile lesions in DNA exposed to cyclophosphamide.

Fig. 4.1 Metabolic pathway of cyclophosphamide



A. Metabolism of cyclophosphamide to phosphoramidate mustard, acrolein, and normitrogen mustard. Cyclophosphamide is metabolized by CYP enzymes to 4-hydroxycyclophosphamide, which equilibrates with aldophosphamide to spontaneously yield phosphoramidate mustard and acrolein. Aldophosphamide is also metabolized by aldehyde oxidase to carboxyphosphamide, which produces normitrogen mustard. 4-Hydroxy-cyclophosphamide can be oxidized to the inactive 4-keto-cyclophosphamide. B. Phosphoramidate mustard produces multiple monofunctional and bifunctional adducts with guanine, and acrolein forms exocyclic adducts. Normitrogen mustard forms mono- and bifunctional adducts with guanine.

From [Povirk & Shuker \(1994\)](#), [Anderson et al. \(1995\)](#), [Khan et al. \(1998\)](#)  
 CYP, cytochrome P450; nor-G, *N*-(2-chloroethyl)-*N*-[2-(7-guanylyl)ethyl]amine; G-nor-G, *N,N*-bis[2-(7-guanylyl)ethyl]amine; nor-G-OH, *N*-(2-hydroxyethyl)-*N*-[2-(7-guanylyl)ethyl]amine; dR, deoxyribose

*In vitro* studies have demonstrated the comet-forming activity of cyclophosphamide in human hepatoma (Hep G2) cells ([Uhl et al., 2000](#); [Yusuf et al., 2000](#)), in primary cultures of rat and human urinary bladder cells ([Robbiano et al., 2002](#)), in primary cultures of human leukocytes in the presence of metabolic activation system S9 mix ([Hartmann et al., 1995](#); [Hartmann & Speit, 1995](#); [Frenzilli et al., 2000](#)), and in extended-term cultures of human T-lymphocytes, also in the presence of S9 ([Andersson et al., 2003](#)). Comet formation was also detected *in vivo* in the urinary bladder mucosa of rats given cyclophosphamide orally ([Robbiano et al., 2002](#)), and in peripheral blood cells of patients administered the drug ([Hartmann et al., 1995](#)).

#### 4.2.2 Genotoxic effects in humans

There are few reports of DNA-adduct formation by cyclophosphamide in humans. Acrolein-derived DNA adducts, detected by immunochemical methods, were found in blood leukocytes of cancer patients receiving cyclophosphamide ([McDiarmid et al., 1991](#)). In another study, mono-adducts and inter-strand cross-links derived from phosphoramidate mustard were detected in a single patient administered 1 g/m<sup>2</sup> cyclophosphamide ([Souliotis et al., 2003](#)). Increased DNA damage (comet formation) was also observed in the lymphocytes of patients administered cyclophosphamide ([Hartmann et al., 1995](#)).

Increased frequencies of several biomarkers of genotoxicity have been observed in the lymphocytes of patients treated with cyclophosphamide, relative to control subjects. These include mutations at the hypoxanthine-(guanine) phosphoribosyl transferase (*HPRT*) locus ([Palmer et al., 1986](#), [1988](#); [Tates et al., 1994](#); [Sanderson et al., 2001](#)), and sister chromatid exchange ([Raposa & Várkonyi, 1987](#); [McDiarmid et al., 1990](#); [Sardaş et al., 1994](#); [Mertens et al., 1995](#); [Hartmann et al., 1995](#)).

Other studies reported positive findings for elevated chromosomal aberrations frequencies ([Sessink et al., 1994](#); [Rubes et al., 1998](#); [Burgaz et al., 2002](#)), and micronuclei ([Yager et al., 1988](#); [Tates et al., 1994](#); [Zúñiga et al., 1996](#); [Burgaz et al., 1999](#); [Rekhadevi et al., 2007](#)) in medical personnel exposed to cyclophosphamide. Increases in frequencies of micronuclei were also detected in buccal cells in some studies ([Cavallo et al., 2005](#); [Rekhadevi et al., 2007](#)), but not in another ([Burgaz et al., 1999](#)).

#### 4.2.3 Genotoxic effects in experimental systems

##### (a) Mutagenic effects *in vitro*

The previous *IARC Monograph* ([IARC, 1987b](#)) states that cyclophosphamide induced chromosomal aberrations, sister chromatid exchange, and DNA damage in human cells *in vitro*. It also induced morphological transformation, chromosomal aberrations, sister chromatid exchange, mutation, and unscheduled DNA synthesis (UDS) in rodent cells *in vitro*. It further induced aneuploidy, mutation, recombination, gene conversion, and DNA damage in fungi. It was also reported to act as a mutagen and DNA-damaging agent in bacteria.

The mutagenicity of cyclophosphamide in *Salmonella typhimurium* was enhanced by increased induction of CYPs in S9 liver fractions by a combination of  $\beta$ -naphthoflavone and sodium phenobarbital ([Paolini et al., 1991a](#)). Comparison of S9 from liver and kidney of pregnant mice revealed that liver S9 was more effective in activating cyclophosphamide to mutagenic metabolites in *S. typhimurium*, and also in inducing sister chromatid exchange in human peripheral lymphocytes, and Chinese hamster ovary (CHO) cells ([Winckler et al., 1987](#)).

In *Saccharomyces cerevisiae*, higher rates of mitotic gene conversion and point mutation by cyclophosphamide were associated with induction of class 2B CYPs in co-cultured epithelial cell

lines from fetal mouse liver (Paolini *et al.*, 1991b). A recombinant plasmid containing a full-length cDNA encoding the rat cytochrome CYP2B1 introduced into *S. cerevisiae* also increased the mutation frequency induced by cyclophosphamide (Black *et al.*, 1989).

CYP2B1 expressed in Chinese hamster V79-derived SD1 cell lines also potentiated cyclophosphamide mutagenesis (6-thioguanine resistance), whereas CYP1A1 expressed in V79-derived XEM<sub>2</sub> cell lines did not (Doehmer *et al.*, 1990, 1992).

Cyclophosphamide was weakly mutagenic (detected by induction of resistance to 6-thioguanine) in differentiated Reuber hepatoma cells H4IIEC3/G<sup>-</sup>, but markedly cytotoxic and clastogenic (micronucleus formation) (Roscher & Wiebel, 1988), and also mutagenic in a Chinese hamster epithelial liver cell line (6-thioguanine resistance) (Turchi *et al.*, 1992), and in Chinese hamster lung (CHL) cells in the presence of S9, as measured at microsatellite loci (Kikuno *et al.*, 1995).

Using 4-hydroperoxycyclophosphamide and phosphoradiazidic mustard, the role of different repair enzymes in defining sensitivity was investigated by Andersson *et al.* (1996) in CHO cells. Mutations in excision repair cross-complementing *ERCC1* and *ERCC4* genes caused hypersensitivity to the cyclophosphamide analogues.

Cyclophosphamide induced sister chromatid exchange in mouse primary bone-marrow and spleen cells (Soler-Niedziela *et al.*, 1989), and micronuclei in mouse lymphoma in L5178Y tk<sup>+/+</sup> cells (Kirsch-Volders *et al.*, 2003), and in parental V79 cells (Kalweit *et al.*, 1999) in the presence of rat liver S9. Of several V79 cell lines engineered to express rat CYPs, increases in micronuclei (Ellard *et al.*, 1991) and sister chromatid exchange (Kulka *et al.*, 1993) were seen in the cells expressing CYP2B1. The rat hepatoma cells lines H4IIEC3/G<sup>-</sup> and 2sFou were also susceptible to micronuclei induction by cyclophosphamide (Tafazoli *et al.*, 1995).

Human T-lymphocytes were more susceptible than B-lymphocytes to both chromosomal aberrations and sister chromatid exchange induction by cyclophosphamide in the presence of rat liver S9 (Miller 1991a, b). This difference between T- and B-lymphocytes was not found with mouse cells treated with 4-hydroxycyclophosphamide or phosphoramidic mustard (Kwanyuen *et al.*, 1990). In another study (Kugler *et al.*, 1987), rat liver microsomal mix was more effective than rat liver S9 in activating cyclophosphamide to induce chromosomal aberrations. Human lymphocytes from women carrying mutations in the breast cancer susceptibility gene *BRCA1* were more susceptible to micronuclei induction than cells from non-carriers (Trenz *et al.*, 2003). Hep G2 human hepatoma cells were susceptible to sister chromatid exchange and micronuclei induction by cyclophosphamide (Natarajan & Darroudi, 1991) and, in analogous studies, the S9 microsomal fraction of these cells were shown to be capable in activating cyclophosphamide to induce sister chromatid exchange and micronuclei in CHO cells (Darroudi & Natarajan, 1993). Human dental pulp cells formed chromosomal aberrations when exposed to cyclophosphamide in the presence of rat liver S9 (Tsutsui *et al.*, 2006).

In the presence of rat liver S9, cyclophosphamide induced morphological transformation of BALB/3T3 mouse embryonic fibroblast cells (McCarvill *et al.*, 1990).

#### (b) Mutagenic effects *in vivo*

The previous IARC Monograph (IARC, 1987b) states that cyclophosphamide was found to bind to kidney, liver and lung DNA in mice. It also induced dominant lethality, chromosomal aberrations, micronuclei, sister chromatid exchange, mutations, and DNA damage in rodents *in vivo*. In *Drosophila*, it induced aneuploidy, heritable translocations, and somatic and sex-linked recessive lethal mutations. In patients administered cyclophosphamide, increased incidences of chromosomal aberrations and sister chromatid

exchange in peripheral lymphocytes and bone marrow were observed.

In *Drosophila melanogaster*, cyclophosphamide tested positive for the somatic white-ivory mutation ([Batiste-Alentorn et al., 1994](#)), and produced chromosome breaks in spermatocytes ([Zijlstra & Vogel, 1989](#)).

Several studies have examined the mutagenic effects of cyclophosphamide in transgenic mice. In MutaMouse, mutation induction was observed in bone marrow (other tissues not studied) ([Hoorn et al., 1993](#)). In Big Blue mice, mutation frequencies were elevated in the liver, but not in the testis or spleen in one study ([Hoyes et al., 1998](#)), and in another study, in the lung and urinary bladder, but not in the kidney, bone-marrow or splenic T-cells ([Gorelick et al., 1999](#)). Another study compared the *lacI* locus in Big Blue mice with the *Hprt* locus in conventional B6C3F1 mice, and cyclophosphamide induced mutations in the endogenous gene in splenic lymphocytes, but not in the transgene ([Walker et al., 1999](#)). In rats, cyclophosphamide produced the ‘common deletion’ mutation in liver mitochondrial DNA, and folic acid supplementation was found to be protective against this damage ([Branda et al., 2002](#)).

In two related studies investigating oncogene and tumour-suppressor gene expression in mice, cyclophosphamide was found to induce expression of several genes, including *c-Myc* and *Tp53*, in the spleen and thymus, but not in other tissues ([Ember et al., 1995](#); [Ember & Kiss, 1997](#)).

Many studies have investigated the cytogenicity of cyclophosphamide in newts, rodents, dogs, and non-human primates. Results are invariably positive for this compound, and are summarized in [Table 4.1](#).

#### (c) Mutagenic effects in germ cells

[Anderson et al. \(1995\)](#) reviewed the activity of cyclophosphamide in germ cells, and in summary, the germ cell stages that are most sensitive to cyclophosphamide are the postmeiotic stages.

Tests for germ-cell damage that examine effects in  $F_1$  progeny in which cyclophosphamide gave positive results include dominant lethality, heritable translocations, specific locus mutations, and malformations. Although cyclophosphamide is not an effective aneugen, it causes structural and numerical chromosomal damage in second meiotic metaphases and first cleavage metaphases, and in  $F_1$  embryos. It is also positive for inducing sister chromatid exchange in germ cells and causes abnormal sperm-head morphology. Most studies have been carried out in mice, but positive results have also been observed in rats and rabbits, e.g. induction of unscheduled DNA synthesis in the testes (reviewed in [Anderson et al., 1995](#)), and also in hamsters ([Waters & Nolan, 1995](#)).

More recent studies in mice have demonstrated the dominant lethal effects of cyclophosphamide ([Dobrzyńska et al., 1998](#)) as well as intrachromosomal gene conversion and mutation events primarily in meiotic stage cells ([Schimenti et al., 1997](#)). In female rats, administration of cyclophosphamide at 16 days of gestation significantly increased nucleolar and synaptonemal complex fragmentation ([Cusidó et al., 1995](#)), and in male rats chronic exposure to cyclophosphamide disrupted meiotic events before pachynema during spermatogenesis ([Barton et al., 2003](#)).

#### (d) Modulation of mutagenicity by other agents

A large number of studies have investigated the effects of agents in modulating the genotoxicity of cyclophosphamide, and are summarized in [Table 4.2](#).

### 4.3 Mechanisms of carcinogenesis

All of the available evidence indicates that cyclophosphamide exerts its carcinogenic activity via a genotoxic mechanism ([McCarroll et al., 2008](#)). The metabolite widely thought to be responsible for the antitumour activity

**Table 4.1 Positive cytogenicity studies of cyclophosphamide in newts, rodents, dogs, and non-human primates**

<b>Species</b>	<b>Cytogenetic end-point investigated</b>	<b>Additional considerations</b>	<b>Reference</b>
Mouse	SCE	Bone-marrow cells. Reduction in frequency with increasing numbers of cell division	<a href="#">Morales-Ramírez et al. (1990)</a>
Mouse	SCE	Bone-marrow cells. A comparison of wild and laboratory mice	<a href="#">Huang et al. (1990)</a>
Mouse	MN	Bladder epithelial cells	<a href="#">Konopacka (1994)</a>
Mouse	CA	Bone-marrow cells. Effects of malnutrition and alcohol	<a href="#">Terreros et al. (1995)</a>
Mouse	MN	Peripheral blood reticulocytes and PCE in bone marrow	<a href="#">Hatanaka et al. (1992)</a>
Mouse	MN	Splenocytes	<a href="#">Benning et al. (1992)</a>
Mouse	MN	Bone-marrow PCE. Comparison of i.p. and p.o. administration	<a href="#">Wakata et al. (1989)</a>
Mouse	MN	7 organs compared (bone marrow, forestomach, stomach, small intestine, large intestine, urinary bladder, lung)	<a href="#">Sycheva (2001)</a>
Mouse	Intrachromosomal recombination	Spleen cells Transgenic mouse model with <i>lacZ</i> transgenic expression depending on somatic interchromosomal inversion	<a href="#">Sykes et al. (1998)</a>
Mouse	MN	PCE in adult bone-marrow cells and fetal liver cells. Male, female, pregnant female, and fetal mice compared	<a href="#">Harper et al. (1989)</a>
Mouse	MN SCE	Transplacental exposure; fetal liver cells	<a href="#">Porter &amp; Singh (1988)</a>
Mouse	MN CA	Bone-marrow and peripheral blood cells (CA) and peripheral blood erythrocytes (MN). Chronic ingestion of cyclophosphamide; results positive for MN, negative for CA	<a href="#">Director et al. (1998)</a>
Mouse	MN	Bone-marrow cells. In-vivo/in-vitro assay	<a href="#">Odagiri et al. (1994)</a>
Mouse	CA SCE	Bone-marrow and spleen cells. In-vivo/in-vitro assay vs in-vivo assay	<a href="#">Krishna et al. (1987)</a>
Mouse	SCE	Bone-marrow and spleen cells. In-vivo/in-vitro assay vs in-vivo assay	<a href="#">Krishna et al. (1988)</a>
Rat	CA	Liver cells of neonates exposed <i>in utero</i>	<a href="#">Saxena &amp; Singh (1998)</a>
Rat	CA	Bone-marrow cells. Comparison in liver cells before and after partial hepatectomy of treated rats	<a href="#">Rossi et al. (1987)</a>
Rat	CA SCE	Bone-marrow cells. Regenerating hepatocytes (SCE)	<a href="#">Masuda et al. (1990)</a>
Rat	MN	Peripheral blood reticulocytes and bone-marrow cells comparison	<a href="#">Hayashi et al. (1992)</a>
Mouse	MN	Bone-marrow PCE (positive), hepatocytes (negative)	<a href="#">Parton &amp; Garriott (1997)</a>
Rat	MN	Bone-marrow cells and peripheral blood reticulocytes. 14 rat strains compared	<a href="#">Hamada et al. (2001)</a>
Rat	MN	Bone-marrow cells and peripheral blood reticulocytes. Effect of ageing studied	<a href="#">Hamada et al. (2003)</a>
Rat	MN	Pre-estrous vaginal cells	<a href="#">Zúñiga-González et al. (2003)</a>

Table 4.1 (continued)

Species	Cytogenetic end-point investigated	Additional considerations	Reference
Rat	CA MN	Bone-marrow cells. Simultaneous evaluation of two end-points in the same animal	<a href="#">Krishna et al. (1991)</a>
Rat	MN	Bone-marrow, spleen, peripheral blood cells	<a href="#">Abramsson-Zetterberg et al. (1999)</a>
Rat	MN	Embryos, treatment during pre-implantation period	<a href="#">Giavini et al. (1990)</a>
Rat	MN CA	Bone-marrow and spleen cells. In-vivo/in-vitro assay	<a href="#">Moore et al. (1995)</a>
Newt	MN	Larvae exposed to agent. Red blood cells	<a href="#">Fernandez et al. (1989)</a>
Mouse, Chinese hamster	CA SCE	Bone-marrow cells Comparison of different routes of administration	<a href="#">Jenderny et al. (1988)</a>
Rat, mouse	MN SCE	Bone-marrow cells (MN). Splenocytes (SCE). Rats more susceptible than mice	<a href="#">Simula &amp; Priestly (1992)</a>
Rat, mouse, Chinese hamster	MN SCE	Bone-marrow cells. Species comparison Susceptibility ranked into the order rat > mouse > Chinese hamster	<a href="#">Madle et al. (1986)</a>
Mouse, rat, Chinese hamster, Armenian hamster, guinea-pig	CA	Bone-marrow cells. Interspecies comparison Susceptibility ranked into the order guinea-pig > rat > mouse > Chinese hamster > Armenian hamster	<a href="#">Nersessian et al. (1992)</a>
Dog (beagle)	MN	Peripheral blood reticulocytes and bone-marrow cells comparison	<a href="#">Harper et al. (2007)</a>
Monkey	MN	Peripheral blood reticulocytes and bone-marrow cells comparison	<a href="#">Hotchkiss et al. (2008)</a>
Marmoset	MN	Peripheral blood erythrocytes	<a href="#">Zúñiga-González et al. (2005)</a>

CA, chromosomal aberrations; i.p., intraperitoneal; MN, micronuclei; PCE, polychromatic erythrocytes; p.o., per oral; SCE, sister chromatid exchange; vs, versus

Table 4.2 Studies of modulation of cyclophosphamide genotoxicity *in vivo* and *in vitro*

Agent	Experimental system	End-point measured	Effect	Reference
Retinol	CHEL cells <i>in vitro</i>	SCE	Inhibitory	<a href="#">Cozzi et al. (1990)</a>
Retinoic acid				
Apigenin	Human lymphocytes + S9 <i>in vitro</i>	SCE CA	Inhibitory	<a href="#">Siddique et al. (2008)</a>
$\beta$ -carotene	Human lymphocytes + S9 <i>in vitro</i>	SCE	Inhibitory	<a href="#">Edenharder et al. (1998)</a>
Retinal				
$\alpha$ -tocopherol				
Riboflavin				
Vitamin C	Human lymphocytes <i>in vitro</i>	SCE	Enhancing	<a href="#">Edenharder et al. (1998)</a>
Vitamin K <sub>1</sub>	Human lymphocytes <i>in vitro</i>	SCE	Inhibitory or enhancing (dependent on timing)	<a href="#">Edenharder et al. (1998)</a>
Melatonin	CHO cells + S9 <i>in vitro</i>	SCE CA	Inhibitory	<a href="#">De Salvia et al. (1999)</a>
Melatonin	CHO cells + S9 <i>in vitro</i>	Comet formation (DNA damage)	Inhibitory	<a href="#">Musatov et al. (1998)</a>
O <sup>6</sup> -alkylguanine-DNA alkyltransferase (AGT)	CHO cells <i>in vitro</i>	<i>Hprt</i> mutation	Inhibitory	<a href="#">Cai et al. (1999)</a>
Buthionine sulfoximine	V79 cells and CHO +S9 <i>in vitro</i>	SCE	Enhancing	<a href="#">Köberle &amp; Speit (1990)</a>
Prostaglandin E <sub>2</sub>	Mouse lymphoid L1210 leukaemia cells <i>in vivo</i>	SCE	Enhancing	<a href="#">Mourdelatos et al. (1995)</a>
Garlic extract	Swiss albino mice <i>in vivo</i>	CA (bone-marrow cells)	Inhibitory	<a href="#">Shukla &amp; Taneja (2002)</a>
Indole-3-carbinol	Swiss albino mice <i>in vivo</i>	CA (bone-marrow cells)	Inhibitory	<a href="#">Shukla et al. (2004)</a>
Ascorbic acid	Pregnant CBA/CaH mice <i>in vivo</i>	CA SCE (pre-implantation embryos)	Inhibitory (SCE no effect)	<a href="#">Kola et al. (1989)</a>
Ascorbic acid	Pregnant NMRI Kisslegg mice <i>in vivo</i>	CA SCE (pre-implantation embryos)	Inhibitory (SCE no effect)	<a href="#">Vogel &amp; Spielmann (1989)</a>
$\beta$ -glucan	Male CD-1 mice <i>in vivo</i>	CA (bone-marrow and spermatogonial cells)	Inhibitory	<a href="#">Tohamy et al. (2003)</a>

Table 4.2 (continued)

Agent	Experimental system	End-point measured	Effect	Reference
Nafenopin	Male Wistar rats <i>in vivo</i>	CA MN	Enhancing CA in bone marrow and MN in hepatocytes. Inhibitory on MN in bone marrow	<a href="#">Voskoboinik et al. (1997)</a>
Prostaglandin E <sub>2</sub>	BALB/c mice inoculated with Ehrlich ascites tumour cells <i>in vivo</i>	SCE (Ehrlich ascites tumour cells)	Inhibitory	<a href="#">Mourelatos et al. (1993)</a>
Ginsenoside Rh <sub>2</sub>	Male C57BL/6 mice <i>in vivo</i>	MN (bone-marrow cells) Comet formation (DNA damage) (white blood cells)	Inhibitory	<a href="#">Wang et al. (2006)</a>
Verapamil	Male BALB/c and C57BL/6 mice <i>in vivo</i>	CA (bone-marrow cells)	Enhancing	<a href="#">Nesterova et al. (1999)</a>
Citrus extract	Male BALB/c mice <i>in vivo</i>	MN (bone-marrow cells)	Inhibitory	<a href="#">Hosseinimehr &amp; Karami (2005a)</a>
Captopril	Male NMRI mice <i>in vivo</i>	MN (bone-marrow cells)	Inhibitory	<a href="#">Hosseinimehr &amp; Karami (2005b)</a>
<i>Spirulina fusiformis</i>	Male Swiss albino mice <i>in vivo</i>	MN (bone-marrow cells)	Inhibitory	<a href="#">Premkumar et al. (2001a)</a>
Saffron ( <i>Crocus sativus</i> L.)	Male Swiss albino mice <i>in vivo</i>	MN (bone-marrow cells)	Inhibitory	<a href="#">Premkumar et al. (2001b)</a>
Melatonin and its derivatives	Male albino mice <i>in vivo</i>	MN (bone-marrow cells)	Inhibitory	<a href="#">Elmegeed et al. (2008)</a>
Vitamin C	Male Swiss albino mice <i>in vivo</i>	MN (bone-marrow cells)	Inhibitory	<a href="#">Ghaskadbi et al. (1992)</a>
Malaria infection	Female C57BL/6 mice	MN (bone-marrow cells)	Inhibitory	<a href="#">Poça et al. (2008)</a>
Lipoic acid	Male Wistar rats <i>in vivo</i>	MN (bone-marrow cells and peripheral blood cells)	Inhibitory	<a href="#">Selvakumar et al. (2006)</a>
Folic acid	Newborn Wistar rats (fetal exposure) <i>in vivo</i>	MN (peripheral blood erythrocytes)	Inhibitory	<a href="#">Gómez-Meda et al. (2004)</a>
<i>Taenia taeniformis</i> infection	Sprague-Dawley rats	MN (peripheral blood erythrocytes)	Enhancing	<a href="#">Montero et al. (2003)</a>
O <sup>6</sup> -methylguanine-DNA methyltransferase	C57BL/6 wild type and <i>Mgmt</i> <sup>-/-</sup> mice	<i>Hprt</i> mutation (splenic lymphocytes)	Inhibitory (non-significant)	<a href="#">Hansen et al. (2007)</a>

CA, chromosomal aberrations; CHEL, Chinese hamster epithelial liver; CHO, Chinese hamster ovary; *Hprt*, hypoxanthine(guanine)phosphoribosyl transferase; MN, micronuclei; SCE, sister chromatid exchange

of cyclophosphamide is the phosphoramidate mustard (Povirk & Shuker, 1994). This metabolite is also generally considered to be the most genotoxic, but the contribution of acrolein, which is highly toxic, to the genotoxic activity of cyclophosphamide is less clear.

It is well established that the treatment of cancer patients with cyclophosphamide results in inflammation of the urinary bladder (haemorrhagic cystitis), which is not seen with other alkylating agents (Forni *et al.*, 1964; Liedberg *et al.*, 1970). In rats, cyclophosphamide treatment resulted in cystitis as well (Crocitto *et al.*, 1996), and in mice, mutagenic activity has been detected in urine following cyclophosphamide treatment (Teet *et al.*, 1997). The ultimate alkylating metabolite of cyclophosphamide, phosphoramidate mustard, is metabolized but was not shown to cause cytotoxicity and had minimal morphological effects on the mouse bladder, but an intermediate in the formation of the acrolein metabolite, diethylcyclophosphamide administered by intraperitoneal injection, caused severe cystitis in male rats, and less extensive inflammation in female rats (Cox, 1979). Acrolein administered to rats by intraperitoneal injections increased urothelial cell proliferation (Sakata *et al.*, 1989). Acrolein is the only metabolite of cyclophosphamide that is known to be both reactive and cytotoxic (IARC, 1995). Collectively, these data indicate that acrolein is the likely causative agent in cyclophosphamide-induced cystitis. Cystitis is an established condition associated with the development of both squamous cell and urothelial bladder cancers (Michaud, 2007). However, intraperitoneal injections of acrolein by itself only induced bladder hyperplasia, not cancer (Cohen *et al.*, 1992), and oral administration studies in mice and rats did not result in carcinogenic effects (IARC, 1995). Thus it is plausible that acrolein-induced cystitis plays a promoting role in cyclophosphamide bladder tumorigenesis that is initiated by other cyclophosphamide metabolites.

The protective effect of O<sup>6</sup>-alkylguanine-DNA alkyltransferase (AGT) against cyclophosphamide mutagenicity (*Hprt* mutations) (Cai *et al.*, 1999), and cytotoxicity (Friedman *et al.*, 1999) in CHO cells implies some involvement of acrolein-derived DNA damage. However, mice deficient in this protein (called O<sup>6</sup>-methylguanine-DNA methyl transferase [MGMT] in this study) were less susceptible to cyclophosphamide tumorigenesis, not more (Nagasubramanian *et al.*, 2008). Studies of sister chromatid exchange induced in human lymphocytes by acrolein and phosphoramidate mustard suggest that phosphoramidate mustard is the more potent genotoxic agent (Wilmer *et al.*, 1990). Furthermore, analysis of *TP53* mutations in cyclophosphamide-associated human bladder cancers suggests that the mutations detected are characteristic of DNA damage caused by phosphoramidate mustard, rather than by acrolein (Khan *et al.*, 1998).

## 4.4 Synthesis

Cyclophosphamide, after its bioactivation to alkylating metabolites, is carcinogenic via a genotoxic mechanism.

## 5. Evaluation

There is *sufficient evidence* in humans for the carcinogenicity of cyclophosphamide. Cyclophosphamide causes cancer of the bladder, and acute myeloid leukaemia.

There is *sufficient evidence* in experimental animals for the carcinogenicity of cyclophosphamide.

Cyclophosphamide is *carcinogenic to humans* (Group 1).

## References

- Abramsson-Zetterberg L, Grawé J, Zetterberg G (1999). The micronucleus test in rat erythrocytes from bone marrow, spleen and peripheral blood: the response to low doses of ionizing radiation, cyclophosphamide and vincristine determined by flow cytometry. *Mutat Res*, 423: 113–124. PMID:10029688
- Anderson D, Bishop JB, Garner RC *et al.* (1995). Cyclophosphamide: review of its mutagenicity for an assessment of potential germ cell risks. *Mutat Res*, 330: 115–181. PMID:7623863
- Andersson BS, Sadeghi T, Siciliano MJ *et al.* (1996). Nucleotide excision repair genes as determinants of cellular sensitivity to cyclophosphamide analogs. *Cancer Chemother Pharmacol*, 38: 406–416. doi:10.1007/s002800050504 PMID:8765433
- Andersson M, Agurell E, Vaghef H *et al.* (2003). Extended-term cultures of human T-lymphocytes and the comet assay: a useful combination when testing for genotoxicity in vitro? *Mutat Res*, 540: 43–55. PMID:12972057
- Balu N, Gamcsik MP, Colvin ME *et al.* (2002). Modified guanines representing O<sup>6</sup>-alkylation by the cyclophosphamide metabolites acrolein and chloroacetaldehyde: synthesis, stability, and ab initio studies. *Chem Res Toxicol*, 15: 380–387. doi:10.1021/tx0101503 PMID:11896686
- Barton TS, Wyrobek AJ, Hill FS *et al.* (2003). Numerical chromosomal abnormalities in rat epididymal spermatozoa following chronic cyclophosphamide exposure. *Biol Reprod*, 69: 1150–1157. doi:10.1095/biol-reprod.103.016261 PMID:12773405
- Batiste-Alentorn M, Xamena N, Creus A, Marcos R (1994). Further studies with the somatic white-ivory system of *Drosophila melanogaster*: genotoxicity testing of ten carcinogens. *Environ Mol Mutagen*, 24: 143–147. doi:10.1002/em.2850240210 PMID:7925328
- Benning V, Depasse F, Melcion C, Cordier A (1992). Detection of micronuclei after exposure to mitomycin C, cyclophosphamide and diethylnitrosamine by the in vivo micronucleus test in mouse splenocytes. *Mutat Res*, 280: 137–142. doi:10.1016/0165-1218(92)90009-O PMID:1378538
- Benson AJ, Martin CN, Garner RC (1988). N-(2-hydroxyethyl)-N-[2-(7-guaninyl)ethyl]amine, the putative major DNA adduct of cyclophosphamide in vitro and in vivo in the rat. *Biochem Pharmacol*, 37: 2979–2985. doi:10.1016/0006-2952(88)90285-7 PMID:3395373
- Black SM, Ellard S, Meehan RR *et al.* (1989). The expression of cytochrome P450IIB1 in *Saccharomyces cerevisiae* results in an increased mutation frequency when exposed to cyclophosphamide. *Carcinogenesis*, 10: 2139–2143. doi:10.1093/carcin/10.11.2139 PMID:2680147
- Branda RF, Brooks EM, Chen Z *et al.* (2002). Dietary modulation of mitochondrial DNA deletions and copy number after chemotherapy in rats. *Mutat Res*, 501: 29–36. PMID:11934435
- Burgaz S, Karahalil B, Bayrak P *et al.* (1999). Urinary cyclophosphamide excretion and micronuclei frequencies in peripheral lymphocytes and in exfoliated buccal epithelial cells of nurses handling antineoplastics. *Mutat Res*, 439: 97–104. PMID:10029685
- Burgaz S, Karahalil B, Canli Z *et al.* (2002). Assessment of genotoxic damage in nurses occupationally exposed to antineoplastics by the analysis of chromosomal aberrations. *Hum Exp Toxicol*, 21: 129–135. doi:10.1191/0960327102ht230oa PMID:12102538
- Cai Y, Wu MH, Ludeman SM *et al.* (1999). Role of O<sup>6</sup>-alkylguanine-DNA alkyltransferase in protecting against cyclophosphamide-induced toxicity and mutagenicity. *Cancer Res*, 59: 3059–3063. PMID:10397244
- Cavallo D, Ursini CL, Perniconi B *et al.* (2005). Evaluation of genotoxic effects induced by exposure to antineoplastic drugs in lymphocytes and exfoliated buccal cells of oncology nurses and pharmacy employees. *Mutat Res*, 587: 45–51. PMID:16202645
- Chung FL, Young R, Hecht SS (1984). Formation of cyclic 1-N<sup>2</sup>-propanodeoxyguanosine adducts in DNA upon reaction with acrolein or crotonaldehyde. *Cancer Res*, 44: 990–995. PMID:6318992
- Cohen SM, Garland EM, St John M *et al.* (1992). Acrolein initiates rat urinary bladder carcinogenesis. *Cancer Res*, 52: 3577–3581. PMID:1617627
- Cox PJ (1979). Cyclophosphamide cystitis—identification of acrolein as the causative agent. *Biochem Pharmacol*, 28: 2045–2049. doi:10.1016/0006-2952(79)90222-3 PMID:475846
- Cozzi R, Bona R, Polani S, De Salvia R (1990). Retinoids as modulators of metabolism: their inhibitory effect on cyclophosphamide and 7,12-dimethylbenz[a]anthracene induced sister chromatid exchanges in a metabolically competent cell line. *Mutagenesis*, 5: 397–401. doi:10.1093/mutage/5.4.397 PMID:2118976
- Crocitto LE, Simpson JF, Wilson TG (1996). Bladder augmentation in the prevention of cyclophosphamide-induced haemorrhagic cystitis in the rat model. *Br J Urol*, 78: 530–533. doi:10.1046/j.1464-410X.1996.01146.x PMID:8944508
- Curtis RE, Boice JD Jr, Stovall M *et al.* (1992). Risk of leukemia after chemotherapy and radiation treatment for breast cancer. *N Engl J Med*, 326: 1745–1751. doi:10.1056/NEJM199206253262605 PMID:1594016
- Cushnir JR, Naylor S, Lamb JH *et al.* (1990). Identification of phosphoramidate/DNA adducts using tandem mass spectrometry. *Rapid Commun Mass Spectrom*, 4: 410–414. doi:10.1002/rcm.1290041014 PMID:2134189
- Cusidó L, Pujol R, Egozcue J, García M (1995). Cyclophosphamide-induced synaptonemal complex

- damage during meiotic prophase of female *Rattus norvegicus*. *Mutat Res*, 329: 131–141. PMID:7603495
- Darroudi F & Natarajan AT (1993). Metabolic activation of chemicals to mutagenic carcinogens by human hepatoma microsomal extracts in Chinese hamster ovary cells (in vitro). *Mutagenesis*, 8: 11–15. doi:10.1093/mutage/8.1.11 PMID:8383795
- De Salvia R, Fiore M, Aglitti T *et al.* (1999). Inhibitory action of melatonin on H<sub>2</sub>O<sub>2</sub>- and cyclophosphamide-induced DNA damage. *Mutagenesis*, 14: 107–112. doi:10.1093/mutage/14.1.107 PMID:10474831
- Director AE, Tucker JD, Ramsey MJ, Nath J (1998). Chronic ingestion of clastogens by mice and the frequency of chromosome aberrations. *Environ Mol Mutagen*, 32: 139–147. doi:10.1002/(SICI)1098-2280(1998)32:2<139::AID-EM9>3.0.CO;2-O PMID:9776176
- Dobrzyńska MM, Lenarczyk M, Gajewski AK (1998). Induction of dominant lethal mutations after exposure of male mice to cyclophosphamide. *Rocz Panstw Zakl Hig*, 49: 285–291. PMID:9930021
- Doehmer J, Seidel A, Oesch F, Glatt HR (1990). Genetically engineered V79 Chinese hamster cells metabolically activate the cytostatic drugs cyclophosphamide and ifosfamide. *Environ Health Perspect*, 88: 63–65. doi:10.2307/3431052 PMID:2272335
- Doehmer J, Wölfel C, Dogra S *et al.* (1992). Applications of stable V79-derived cell lines expressing rat cytochromes P4501A1, 1A2, and 2B1. *Xenobiotica*, 22: 1093–1099. doi:10.3109/00498259209051863 PMID:1441600
- Eastin WC, Mennear JH, Tennant RW *et al.* (2001). Tg.AC genetically altered mouse: assay working group overview of available data. *Toxicol Pathol*, 29: Suppl60–80. doi:10.1080/019262301753178483 PMID:11695563
- Edenharder R, Kerkhoff G, Dunkelberg H (1998). Effects of beta-carotene, retinal, riboflavin, alpha-tocopherol and vitamins C and K1 on sister-chromatid exchanges induced by 3-amino-1-methyl-5H-pyrido[4,3-b]indole (Trp-P-2) and cyclophosphamide in human lymphocyte cultures. *Food Chem Toxicol*, 36: 897–906. doi:10.1016/S0278-6915(98)00068-4 PMID:9771550
- Ellard S, Mohammed Y, Dogra S *et al.* (1991). The use of genetically engineered V79 Chinese hamster cultures expressing rat liver *CYP1A1*, *1A2* and *2B1* cDNAs in micronucleus assays. *Mutagenesis*, 6: 461–470. doi:10.1093/mutage/6.6.461 PMID:1800893
- Elmegeed GA, Khalil WK, Raouf AA, Abdelhalim MM (2008). Synthesis and in vivo anti-mutagenic activity of novel melatonin derivatives. *Eur J Med Chem*, 43: 763–770. doi:10.1016/j.ejmech.2007.06.003 PMID:17706326
- Ember I & Kiss I (1997). In vivo effects of cyclophosphamide on oncogene and suppressor gene expression in a “follow up” study. *Anticancer Res*, 17: 3593–3597. PMID:9413208
- Ember I, Raposa T, Varga C, Kiss I (1995). Effect of different cytostatic protocols on oncogene expression in CBA/Ca mice. *Anticancer Res*, 15: 1285–1288. PMID:7654010
- Fernandez M, Gauthier L, Jaylet A (1989). Use of new larvae for in vivo genotoxicity testing of water: results on 19 compounds evaluated by the micronucleus test. *Mutagenesis*, 4: 17–26. doi:10.1093/mutage/4.1.17 PMID:2654548
- Foiles PG, Akerkar SA, Miglietta LM, Chung F-L (1990). Formation of cyclic deoxyguanosine adducts in Chinese hamster ovary cells by acrolein and crotonaldehyde. *Carcinogenesis*, 11: 2059–2061. doi:10.1093/carcin/11.11.2059 PMID:2225341
- Forni AM, Koss LG, Geller W (1964). Cytological study of the effect of cyclophosphamide on the epithelium of the urinary bladder in man. *Cancer*, 17: 1348–1355. doi:10.1002/1097-0142(196410)17:10<1348::AID-CNCR2820171017>3.0.CO;2-0 PMID:14236768
- Frenzilli G, Bosco E, Barale R (2000). Validation of single cell gel assay in human leukocytes with 18 reference compounds. *Mutat Res*, 468: 93–108. PMID:10882888
- Friedman HS, Pegg AE, Johnson SP *et al.* (1999). Modulation of cyclophosphamide activity by O<sup>6</sup>-alkylguanine-DNA alkyltransferase. *Cancer Chemother Pharmacol*, 43: 80–85. doi:10.1007/s002800050866 PMID:9923545
- Ghaskadbi S, Rajmachikar S, Agate C *et al.* (1992). Modulation of cyclophosphamide mutagenicity by vitamin C in the in vivo rodent micronucleus assay. *Teratog Carcinog Mutagen*, 12: 11–17. doi:10.1002/tcm.1770120103 PMID:1354896
- Giavini E, Lemonica IP, Lou Y *et al.* (1990). Induction of micronuclei and toxic effects in embryos of pregnant rats treated before implantation with anticancer drugs: cyclophosphamide, cis-platinum, adriamycin. *Teratog Carcinog Mutagen*, 10: 417–426. doi:10.1002/tcm.1770100507 PMID:1981952
- Gómez-Meda BC, Zúñiga-González GM, Zamora-Perez A *et al.* (2004). Folate supplementation of cyclophosphamide-treated mothers diminishes micronucleated erythrocytes in peripheral blood of newborn rats. *Environ Mol Mutagen*, 44: 174–178. doi:10.1002/em.20037 PMID:15278921
- Gorelick NJ, Andrews JL, deBoer JG *et al.* (1999). Tissue-specific mutant frequencies and mutational spectra in cyclophosphamide-treated *lacI* transgenic mice. *Environ Mol Mutagen*, 34: 154–166. doi:10.1002/(SICI)1098-2280(1999)34:2/3<154::AID-EM15>3.0.CO;2-0 PMID:10529740
- Greene MH, Harris EL, Gershenson DM *et al.* (1986). Melphalan may be a more potent leukemogen than cyclophosphamide. *Ann Intern Med*, 105: 360–367. PMID:3740675
- Haas JF, Kittelmann B, Mehnert WH *et al.* (1987). Risk of leukaemia in ovarian tumour and breast cancer patients following treatment by cyclophosphamide. *Br J Cancer*, 55: 213–218. PMID:3814491

- Habs MR & Schmähld (1983). Prevention of urinary bladder tumors in cyclophosphamide-treated rats by additional medication with the uroprotectors sodium 2-mercaptoethane sulfonate (mesna) and disodium 2,2'-dithiobis-ethane sulfonate (dimesna). *Cancer*, 51: 606–609. doi:10.1002/1097-0142(19830215)51:4<606::AID-CNCR2820510409>3.0.CO;2-S PMID:6401591
- Hamada S, Nakajima K, Serikawa T, Hayashi M (2003). The effect of aging on the results of the rat micronucleus assay. *Mutagenesis*, 18: 273–275. doi:10.1093/mutage/18.3.273 PMID:12714693
- Hamada S, Yamasaki KI, Nakanishi S *et al.* (2001). Evaluation of the general suitability of the rat for the micronucleus assay: the effect of cyclophosphamide in 14 strains. *Mutat Res*, 495: 127–134. PMID:11448650
- Hansen RJ, Nagasubramanian R, Delaney SM *et al.* (2007). Role of O<sup>6</sup>-methylguanine-DNA methyltransferase in protecting from alkylating agent-induced toxicity and mutations in mice. *Carcinogenesis*, 28: 1111–1116. doi:10.1093/carcin/bgl218 PMID:17116724
- Harper BL, Ramanujam VM, Legator MS (1989). Micronucleus formation by benzene, cyclophosphamide, benzo(a)pyrene, and benzidine in male, female, pregnant female, and fetal mice. *Teratog Carcinog Mutagen*, 9: 239–252. doi:10.1002/tcm.1770090406 PMID:2572067
- Harper SB, Dertinger SD, Bishop ME *et al.* (2007). Flow cytometric analysis of micronuclei in peripheral blood reticulocytes III. An efficient method of monitoring chromosomal damage in the beagle dog. *Toxicol Sci*, 100: 406–414. doi:10.1093/toxsci/kfm241 PMID:17872896
- Hartmann A, Herkommer K, Glück M, Speit G (1995). DNA-damaging effect of cyclophosphamide on human blood cells in vivo and in vitro studied with the single-cell gel test (comet assay). *Environ Mol Mutagen*, 25: 180–187. doi:10.1002/em.2850250303 PMID:7737135
- Hartmann A & Speit G (1995). Genotoxic effects of chemicals in the single cell gel (SCG) test with human blood cells in relation to the induction of sister-chromatid exchanges (SCE). *Mutat Res*, 346: 49–56. doi:10.1016/0165-7992(95)90068-3 PMID:7530329
- Hatanaka Y, Kitagawa Y, Toyoda Y *et al.* (1992). Micronucleus test with cyclophosphamide using mouse peripheral blood reticulocytes. *Mutat Res*, 278: 99–101. doi:10.1016/0165-1218(92)90216-M PMID:1372710
- Hayashi M, Kodama Y, Awogi T *et al.* (1992). The micronucleus assay using peripheral blood reticulocytes from mitomycin C- and cyclophosphamide-treated rats. *Mutat Res*, 278: 209–213. doi:10.1016/0165-1218(92)90236-S PMID:1372708
- Hemminki K (1987). DNA-binding products of normitrogen mustard, a metabolite of cyclophosphamide. *Chem Biol Interact*, 61: 75–88. doi:10.1016/0009-2797(87)90020-2 PMID:3815587
- Hoorn AJ, Custer LL, Myhr BC *et al.* (1993). Detection of chemical mutagens using Muta<sup>+</sup> Mouse: a transgenic mouse model. *Mutagenesis*, 8: 7–10. doi:10.1093/mutage/8.1.7 PMID:8450770
- Hosseinimehr SJ & Karami M (2005a). Citrus extract modulates genotoxicity induced by cyclophosphamide in mice bone marrow cells. *J Pharm Pharmacol*, 57: 505–509. doi:10.1211/0022357055849 PMID:15831212
- Hosseinimehr SJ & Karami M (2005b). Chemoprotective effects of captopril against cyclophosphamide-induced genotoxicity in mouse bone marrow cells. *Arch Toxicol*, 79: 482–486. doi:10.1007/s00204-005-0655-7 PMID:15856182
- Hotchkiss CE, Bishop ME, Dertinger SD *et al.* (2008). Flow cytometric analysis of micronuclei in peripheral blood reticulocytes IV: an index of chromosomal damage in the rhesus monkey (*Macaca mulatta*). *Toxicol Sci*, 102: 352–358. doi:10.1093/toxsci/kfn013 PMID:18211907
- Hoyes KP, Wadeson PJ, Sharma HL *et al.* (1998). Mutation studies in *lacI* transgenic mice after exposure to radiation or cyclophosphamide. *Mutagenesis*, 13: 607–612. doi:10.1093/mutage/13.6.607 PMID:9862192
- Huang CC, Tan JC, Sirianni SR *et al.* (1990). Comparison of baseline sister-chromatid exchanges (SCE), cyclophosphamide-, ethylnitrosourea (ENU)-induced SCE, ENU-induced cell-cycle delay and chromosome aberrations between Peru and laboratory mice. *Mutat Res*, 230: 93–100. PMID:2342502
- IARC (1981). Some antineoplastic and immunosuppressive agents. *IARC Monogr Eval Carcinog Risk Chem Hum*, 26: 1–411. PMID:6944253
- IARC (1987a). Overall evaluations of carcinogenicity: an updating of IARC Monographs volumes 1 to 42. *IARC Monogr Eval Carcinog Risks Hum Suppl*, 7: 1–440. PMID:3482203
- IARC (1987b). Genetic and related effects: An updating of selected IARC monographs from Volumes 1 to 42. *IARC Monogr Eval Carcinog Risks Hum Suppl*, 6: 1–729. PMID:3504843
- IARC (1995). Dry cleaning, some chlorinated solvents and other industrial chemicals. *IARC Monogr Eval Carcinog Risks Hum*, 63: 1–551.
- Jenderny J, Walk RA, Hackenberg U, Röhrborn G (1988). Chromosomal abnormalities and sister-chromatid exchange in bone marrow cells of mice and Chinese hamsters after inhalation and intraperitoneal administration. II. Cyclophosphamide. *Mutat Res*, 203: 1–10. PMID:3340088
- Kaldor JM, Day NE, Kittelmann B *et al.* (1995). Bladder tumours following chemotherapy and radiotherapy for ovarian cancer: a case-control study. *Int J Cancer*, 63: 1–6. doi:10.1002/ijc.2910630102 PMID:7558434
- Kaldor JM, Day NE, Pettersson F *et al.* (1990). Leukemia following chemotherapy for ovarian cancer. *N Engl J Med*, 322: 1–6. doi:10.1056/NEJM199001043220101 PMID:2104664
- Kalweit S, Utesch D, von der Hude W, Madle S (1999). Chemically induced micronucleus formation in V79

- cells—comparison of three different test approaches. *Mutat Res*, 439: 183–190. PMID:10023054
- Kelly WA, Nelson LW, Hawkins HC, Weikel JH Jr (1974). An evaluation of the tumorigenicity of cyclophosphamide and urethan in newborn mice. *Toxicol Appl Pharmacol*, 27: 629–640. doi:10.1016/0041-008X(74)90042-8 PMID:4850539
- Khan MA, Travis LB, Lynch CF *et al.* (1998). *p53* mutations in cyclophosphamide-associated bladder cancer. *Cancer Epidemiol Biomarkers Prev*, 7: 397–403. PMID:9610789
- Kikuno T, Honma M, Ogura S *et al.* (1995). DNA fingerprint analysis in chemically mutagenized Chinese hamster lung cells. *Mutat Res*, 338: 87–93. PMID:7565885
- Kinlen LJ (1985). Incidence of cancer in rheumatoid arthritis and other disorders after immunosuppressive treatment. *Am J Med*, 78: 1A44–49. doi:10.1016/0002-9343(85)90245-1 PMID:3970040
- Kirsch-Volders M, Sofuni T, Aardema M *et al.* (2003). Report from the in vitro micronucleus assay working group. *Mutat Res*, 540: 153–163. PMID:14550499
- Köberle B & Speit G (1990). The effect of glutathione depletion on sister-chromatid exchange induction by cytostatic drugs. *Mutat Res*, 243: 225–231. doi:10.1016/0165-7992(90)90095-2 PMID:2308598
- Kola I, Vogel R, Spielmann H (1989). Co-administration of ascorbic acid with cyclophosphamide (CPA) to pregnant mice inhibits the clastogenic activity of CPA in preimplantation murine blastocysts. *Mutagenesis*, 4: 297–301. doi:10.1093/mutage/4.4.297 PMID:2674608
- Konopacka M (1994). Evaluation of frequency of micronuclei in exfoliated cells from bladder of mice treated with benzo(a) pyrene, 2-acetylaminofluorene and cyclophosphamide. *Cell Biol Int*, 18: 669–672. doi:10.1006/cbir.1994.1094 PMID:8075628
- Krishna G, Kropko ML, Ciaravino V, Theiss JC (1991). Simultaneous micronucleus and chromosome aberration assessment in the rat. *Mutat Res*, 264: 29–35. doi:10.1016/0165-7992(91)90042-3 PMID:1881414
- Krishna G, Nath J, Petersen M, Ong T (1987). Cyclophosphamide-induced cytogenetic effects in mouse bone marrow and spleen cells in vivo and in vivo/in vitro assays. *Teratog Carcinog Mutagen*, 7: 183–195. doi:10.1002/tcm.1770070209 PMID:2885941
- Krishna G, Nath J, Petersen M, Ong T (1988). In vivo and in vivo/in vitro kinetics of cyclophosphamide-induced sister-chromatid exchanges in mouse bone marrow and spleen cells. *Mutat Res*, 204: 297–305. doi:10.1016/0165-1218(88)90103-6 PMID:3343979
- Kugler U, Bauchinger M, Schmid E, Göggelmann W (1987). The effectiveness of S9 and microsomal mix on activation of cyclophosphamide to induce genotoxicity in human lymphocytes. *Mutat Res*, 187: 151–156. doi:10.1016/0165-1218(87)90082-6 PMID:3821768
- Kulka U, Doehmer J, Glatt HR, Bauchinger M (1993). Cytogenetic effects of promutagens in genetically engineered V79 Chinese hamster cells expressing cytochromes P450. *Eur J Pharmacol*, 228: 299–304. PMID:8482321
- Kwanyuen P, Erexson GL, Bryant MF, Kligerman AD (1990). Comparison of sister-chromatid exchange frequencies in mouse T- and B-lymphocytes after exposure to 4-hydroxycyclophosphamide or phosphoramid mustard. *Mutat Res*, 245: 293–297. doi:10.1016/0165-7992(90)90159-H PMID:2266981
- Liedberg CF, Rausing A, Langeland P (1970). Cyclophosphamide hemorrhagic cystitis. *Scand J Urol Nephrol*, 4: 183–190. doi:10.3109/00365597009137596 PMID:5518247
- Maccubbin AE, Caballes L, Riordan JM *et al.* (1991). A cyclophosphamide/DNA phosphoester adduct formed in vitro and in vivo. *Cancer Res*, 51: 886–892. PMID:1988129
- Madle E, Korte A, Beek B (1986). Species differences in mutagenicity testing. II. Sister-chromatid exchange and micronucleus induction in rats, mice and Chinese hamsters treated with cyclophosphamide. *Mutagenesis*, 1: 419–422. doi:10.1093/mutage/1.6.419 PMID:3331680
- Mahgoub N, Taylor BR, Le Beau MM *et al.* (1999). Myeloid malignancies induced by alkylating agents in Nf1 mice. *Blood*, 93: 3617–3623. PMID:10339466
- Masuda R, Abe S, Yoshida MC *et al.* (1990). Cytochrome P-450 and chromosome damage by cyclophosphamide in LEC strain rats predisposed to hereditary hepatitis and liver cancer. *Mutat Res*, 244: 309–316. doi:10.1016/0165-7992(90)90078-X PMID:2385246
- McCarroll N, Keshava N, Cimino M *et al.* (2008). An evaluation of the mode of action framework for mutagenic carcinogens case study: Cyclophosphamide. *Environ Mol Mutagen*, 49: 117–131. doi:10.1002/em.20372 PMID:18240158
- McCarvill JT, Lubet RA, Schechtman LM *et al.* (1990). Morphological transformation of BALB/3T3 cells by various procarcinogens in the presence of a rat liver S-9 activation system. *Environ Mol Mutagen*, 16: 304–310. doi:10.1002/em.2850160410 PMID:2253607
- McClain RM, Keller D, Casciano D *et al.* (2001). Neonatal mouse model: review of methods and results. *Toxicol Pathol*, 29: Suppl128–137. doi:10.1080/019262301753178537 PMID:11695548
- McDiarmid MA, Iype PT, Kolodner K *et al.* (1991). Evidence for acrolein-modified DNA in peripheral blood leukocytes of cancer patients treated with cyclophosphamide. *Mutat Res*, 248: 93–99. PMID:2030715
- McDiarmid MA, Strickland PT, Kolodner K *et al.* (1990). Baseline and phosphoramid mustard-induced sister-chromatid exchanges in cancer patients treated with cyclophosphamide. *Mutat Res*, 241: 273–278. doi:10.1016/0165-1218(90)90024-V PMID:2366806
- McEvoy GK, editor (2007). *American Hospital Formulary Service Drug Information*. Bethesda, MD: American Society of Health-System Pharmacists, pp. 984–988.

- Mertens R, Rubbert F, Büssing A (1995). Childhood acute lymphoblastic leukemia (ALL): sister chromatid exchange (SCE) frequency and lymphocyte subpopulations during therapy. *Leukemia*, 9: 501–505. PMID:7885047
- Michaud DS (2007). Chronic inflammation and bladder cancer. *Urol Oncol*, 25: 260–268. PMID:17483025
- Miller K (1991a). Sister-chromatid exchange in human B- and T-lymphocytes exposed to bleomycin, cyclophosphamide, and ethyl methanesulfonate. *Mutat Res*, 247: 175–182. PMID:1706068
- Miller K (1991b). Clastogenic effects of bleomycin, cyclophosphamide, and ethyl methanesulfonate on resting and proliferating human B- and T-lymphocytes. *Mutat Res*, 251: 241–251. PMID:1720874
- Mirkes PE, Brown NA, Kajbaf M *et al.* (1992). Identification of cyclophosphamide-DNA adducts in rat embryos exposed in vitro to 4-hydroperoxycyclophosphamide. *Chem Res Toxicol*, 5: 382–385. doi:10.1021/tx00027a010 PMID:1504261
- Montero R, Serrano L, Dávila VM *et al.* (2003). Infection of rats with *Taenia taeniformis* metacestodes increases hepatic CYP450, induces the activity of CYP1A1, CYP2B1 and COH isoforms and increases the genotoxicity of the procarcinogens benzo[a]pyrene, cyclophosphamide and aflatoxin B(1). *Mutagenesis*, 18: 211–216. doi:10.1093/mutage/18.2.211 PMID:12621079
- Moore FR, Urda GA, Krishna G, Theiss JC (1995). An in vivo/in vitro method for assessing micronucleus and chromosome aberration induction in rat bone marrow and spleen. 1. Studies with cyclophosphamide. *Mutat Res*, 335: 191–199. PMID:7477050
- Morales-Ramírez P, Rodríguez-Reyes R, Vallarino-Kelly T (1990). Fate of DNA lesions that elicit sister-chromatid exchanges. *Mutat Res*, 232: 77–88. PMID:2117709
- Mourelatos D, Kritsi Z, Mioglou E, Dozi-Vassiliades J (1993). Enhancement of antineoplastic effect and attenuation of sister chromatid exchanges by prostaglandin E2 in Ehrlich ascites tumour cells treated with cyclophosphamide in vivo. *Prostaglandins Leukot Essent Fatty Acids*, 49: 707–710. doi:10.1016/0952-3278(93)90082-8 PMID:8248278
- Mourelatos D, Mioglou E, Kristi Z, Dozi-Vassiliades J (1995). Enhancement of cytogenetic damage and of antineoplastic effect in lymphoid L1210 leukemia cells treated with prostaglandin E2 and cyclophosphamide in vivo. *Mutat Res*, 326: 125–129. PMID:7528880
- Musatov SA, Anisimov VN, André V *et al.* (1998). Modulatory effects of melatonin on genotoxic response of reference mutagens in the Ames test and the comet assay. *Mutat Res*, 417: 75–84. PMID:9733925
- Nagasubramanian R, Hansen RJ, Delaney SM *et al.* (2008). Survival and tumorigenesis in O6-methylguanine DNA methyltransferase-deficient mice following cyclophosphamide exposure. *Mutagenesis*, 23: 341–346. doi:10.1093/mutage/gen018 PMID:18477655
- Nandakumar A, Davis S, Moolgavkar S *et al.* (1991). Myeloid leukaemia following therapy for a first primary cancer. *Br J Cancer*, 63: 782–788. PMID:2039704
- Natarajan AT & Darroudi F (1991). Use of human hepatoma cells for in vitro metabolic activation of chemical mutagens/carcinogens. *Mutagenesis*, 6: 399–403. doi:10.1093/mutage/6.5.399 PMID:1665540
- Nersessian AK, Zilfian VN, Koumkoumadjian VA (1992). Comparative investigation of cyclophosphamide action on bone marrow cells of the Armenian hamster and 4 other species of rodents. *Mutat Res*, 268: 211–215. PMID:1379326
- Nesterova EV, Durnev AD, Seredenin SB (1999). Verapamil contributes to the clastogenic effects of acrylamide, cyclophosphamide, and dioxidine on somatic cells of BALB/C and C57BL/6 mice. *Mutat Res*, 440: 171–179. PMID:10209340
- Noort D, Hulst AG, Jansen R (2002). Covalent binding of nitrogen mustards to the cysteine-34 residue in human serum albumin. *Arch Toxicol*, 76: 83–88. doi:10.1007/s00204-001-0318-2 PMID:11914777
- O’Neil MJ, editor (2006). *The Merck Index*, 14<sup>th</sup> ed. Whitehouse Station, NJ: Merck & Co., Inc., p. 1554.
- Odagiri Y, Takemoto K, Fenech M (1994). Micronucleus induction in cytokinesis-blocked mouse bone marrow cells in vitro following in vivo exposure to X-irradiation and cyclophosphamide. *Environ Mol Mutagen*, 24: 61–67. doi:10.1002/em.2850240108 PMID:8050417
- Palmer RG, Smith-Burchnell CA, Dore CJ, Denman AM (1986). Thioguanine-resistant mutations induced by cytotoxic drugs in lymphocytes of patients with connective tissue diseases. *Br J Rheumatol*, 25: 376–379. doi:10.1093/rheumatology/25.4.376 PMID:3779323
- Palmer RG, Smith-Burchnell CA, Pelton BK *et al.* (1988). Use of T cell cloning to detect in vivo mutations induced by cyclophosphamide. *Arthritis Rheum*, 31: 757–761. doi:10.1002/art.1780310609 PMID:3289548
- Paolini M, Barone E, Corsi C *et al.* (1991b). Expression and inducibility of drug-metabolizing enzymes in novel murine liver epithelial cell lines and their ability to activate procarcinogens. *Cancer Res*, 51: 301–309. PMID:1988092
- Paolini M, Sapigni E, Hrelia P *et al.* (1991a). Wide spectrum detection of precarcinogens in short-term bioassays by simultaneous superinduction of multiple forms of cytochrome P450 isoenzymes. *Carcinogenesis*, 12: 759–766. doi:10.1093/carcin/12.5.759 PMID:1903089
- Parton JW & Garriott ML (1997). An evaluation of micronucleus induction in bone marrow and in hepatocytes isolated from collagenase perfused liver or from formalin-fixed liver using four-week-old rats treated with known clastogens. *Environ Mol Mutagen*, 29: 379–385. doi:10.1002/(SICI)1098-2280(1997)29:4<379::AID-EM6>3.0.CO;2-5 PMID:9212789
- Pedersen-Bjergaard J, Ersbøll J, Sørensen HM *et al.* (1985). Risk of acute nonlymphocytic leukemia and

- preleukemia in patients treated with cyclophosphamide for non-Hodgkin's lymphomas. Comparison with results obtained in patients treated for Hodgkin's disease and ovarian carcinoma with other alkylating agents. *Ann Intern Med*, 103: 195–200. PMID:4014901
- Petru E, Berger MR, Schmähl D (1989). Long-term carcinogenicity of cyclophosphamide in two mouse strains with different spontaneous leukemia incidence. *Cancer Lett*, 44: 221–226. doi:10.1016/0304-3835(89)90065-7 PMID:2924289
- Poça KS, De-Oliveira AC, Santos MJ, Paumgartten FJ (2008). Malaria infection modulates effects of genotoxic chemicals in the mouse bone-marrow micronucleus test. *Mutat Res*, 649: 28–33. PMID:17851116
- Porter AJ & Singh SM (1988). Transplacental teratogenesis and mutagenesis in mouse fetuses treated with cyclophosphamide. *Teratog Carcinog Mutagen*, 8: 191–203. doi:10.1002/tcm.1770080403 PMID:2906177
- Povirk LF & Shuker DE (1994). DNA damage and mutagenesis induced by nitrogen mustards. *Mutat Res*, 318: 205–226. PMID:7527485
- Premkumar K, Abraham SK, Santhiya ST *et al.* (2001b). Inhibition of genotoxicity by saffron (*Crocus sativus L.*) in mice. *Drug Chem Toxicol*, 24: 421–428. doi:10.1081/DCT-100106266 PMID:11665650
- Premkumar K, Pachappan A, Abraham SK *et al.* (2001a). Effect of *Spirulina fusiformis* on cyclophosphamide and mitomycin-C induced genotoxicity and oxidative stress in mice. *Fitoterapia*, 72: 906–911. doi:10.1016/S0367-326X(01)00340-9 PMID:11731115
- Raposa T & Várkonyi J (1987). The relationship between sister chromatid exchange induction and leukemogenicity of different cytostatics. *Cancer Detect Prev*, 10: 141–151. PMID:3568006
- Rekhadevi PV, Sailaja N, Chandrasekhar M *et al.* (2007). Genotoxicity assessment in oncology nurses handling anti-neoplastic drugs. *Mutagenesis*, 22: 395–401. doi:10.1093/mutage/gem032 PMID:17855733
- Robbiano L, Carrozzino R, Bacigalupo M *et al.* (2002). Correlation between induction of DNA fragmentation in urinary bladder cells from rats and humans and tissue-specific carcinogenic activity. *Toxicology*, 179: 115–128. doi:10.1016/S0300-483X(02)00354-2 PMID:12204548
- Roscher E & Wiebel FJ (1988). Mutagenicity, clastogenicity and cytotoxicity of procarcinogens in a rat hepatoma cell line competent for xenobiotic metabolism. *Mutagenesis*, 3: 269–276. doi:10.1093/mutage/3.3.269 PMID:3045489
- Roschlau G & Justus J (1971). Carcinogenic effect of methotrexate and cyclophosphamide in animal experiment. *Dtsch Gesundheitsw*, 26: 219–222. PMID:5095181
- Rossi AM, Romano M, Zaccaro L *et al.* (1987). DNA synthesis, mitotic index, drug-metabolising systems and cytogenetic analysis in regenerating rat liver. Comparison with bone marrow test after 'in vivo' treatment with cyclophosphamide. *Mutat Res*, 182: 75–82. PMID:3561429
- Royal Pharmaceutical Society of Great Britain (2007). *British National Formulary*, No. 54, London: BMJ Publishing Group Ltd./RPS Publishing.
- Rubes J, Kucharová S, Vozdová M *et al.* (1998). Cytogenetic analysis of peripheral lymphocytes in medical personnel by means of FISH. *Mutat Res*, 412: 293–298. PMID:9600697
- Sakata T, Smith RA, Garland EM, Cohen SM (1989). Rat urinary bladder epithelial lesions induced by acrolein. *J Environ Pathol Toxicol Oncol*, 9: 159–169. PMID:2732910
- Sanderson BJ, Johnson KJ, Henner WD (2001). Induction of mutant lymphocytes in cyclophosphamide- and chlorambucil-treated patients. *Mutagenesis*, 16: 197–202. doi:10.1093/mutage/16.3.197 PMID:11320143
- Sardaş S, Erdoğan F, Sardaş OS *et al.* (1994). Sister chromatid exchange studies for monitoring DNA damage in lymphocytes of malignant lymphoma patients under cytostatic therapy. *Anticancer Drugs*, 5: 487–489. doi:10.1097/00001813-199408000-00017 PMID:7524801
- Saxena AK & Singh G (1998). Cyclophosphamide-induced chromosomal aberrations and associated congenital malformations in rats. *In Vitro Cell Dev Biol Anim*, 34: 751–752. doi:10.1007/s11626-998-0027-8 PMID:9870522
- Schimenti KJ, Hanneman WH, Schimenti JC (1997). Evidence for cyclophosphamide-induced gene conversion and mutation in mouse germ cells. *Toxicol Appl Pharmacol*, 147: 343–350. doi:10.1006/taap.1997.8292 PMID:9439729
- Schmähl D (1967). Carcinogenic action of cyclophosphamide and triaziquone in rats. *Dtsch Med Wochenschr*, 92: 1150–1152.
- Schmähl D (1974). Investigations on the influence of immunodepressive means on the chemical carcinogenesis in rats. *Z Krebsforsch Klin Onkol Cancer Res Clin Oncol*, 81: 211–215. doi:10.1007/BF00305020 PMID:4279516
- Schmähl D & Habs M (1979). Carcinogenic action of low-dose cyclophosphamide given orally to Sprague-Dawley rats in a lifetime experiment. *Int J Cancer*, 23: 706–712. doi:10.1002/ijc.2910230518 PMID:572348
- Schmähl D & Habs MR (1983). Prevention of cyclophosphamide-induced carcinogenesis in the urinary bladder of rats by administration of mesna. *Cancer Treat Rev*, 10: Suppl A57–61. doi:10.1016/S0305-7372(83)80008-5 PMID:6414697
- Schmähl D & Osswald H (1970). Experimental studies on the carcinogenic effects of anticancer chemotherapeutics and immunosuppressive agents. *Arzneimittelforschung*, 20: 1461–1467. PMID:5536412
- Selvakumar E, Prahalathan C, Sudharsan PT, Varalakshmi P (2006). Protective effect of lipoic acid

- on micronuclei induction by cyclophosphamide. *Arch Toxicol*, 80: 115–119. doi:10.1007/s00204-005-0015-7 PMID:16088343
- Sessink PJ, Cerná M, Rössner P *et al.* (1994). Urinary cyclophosphamide excretion and chromosomal aberrations in peripheral blood lymphocytes after occupational exposure to antineoplastic agents. *Mutat Res*, 309: 193–199. PMID:7520976
- Shimkin MB, Weisburger JH, Weisburger EK *et al.* (1966). Bioassay of 29 Alkylating Chemicals by the Pulmonary-Tumor Response in Strain A Mice. *J Natl Cancer Inst*, 36: 915–935.
- Shukla Y, Srivastava B, Arora A, Chauhan LK (2004). Protective effects of indole-3-carbinol on cyclophosphamide-induced clastogenicity in mouse bone marrow cells. *Hum Exp Toxicol*, 23: 245–250. doi:10.1191/0960327104ht441oa PMID:15222402
- Shukla Y & Taneja P (2002). Antimutagenic effects of garlic extract on chromosomal aberrations. *Cancer Lett*, 176: 31–36. doi:10.1016/S0304-3835(01)00774-1 PMID:11790451
- Siddique YH, Beg T, Afzal M (2008). Antigenotoxic effect of apigenin against anti-cancerous drugs. *Toxicol In Vitro*, 22: 625–631. doi:10.1016/j.tiv.2007.12.002 PMID:18206345
- Simula AP & Priestly BG (1992). Species differences in the genotoxicity of cyclophosphamide and styrene in three in vivo assays. *Mutat Res*, 271: 49–58. PMID:1371829
- Smith RA, Williamson DS, Cerny RL, Cohen SM (1990). Detection of 1,N6-propanodeoxyadenosine in acrolein-modified polydeoxyadenylic acid and DNA by 32P post-labeling. *Cancer Res*, 50: 3005–3012. PMID:2334905
- Smith RE, Bryant J, DeCillis A, Anderson S National Surgical Adjuvant Breast and Bowel Project Experience. (2003). Acute myeloid leukemia and myelodysplastic syndrome after doxorubicin-cyclophosphamide adjuvant therapy for operable breast cancer: the National Surgical Adjuvant Breast and Bowel Project Experience. *J Clin Oncol*, 21: 1195–1204. doi:10.1200/JCO.2003.03.114 PMID:12663705
- Soler-Niedziela L, Ong T, Krishna G *et al.* (1989). Sister-chromatid exchange studies on direct- and indirect-acting clastogens in mouse primary cell cultures. *Mutat Res*, 224: 465–470. doi:10.1016/0165-1218(89)90071-2 PMID:2586544
- Souliotis VL, Dimopoulos MA, Sfrikakis PP (2003). Gene-specific formation and repair of DNA monoadducts and interstrand cross-links after therapeutic exposure to nitrogen mustards. *Clin Cancer Res*, 9: 4465–4474. PMID:14555520
- Sweetman SC, editor (2008). *Martindale: The Complete Drug Reference*. London: Pharmaceutical Press. Available at: <http://www.medicinescomplete.com/mc/>
- Sycheva LP (2001). Evaluation of organ specificity of mutagenic effects of cyclophosphamide in mice by micronucleus test. *Bull Exp Biol Med*, 131: 374–376. doi:10.1023/A:1017916622451 PMID:11550030
- Sykes PJ, Hooker AM, Harrington CS *et al.* (1998). Induction of somatic intrachromosomal recombination inversion events by cyclophosphamide in a transgenic mouse model. *Mutat Res*, 397: 209–219. PMID:9541645
- Tafazoli M, Van Hummelen P, Kiefer F, Kirsch-Volders M (1995). Induction of micronuclei in metabolically competent rat hepatoma cell lines by the promutagens 7,12-dimethylbenz[a]anthracene, benzo[a]pyrene and cyclophosphamide. *Mutagenesis*, 10: 15–21. doi:10.1093/mutage/10.1.15 PMID:7739396
- Tates AD, van Dam FJ, Natarajan AT *et al.* (1994). Frequencies of HPRT mutants and micronuclei in lymphocytes of cancer patients under chemotherapy: a prospective study. *Mutat Res*, 307: 293–306. PMID:7513809
- Te C, Gentile JM, Baguley BC *et al.* (1997). In vivo effects of chlorophyllin on the antitumour agent cyclophosphamide. *Int J Cancer*, 70: 84–89. doi:10.1002/(SICI)1097-0215(19970106)70:1<84::AID-IJC13>3.0.CO;2-D PMID:8985095
- Terreros MC, de Luca JC, Furnus CC, Dulout FN (1995). The relation between protein malnutrition, ethanol consumption and chromosomal damage induced by cyclophosphamide in bone marrow cells of mice. *J Vet Med Sci*, 57: 5–8. PMID:7756424
- Tohamy AA, El-Ghor AA, El-Nahas SM, Noshay MM (2003). Beta-glucan inhibits the genotoxicity of cyclophosphamide, adriamycin and cisplatin. *Mutat Res*, 541: 45–53. PMID:14568293
- Tokuoka S (1965). Induction of tumor in mice with N,N-bis(2-chloroethyl)-N',O-propylenephosphoric acid ester diamide (cyclophosphamide). *Gann*, 56: 537–541. PMID:5893414
- Travis LB, Curtis RE, Glimelius B *et al.* (1995). Bladder and kidney cancer following cyclophosphamide therapy for non-Hodgkin's lymphoma. *J Natl Cancer Inst*, 87: 524–530. doi:10.1093/jnci/87.7.524 PMID:7707439
- Travis LB, Curtis RE, Stovall M *et al.* (1994). Risk of leukemia following treatment for non-Hodgkin's lymphoma. *J Natl Cancer Inst*, 86: 1450–1457. doi:10.1093/jnci/86.19.1450 PMID:8089863
- Trenz K, Lugowski S, Jahrsdörfer U *et al.* (2003). Enhanced sensitivity of peripheral blood lymphocytes from women carrying a BRCA1 mutation towards the mutagenic effects of various cytostatics. *Mutat Res*, 544: 279–288. doi:10.1016/j.mrrev.2003.06.011 PMID:14644329
- Tsutsui TW, Inaba T, Fisher LW *et al.* (2006). In vitro chromosome aberration tests using human dental pulp cells to detect the carcinogenic potential of chemical agents. *Odontology*, 94: 44–50. doi:10.1007/s10266-006-0065-1 PMID:16998617
- Turchi G, Nardone A, Palitti F (1992). Application of an epithelial liver cell line, metabolically competent,

- for mutation studies of promutagens. *Mutat Res*, 271: 79–88. PMID:1371832
- Uhl M, Helma C, Knasmüller S (2000). Evaluation of the single cell gel electrophoresis assay with human hepatoma (Hep G2) cells. *Mutat Res*, 468: 213–225. PMID:10882898
- Valagussa P, Moliterni A, Terenzi M *et al.* (1994). Second malignancies following CMF-based adjuvant chemotherapy in resectable breast cancer. *Ann Oncol*, 5: 803–808. PMID:7848882
- Vogel R & Spielmann H (1989). Beneficial effects of ascorbic acid on preimplantation mouse embryos after exposure to cyclophosphamide in vivo. *Teratog Carcinog Mutagen*, 9: 51–59. doi:10.1002/tcm.1770090107 PMID:2567069
- Voskoboinik I, Drew R, Ahokas JT (1997). Peroxisome proliferator nafenopin potentiated cytotoxicity and genotoxicity of cyclophosphamide in the liver and bone marrow cells. *Chem Biol Interact*, 105: 81–97. doi:10.1016/S0009-2797(97)00039-2 PMID:9251722
- Wakata A, Yamashita T, Tamaoki M *et al.* (1989). Micronucleus test with cyclophosphamide administered by intraperitoneal injection and oral gavage. *Mutat Res*, 223: 369–372. doi:10.1016/0165-1218(89)90088-8 PMID:2747720
- Walker SE & Anver MR (1979). Accelerated appearance of neoplasms in female NZB/NZW mice treated with high-dose cyclophosphamide. *Arthritis Rheum*, 22: 1338–1343. doi:10.1002/art.1780221204 PMID:391238
- Walker SE & Anver MR (1983). High incidence of neoplasms in female NZB/NZW mice treated with pulse doses of cyclophosphamide. *Vet Immunol Immunopathol*, 5: 97–104. doi:10.1016/0165-2427(83)90035-1 PMID:6606891
- Walker SE & Bole GG (1971). Augmented incidence of neoplasia in female New Zealand black-New Zealand white (NZB-NZW) mice treated with long-term cyclophosphamide. *J Lab Clin Med*, 78: 978–979. PMID:4943505
- Walker SE & Bole GG Jr (1973). Augmented incidence of neoplasia in NZB-NZW mice treated with long-term cyclophosphamide. *J Lab Clin Med*, 82: 619–633. PMID:4755436
- Walker VE, Andrews JL, Upton PB *et al.* (1999). Detection of cyclophosphamide-induced mutations at the *Hprt* but not the *lacI* locus in splenic lymphocytes of exposed mice. *Environ Mol Mutagen*, 34: 167–181. doi:10.1002/(SICI)1098-2280(1999)34:2/3<167::AID-EM16>3.0.CO;2-O PMID:10529741
- Wang Z, Zheng Q, Liu K *et al.* (2006). Ginsenoside Rh<sub>2</sub> enhances antitumour activity and decreases genotoxic effect of cyclophosphamide. *Basic Clin Pharmacol Toxicol*, 98: 411–415. doi:10.1111/j.1742-7843.2006.pto\_348.x PMID:16623867
- Waters MD & Nolan C (1995). EC/US workshop report: assessment of genetic risks associated with exposure to ethylene oxide, acrylamide, 1,3-butadiene and cyclophosphamide. *Mutat Res*, 330: 1–11. PMID:7623860
- Weisburger JH, Griswold DP, Prejean JD *et al.* (1975). The carcinogenic properties of some of the principal drugs used in clinical cancer chemotherapy. *Recent Results Cancer Res*, 52: 1–17. PMID:138176
- Wilmer JL, Erexson GL, Kligerman AD (1990). Effect of acrolein on phosphoramidate mustard-induced sister chromatid exchanges in cultured human lymphocytes. *Cancer Res*, 50: 4635–4638. PMID:2369740
- Wilson VL, Foiles PG, Chung FL *et al.* (1991). Detection of acrolein and crotonaldehyde DNA adducts in cultured human cells and canine peripheral blood lymphocytes by <sup>32</sup>P-postlabeling and nucleotide chromatography. *Carcinogenesis*, 12: 1483–1490. doi:10.1093/carcin/12.8.1483 PMID:1860170
- Winckler K, Obe G, Madle S *et al.* (1987). Cyclophosphamide: interstrain differences in the production of mutagenic metabolites by S9-fractions from liver and kidney in different mutagenicity test systems in vitro and in the teratogenic response in vivo between CBA and C57 BL mice. *Teratog Carcinog Mutagen*, 7: 399–409. doi:10.1002/tcm.1770070407 PMID:2888218
- Yager JW, Sorsa M, Selvin S (1988). Micronuclei in cytokinesis-blocked lymphocytes as an index of occupational exposure to alkylating cytostatic drugs. *IARC Sci Publ*, 89: 213–216. PMID:3198202
- Yamamoto S, Mitsumori K, Kodama Y *et al.* (1996). Rapid induction of more malignant tumors by various genotoxic carcinogens in transgenic mice harboring a human prototype c-Ha-ras gene than in control non-transgenic mice. *Carcinogenesis*, 17: 2455–2461. doi:10.1093/carcin/17.11.2455 PMID:8968063
- Yusuf AT, Vian L, Sabatier R, Cano JP (2000). In vitro detection of indirect-acting genotoxins in the comet assay using Hep G2 cells. *Mutat Res*, 468: 227–234. PMID:10882899
- Zijlstra JA & Vogel EW (1989). Influence of metabolic factors on the mutagenic effectiveness of cyclophosphamide in *Drosophila melanogaster*. *Mutat Res*, 210: 79–92. PMID:2491914
- Zúñiga G, Torres-Bugañín O, Ramírez-Muñoz MP *et al.* (1996). Micronucleated erythrocytes in splenectomized patients with and without chemotherapy. *Mutat Res*, 361: 107–112. PMID:8980695
- Zúñiga-González G, Gómez-Meda BC, Zamora-Perez A *et al.* (2003). Induction of micronuclei in proestrus vaginal cells from colchicine- and cyclophosphamide-treated rats. *Environ Mol Mutagen*, 42: 306–310. doi:10.1002/em.10202 PMID:14673876
- Zúñiga-González GM, Gómez-Meda BC, Zamora-Perez AL *et al.* (2005). Micronucleated erythrocyte frequencies in old and new world primates: measurement of micronucleated erythrocyte frequencies in peripheral blood of *Callithrix jacchus* as a model for evaluating genotoxicity in primates. *Environ Mol Mutagen*, 46: 253–259. doi:10.1002/em.20154 PMID:15971258

