

Human Papilloma Virus

References to Supplementary Web Tables, Section 2

- Abdel-Hady ES, Martin-Hirsch P, Duggan-Keen M *et al.* (2001). Immunological and viral factors associated with the response of vulval intraepithelial neoplasia to photodynamic therapy. *Cancer Res*, 61:192–196. [PMID:11196160](#)
- Adami HO, Kuper H, Andersson SO *et al.* (2003). Prostate cancer risk and serologic evidence of human papilloma virus infection: a population-based case-control study. *Cancer Epidemiol Biomarkers Prev*, 12:872–875. [PMID:14504197](#)
- Aguayo F, Castillo A, Koriyama C *et al.* (2007). Human papillomavirus-16 is integrated in lung carcinomas: a study in Chile. *Br J Cancer*, 97:85–91 [doi:10.1038/sj.bjc.6603848](#). [PMID:17579626](#)
- Al-Ghamdi A, Freedman D, Miller D *et al.* (2002). Vulvar squamous cell carcinoma in young women: a clinicopathologic study of 21 cases. *Gynecol Oncol*, 84:94–101 [doi:10.1006/gyno.2001.6466](#). [PMID:11748983](#)
- Alotaibi L, Provost N, Gagnon S *et al.* (2006). Diversity of cutaneous human papillomavirus types in individuals with and without skin lesion. *J Clin Virol*, 36:133–140 [doi:10.1016/j.jcv.2006.02.007](#). [PMID:16678481](#)
- Ansink AC, Krul MR, De Weger RA *et al.* (1994). Human papillomavirus, lichen sclerosus, and squamous cell carcinoma of the vulva: detection and prognostic significance. *Gynecol Oncol*, 52:180–184 [doi:10.1006/gyno.1994.1028](#). [PMID:8314136](#)
- Aynaud O, Ionesco M, Barrasso R (1994). Penile intraepithelial neoplasia. Specific clinical features correlate with histologic and virologic findings. *Cancer*, 74:1762–1767 [doi:10.1002/1097-0142\(19940915\)74:6<1762::AID-CNCR2820740619>3.0.CO;2-1](#). [PMID:8082079](#)
- Bahnassy AA, Zekri AR, Abdallah S *et al.* (2005). Human papillomavirus infection in Egyptian esophageal carcinoma: correlation with p53, p21, mdm2, C-erbB2 and impact on survival. *Pathol Int*, 55:53–62 [doi:10.1111/j.1440-1827.2005.01804.x](#). [PMID:15693850](#)
- Balis V, Sourvinos G, Soultzis N *et al.* (2007). Prevalence of BK virus and human papillomavirus in human prostate cancer. *Int J Biol Markers*, 22:245–251. [PMID:18161654](#)
- Barghi MR, Hajimohammadmehdiarbab A, Moghaddam SM, Kazemi B (2005). Correlation between human papillomavirus infection and bladder transitional cell carcinoma. *BMC Infect Dis*, 5:102 [doi:10.1186/1471-2334-5-102](#). [PMID:16274490](#)
- Bergh J, Marklund I, Gustavsson C *et al.* (2007). No link between viral findings in the prostate and subsequent cancer development. *Br J Cancer*, 96:137–139 [doi:10.1038/sj.bjc.6603480](#). [PMID:17117176](#)
- Bezerra AL, Lopes A, Landman G *et al.* (2001). Clinicopathologic features and human papillomavirus dna prevalence of warty and squamous cell carcinoma of the penis. *Am J Surg Pathol*, 25:673–678 [doi:10.1097/0000478-200105000-00017](#). [PMID:11342782](#)
- Bhatla N, Dar L, Patro AR *et al.* (2006). Human papillomavirus type distribution in cervical cancer in Delhi, India. *Int J Gynecol Pathol*, 25:398–402 [doi:10.1097/01.pgp.0000209574.62081.e4](#). [PMID:16990719](#)
- Björge T, Dillner J, Anttila T *et al.* (1997). Prospective seroepidemiological study of role of human papillomavirus in non-cervical anogenital cancers. *BMJ*, 315:646–649. [PMID:9310566](#)
- Björge T, Engeland A, Luostarinen T *et al.* (2002). Human papillomavirus infection as a risk factor for anal and perianal skin cancer in a prospective study. *Br J Cancer*, 87:61–64 [doi:10.1038/sj.bjc.6600350](#). [PMID:12085257](#)
- Bulk S, Berkhof J, Bulkman NWJ *et al.* (2006). Preferential risk of HPV16 for squamous cell carcinoma and of HPV18 for adenocarcinoma of the cervix compared to women with normal cytology in The Netherlands. *Br J Cancer*, 94:171–175 [doi:10.1038/sj.bjc.6602915](#). Bulk SBerkhof JBulkman NW [PMID:16404371](#)
- Carter JJ, Madeleine MM, Shera K *et al.* (2001). Human papillomavirus 16 and 18 L1 serology compared across anogenital cancer sites. *Cancer Res*, 61:1934–1940. [PMID:11280749](#)

- Casabonne D, Michael KM, Waterboer T *et al.* (2007). A prospective pilot study of antibodies against human papillomaviruses and cutaneous squamous cell carcinoma nested in the Oxford component of the European Prospective Investigation into Cancer and Nutrition. *Int J Cancer*, 121:1862–1868 [doi:10.1002/ijc.22885](https://doi.org/10.1002/ijc.22885). [PMID:17565742](https://pubmed.ncbi.nlm.nih.gov/17565742/)
- Castillo A, Aguayo F, Koriyama C *et al.* (2006). Human papillomavirus in esophageal squamous cell carcinoma in Colombia and Chile. *World J Gastroenterol*, 12:6188–6192. [PMID:17036393](https://pubmed.ncbi.nlm.nih.gov/17036393/)
- Chan KW, Lam KY, Chan AC *et al.* (1994). Prevalence of human papillomavirus types 16 and 18 in penile carcinoma: a study of 41 cases using PCR. *J Clin Pathol*, 47:823–826 [doi:10.1136/jcp.47.9.823](https://doi.org/10.1136/jcp.47.9.823). [PMID:7962651](https://pubmed.ncbi.nlm.nih.gov/7962651/)
- Cheng YW, Wu MF, Wang J *et al.* (2007). Human papillomavirus 16/18 E6 oncoprotein is expressed in lung cancer and related with p53 inactivation. *Cancer Res*, 67:10686–10693 [doi:10.1158/0008-5472.CAN-07-1461](https://doi.org/10.1158/0008-5472.CAN-07-1461). [PMID:18006810](https://pubmed.ncbi.nlm.nih.gov/18006810/)
- Clifford GM, Gallus S, Herrero R *et al.*; IARC HPV Prevalence Surveys Study Group (2005). Worldwide distribution of human papillomavirus types in cytologically normal women in the International Agency for Research on Cancer HPV prevalence surveys: a pooled analysis. *Lancet*, 366:991–998 [doi:10.1016/S0140-6736\(05\)67069-9](https://doi.org/10.1016/S0140-6736(05)67069-9). [PMID:16168781](https://pubmed.ncbi.nlm.nih.gov/16168781/)
- Clifford GM, Rana RK, Franceschi S *et al.* (2005). Human papillomavirus genotype distribution in low-grade cervical lesions: comparison by geographic region and with cervical cancer. *Cancer Epidemiol Biomarkers Prev*, 14:1157–1164 [doi:10.1158/1055-9965.EPI-04-0812](https://doi.org/10.1158/1055-9965.EPI-04-0812). [PMID:15894666](https://pubmed.ncbi.nlm.nih.gov/15894666/)
- Coissard CJ, Besson G, Polette MC *et al.* (2005). Prevalence of human papillomaviruses in lung carcinomas: a study of 218 cases. *Mod Pathol*, 18:1606–1609. [PMID:16056245](https://pubmed.ncbi.nlm.nih.gov/16056245/)
- Cubilla AL, Reuter VE, Gregoire L *et al.* (1998). Basaloid squamous cell carcinoma: a distinctive human papilloma virus-related penile neoplasm: a report of 20 cases. *Am J Surg Pathol*, 22:755–761 [doi:10.1097/0000478-199806000-00014](https://doi.org/10.1097/0000478-199806000-00014). [PMID:9630184](https://pubmed.ncbi.nlm.nih.gov/9630184/)
- Cupp MR, Malek RS, Goellner JR *et al.* (1995). The detection of human papillomavirus deoxyribonucleic acid in intraepithelial, in situ, verrucous and invasive carcinoma of the penis. *J Urol*, 154:1024–1029 [doi:10.1016/S0022-5347\(01\)66967-2](https://doi.org/10.1016/S0022-5347(01)66967-2). [PMID:7637047](https://pubmed.ncbi.nlm.nih.gov/7637047/)
- D'Souza G, Kreimer AR, Viscidi R *et al.* (2007). Case-control study of human papillomavirus and oropharyngeal cancer. *N Engl J Med*, 356:1944–1956 [doi:10.1056/NEJMoa065497](https://doi.org/10.1056/NEJMoa065497). [PMID:17494927](https://pubmed.ncbi.nlm.nih.gov/17494927/)
- da Silva CE, da Silva ID, Cerri A, Weckx LL (2007). Prevalence of human papillomavirus in squamous cell carcinoma of the tongue. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*, 104:497–500 [doi:10.1016/j.tripleo.2007.04.028](https://doi.org/10.1016/j.tripleo.2007.04.028). [PMID:17656122](https://pubmed.ncbi.nlm.nih.gov/17656122/)
- Daling JR, Madeleine MM, Johnson LG *et al.* (2004). Human papillomavirus, smoking, and sexual practices in the etiology of anal cancer. *Cancer*, 101:270–280 [doi:10.1002/cncr.20365](https://doi.org/10.1002/cncr.20365). [PMID:15241823](https://pubmed.ncbi.nlm.nih.gov/15241823/)
- Daling JR, Madeleine MM, Schwartz SM *et al.* (2002). A population-based study of squamous cell vaginal cancer: HPV and cofactors. *Gynecol Oncol*, 84:263–270 [doi:10.1006/gyno.2001.6502](https://doi.org/10.1006/gyno.2001.6502). [PMID:11812085](https://pubmed.ncbi.nlm.nih.gov/11812085/)
- de Boer MA, Vet JN, Aziz MF *et al.* (2006). Human papillomavirus type 18 and other risk factors for cervical cancer in Jakarta, Indonesia. *Int J Gynecol Cancer*, 16:1809–1814 [doi:10.1111/j.1525-1438.2006.00701.x](https://doi.org/10.1111/j.1525-1438.2006.00701.x). [PMID:17009976](https://pubmed.ncbi.nlm.nih.gov/17009976/)
- de Oliveira DE, Bacchi MM, Macarenco RSS *et al.* (2006). Human papillomavirus and Epstein-Barr virus infection, p53 expression, and cellular proliferation in laryngeal carcinoma. *Am J Clin Pathol*, 126:284–293 [doi:10.1309/UU2JADUEHDWATVM9](https://doi.org/10.1309/UU2JADUEHDWATVM9). [PMID:16891205](https://pubmed.ncbi.nlm.nih.gov/16891205/)
- Della Torre G, Donghi R, Longoni A *et al.* (1992). HPV DNA in intraepithelial neoplasia and carcinoma of the vulva and penis. *Diagn Mol Pathol*, 1:25–30 [doi:10.1097/00019606-199203000-00004](https://doi.org/10.1097/00019606-199203000-00004). [PMID:1342951](https://pubmed.ncbi.nlm.nih.gov/1342951/)
- Eliezri YD, Silverstein SJ, Nuovo GJ (1990). Occurrence of human papillomavirus type 16 DNA in cutaneous squamous and basal cell neoplasms. *J Am Acad Dermatol*, 23:836–842 [doi:10.1016/0190-9622\(90\)70299-W](https://doi.org/10.1016/0190-9622(90)70299-W). [PMID:2174930](https://pubmed.ncbi.nlm.nih.gov/2174930/)
- Ernster JA, Sciotto CG, O'Brien MM *et al.* (2007). Rising incidence of oropharyngeal cancer and the role of oncogenic human papilloma virus. *Laryngoscope*, 117:2115–2128 [doi:10.1097/MLG.0b013e31813e5fbb](https://doi.org/10.1097/MLG.0b013e31813e5fbb). [PMID:17891052](https://pubmed.ncbi.nlm.nih.gov/17891052/)

- Fei Y, Yang J, Hsieh WC *et al.* (2006). Different human papillomavirus 16/18 infection in Chinese non-small cell lung cancer patients living in Wuhan, China. *Jpn J Clin Oncol*, 36:274–279 [doi:10.1093/jjco/hyl017](https://doi.org/10.1093/jjco/hyl017). PMID:16690730
- Feltkamp MC, Broer R, di Summa FM *et al.* (2003). Seroreactivity to epidermodysplasia verruciformis-related human papillomavirus types is associated with nonmelanoma skin cancer. *Cancer Res*, 63:2695–2700. PMID:12750299
- Ferreira M, Crespo M, Martins L, Félix A (2008). HPV DNA detection and genotyping in 21 cases of primary invasive squamous cell carcinoma of the vagina. *Mod Pathol*, 21:968–972. Forslund O (2007). Genetic diversity of cutaneous human papillomaviruses. *J Gen Virol*, 88:2662–2669.
- Forslund O, Iftner T, Andersson K *et al.*; Viraskin Study Group (2007). Cutaneous human papillomaviruses found in sun-exposed skin: Beta-papillomavirus species 2 predominates in squamous cell carcinoma. *J Infect Dis*, 196:876–883 [doi:10.1086/521031](https://doi.org/10.1086/521031). PMID:17703418
- Frisch M, Fenger C, van den Brule AJ *et al.* (1999). Variants of squamous cell carcinoma of the anal canal and perianal skin and their relation to human papillomaviruses. *Cancer Res*, 59:753–757. PMID:9973228
- Frisch M, Glimelius B, van den Brule AJ *et al.* (1997). Sexually transmitted infection as a cause of anal cancer. *N Engl J Med*, 337:1350–1358 [doi:10.1056/NEJM199711063371904](https://doi.org/10.1056/NEJM199711063371904). PMID:9358129
- Gentile V, Vicini P, Giacomelli L *et al.* (2006). Detection of human papillomavirus DNA, p53 and ki67 expression in penile carcinomas. *Int J Immunopathol Pharmacol*, 19:209–215. PMID:16569359
- Goffin F, Mayrand MH, Gauthier P *et al.* (2006). High-risk human papillomavirus infection of the genital tract of women with a previous history or current high-grade vulvar intraepithelial neoplasia. *J Med Virol*, 78:814–819 [doi:10.1002/jmv.20628](https://doi.org/10.1002/jmv.20628). PMID:16628584
- Gregoire L, Cubilla AL, Reuter VE *et al.* (1995). Preferential association of human papillomavirus with high-grade histologic variants of penile-invasive squamous cell carcinoma. *J Natl Cancer Inst*, 87:1705–1709 [doi:10.1093/jnci/87.22.1705](https://doi.org/10.1093/jnci/87.22.1705). PMID:7473819
- Guerrero D, Guarach R, Ojer A *et al.* (2008). Hypermethylation of the thrombospondin-1 gene is associated with poor prognosis in penile squamous cell carcinoma. *BJU Int*, 102:747–755 [doi:10.1111/j.1464-410X.2008.07603.x](https://doi.org/10.1111/j.1464-410X.2008.07603.x). PMID:18336597
- Gungor A, Cincik H, Baloglu H *et al.* (2007). Human papilloma virus prevalence in laryngeal squamous cell carcinoma. *J Laryngol Otol*, 121:772–774 [doi:10.1017/S0022215107006482](https://doi.org/10.1017/S0022215107006482). PMID:17320003
- Guo L, Zhang L, Lian L (1996). [Clinicopathological features of human papilloma virus associated squamous cell carcinomas of the vulva]. *Zhonghua Fu Chan Ke Za Zhi*, 31:551–554. PMID:9275429
- Gustafsson AC, Ren ZP, Asplund A *et al.* (2004). The role of p53 codon 72 and human papilloma virus status of cutaneous squamous cell carcinoma in the Swedish population. *Acta Derm Venereol*, 84:439–444 [doi:10.1080/00015550410021673](https://doi.org/10.1080/00015550410021673). PMID:15844633
- Hammarstedt L, Lindquist D, Dahlstrand H *et al.* (2006). Human papillomavirus as a risk factor for the increase in incidence of tonsillar cancer. *Int J Cancer*, 119:2620–2623 [doi:10.1002/ijc.22177](https://doi.org/10.1002/ijc.22177). PMID:16991119
- HAMPL M, Deckers-Figiel S, Hampl JA *et al.* (2008). New aspects of vulvar cancer: changes in localization and age of onset. *Gynecol Oncol*, 109:340–345 [doi:10.1016/j.ygyno.2008.01.041](https://doi.org/10.1016/j.ygyno.2008.01.041). PMID:18407339
- HAMPL M, Sarajuuri H, Wentzensen N *et al.* (2006). Effect of human papillomavirus vaccines on vulvar, vaginal, and anal intraepithelial lesions and vulvar cancer. *Obstet Gynecol*, 108:1361–1368 [doi:10.1097/01.AOG.0000245786.86267.80](https://doi.org/10.1097/01.AOG.0000245786.86267.80). PMID:17138767
- Hansson BG, Rosenquist K, Antonsson A *et al.* (2005). Strong association between infection with human papillomavirus and oral and oropharyngeal squamous cell carcinoma: a population-based case-control study in southern Sweden. *Acta Otolaryngol*, 125:1337–1344 [doi:10.1080/00016480510043945](https://doi.org/10.1080/00016480510043945). PMID:16303684
- Heideman DA, Waterboer T, Pawlita M *et al.* (2007). Human papillomavirus-16 is the predominant type etiologically involved in penile squamous cell carcinoma. *J Clin Oncol*, 25:4550–4556 [doi:10.1200/JCO.2007.12.3182](https://doi.org/10.1200/JCO.2007.12.3182). PMID:17925550
- Hillemanns P, Wang X (2006). Integration of HPV-16 and HPV-18 DNA in vulvar intraepithelial neoplasia. *Gynecol Oncol*, 100:276–282 [doi:10.1016/j.ygyno.2005.10.003](https://doi.org/10.1016/j.ygyno.2005.10.003). PMID:16300821

- Hørding U, Junge J, Daugaard S *et al.* (1994). Vulvar squamous cell carcinoma and papillomaviruses: indications for two different etiologies. *Gynecol Oncol*, 52:241–246 [doi:10.1006/gyno.1994.1039](https://doi.org/10.1006/gyno.1994.1039). [PMID:8314146](https://pubmed.ncbi.nlm.nih.gov/8314146/)
- Hørding U, Kringsholm B, Andreasson B *et al.* (1993). Human papillomavirus in vulvar squamous-cell carcinoma and in normal vulvar tissues: a search for a possible impact of HPV on vulvar cancer prognosis. *Int J Cancer*, 55:394–396 [doi:10.1002/ijc.2910550310](https://doi.org/10.1002/ijc.2910550310). [PMID:8397162](https://pubmed.ncbi.nlm.nih.gov/8397162/)
- Indinnimeo M, Cicchini C, Stazi A *et al.* (1999). Human papillomavirus infection and p53 nuclear overexpression in anal canal carcinoma. *J Exp Clin Cancer Res*, 18:47–52. [PMID:10374676](https://pubmed.ncbi.nlm.nih.gov/10374676/)
- Insinga RP, Liaw KL, Johnson LG, Madeleine MM (2008). A systematic review of the prevalence and attribution of human papillomavirus types among cervical, vaginal, and vulvar precancers and cancers in the United States. *Cancer Epidemiol Biomarkers Prev*, 17:1611–1622 [doi:10.1158/1055-9965.EPI-07-2922](https://doi.org/10.1158/1055-9965.EPI-07-2922). [PMID:18628412](https://pubmed.ncbi.nlm.nih.gov/18628412/)
- Iwasawa A, Kumamoto Y, Fujinaga K (1993). Detection of human papillomavirus deoxyribonucleic acid in penile carcinoma by polymerase chain reaction and in situ hybridization. *J Urol*, 149:59–63. [PMID:8380212](https://pubmed.ncbi.nlm.nih.gov/8380212/)
- Iwasawa A, Nieminen P, Lehtinen M, Paavonen J (1997). Human papillomavirus in squamous cell carcinoma of the vulva by polymerase chain reaction. *Obstet Gynecol*, 89:81–84 [doi:10.1016/S0029-7844\(96\)00376-6](https://doi.org/10.1016/S0029-7844(96)00376-6). [PMID:8990443](https://pubmed.ncbi.nlm.nih.gov/8990443/)
- Junge J, Poulsen H, Horn T *et al.* (1995). Human papillomavirus (HPV) in vulvar dysplasia and carcinoma in situ. *APMIS*, 103:501–510 [doi:10.1111/j.1699-0463.1995.tb01398.x](https://doi.org/10.1111/j.1699-0463.1995.tb01398.x). [PMID:7576565](https://pubmed.ncbi.nlm.nih.gov/7576565/)
- Junge J, Poulsen H, Horn T *et al.* (1997). Prognosis of vulvar dysplasia and carcinoma in situ with special reference to histology and types of human papillomavirus (HPV). *APMIS*, 105:963–971 [doi:10.1111/j.1699-0463.1997.tb05108.x](https://doi.org/10.1111/j.1699-0463.1997.tb05108.x). [PMID:9463515](https://pubmed.ncbi.nlm.nih.gov/9463515/)
- Kagawa R, Yamaguchi T, Furuta R (2006). Histological features of human papilloma virus 16 and its association with the development and progression of anal squamous cell carcinoma. *Surg Today*, 36:885–891 [doi:10.1007/s00595-006-3266-x](https://doi.org/10.1007/s00595-006-3266-x). [PMID:16998682](https://pubmed.ncbi.nlm.nih.gov/16998682/)
- Kagie MJ, Kenter GG, Zomerdijk-Nooijen Y *et al.* (1997). Human papillomavirus infection in squamous cell carcinoma of the vulva, in various synchronous epithelial changes and in normal vulvar skin. *Gynecol Oncol*, 67:178–183 [doi:10.1006/gyno.1997.4834](https://doi.org/10.1006/gyno.1997.4834). [PMID:9367704](https://pubmed.ncbi.nlm.nih.gov/9367704/)
- Kamangar F, Qiao YL, Schiller JT *et al.* (2006). Human papillomavirus serology and the risk of esophageal and gastric cancers: results from a cohort in a high-risk region in China. *Int J Cancer*, 119:579–584 [doi:10.1002/ijc.21871](https://doi.org/10.1002/ijc.21871). [PMID:16496409](https://pubmed.ncbi.nlm.nih.gov/16496409/)
- Karagas MR, Nelson HH, Sehr P *et al.* (2006). Human papillomavirus infection and incidence of squamous cell and basal cell carcinomas of the skin. *J Natl Cancer Inst*, 98:389–395 [doi:10.1093/jnci/djj092](https://doi.org/10.1093/jnci/djj092). [PMID:16537831](https://pubmed.ncbi.nlm.nih.gov/16537831/)
- Kim JY, Yoon JK, Citardi MJ *et al.* (2007b). The prevalence of human papilloma virus infection in sinonasal inverted papilloma specimens classified by histological grade. *Am J Rhinol*, 21:664–669 [doi:10.2500/ajr.2007.21.3093](https://doi.org/10.2500/ajr.2007.21.3093). [PMID:18201444](https://pubmed.ncbi.nlm.nih.gov/18201444/)
- Kim SH, Koo BS, Kang S *et al.* (2007a). HPV integration begins in the tonsillar crypt and leads to the alteration of p16, EGFR and c-myc during tumor formation. *Int J Cancer*, 120:1418–1425 [doi:10.1002/ijc.22464](https://doi.org/10.1002/ijc.22464). [PMID:17205528](https://pubmed.ncbi.nlm.nih.gov/17205528/)
- Kim YT, Thomas NF, Kessis TD *et al.* (1996). p53 mutations and clonality in vulvar carcinomas and squamous hyperplasias: evidence suggesting that squamous hyperplasias do not serve as direct precursors of human papillomavirus-negative vulvar carcinomas. *Hum Pathol*, 27:389–395 [doi:10.1016/S0046-8177\(96\)90113-6](https://doi.org/10.1016/S0046-8177(96)90113-6). [PMID:8617483](https://pubmed.ncbi.nlm.nih.gov/8617483/)
- Kiyabu MT, Shibata D, Arnheim N *et al.* (1989). Detection of human papillomavirus in formalin-fixed, invasive squamous carcinomas using the polymerase chain reaction. *Am J Surg Pathol*, 13:221–224 [doi:10.1097/00000478-198903000-00007](https://doi.org/10.1097/00000478-198903000-00007). [PMID:2537583](https://pubmed.ncbi.nlm.nih.gov/2537583/)
- Knopp S, Nesland JM, Tropé C, Holm R (2006). p14ARF, a prognostic predictor in HPV-negative vulvar carcinoma. *Am J Clin Pathol*, 126:266–276 [doi:10.1309/E9E8Y9WG5QE3LK29](https://doi.org/10.1309/E9E8Y9WG5QE3LK29). [PMID:16891203](https://pubmed.ncbi.nlm.nih.gov/16891203/)
- Kohlberger PD, Kirnbauer R, Bancher D *et al.* (1998). Absence of p53 protein overexpression in precancerous lesions of the vulva. *Cancer*, 82:323–327 [doi:10.1002/\(SICI\)1097-0142\(19980115\)82:2<328::AID-CNCR12>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1097-0142(19980115)82:2<328::AID-CNCR12>3.0.CO;2-0). [PMID:9445189](https://pubmed.ncbi.nlm.nih.gov/9445189/)

- Koppikar P, deVilliers EM, Mulherkar R (2005). Identification of human papillomaviruses in tumors of the oral cavity in an Indian community. *Int J Cancer*, 113:946–950 [doi:10.1002/ijc.20664](https://doi.org/10.1002/ijc.20664). [PMID:15514945](https://pubmed.ncbi.nlm.nih.gov/15514945/)
- Korodi Z, Dillner J, Jellum E *et al.* (2005). Human papillomavirus 16, 18, and 33 infections and risk of prostate cancer: a Nordic nested case-control study. *Cancer Epidemiol Biomarkers Prev*, 14:2952–2955 [doi:10.1158/1055-9965.EPI-05-0602](https://doi.org/10.1158/1055-9965.EPI-05-0602). [PMID:16365015](https://pubmed.ncbi.nlm.nih.gov/16365015/)
- Koskinen WJ, Brøndbo K, Mellin Dahlstrand H *et al.* (2007). Alcohol, smoking and human papillomavirus in laryngeal carcinoma: a Nordic prospective multicenter study. *J Cancer Res Clin Oncol*, 133:673–678 [doi:10.1007/s00432-007-0219-8](https://doi.org/10.1007/s00432-007-0219-8). [PMID:17486368](https://pubmed.ncbi.nlm.nih.gov/17486368/)
- Koyamatsu Y, Yokoyama M, Nakao Y *et al.* (2003). A comparative analysis of human papillomavirus types 16 and 18 and expression of p53 gene and Ki-67 in cervical, vaginal, and vulvar carcinomas. *Gynecol Oncol*, 90:547–551 [doi:10.1016/S0090-8258\(03\)00401-3](https://doi.org/10.1016/S0090-8258(03)00401-3). [PMID:13678722](https://pubmed.ncbi.nlm.nih.gov/13678722/)
- Kreimer AR, Clifford GM, Boyle P, Franceschi S (2005). Human papillomavirus types in head and neck squamous cell carcinomas worldwide: a systematic review. *Cancer Epidemiol Biomarkers Prev*, 14:467–475 [doi:10.1158/1055-9965.EPI-04-0551](https://doi.org/10.1158/1055-9965.EPI-04-0551). [PMID:15734974](https://pubmed.ncbi.nlm.nih.gov/15734974/)
- Lai MD, Luo MJ, Yao JE, Chen PH (1998). Anal cancer in Chinese: human papillomavirus infection and altered expression of p53. *World J Gastroenterol*, 4:298–302. [PMID:11819303](https://pubmed.ncbi.nlm.nih.gov/11819303/)
- Laytragoon-Lewin N, Nilsson PJ, Castro J *et al.* (2007). Human papillomavirus (HPV), DNA aberrations and cell cycle progression in anal squamous cell carcinoma patients. *Anticancer Res*, 27 6C:4473–4479. [PMID:18214063](https://pubmed.ncbi.nlm.nih.gov/18214063/)
- Lee YY, Wilczynski SP, Chumakov A *et al.* (1994). Carcinoma of the vulva: HPV and p53 mutations. *Oncogene*, 9:1655–1659. [PMID:8183560](https://pubmed.ncbi.nlm.nih.gov/8183560/)
- Leiros GJ, Galliano SR, Sember ME *et al.* (2005). Detection of human papillomavirus DNA and p53 codon 72 polymorphism in prostate carcinomas of patients from Argentina. *BMC Urol*, 5:15 [doi:10.1186/1471-2490-5-15](https://doi.org/10.1186/1471-2490-5-15). [PMID:16307686](https://pubmed.ncbi.nlm.nih.gov/16307686/)
- Lerma E, Matias-Guiu X, Lee SJ, Prat J (1999). Squamous cell carcinoma of the vulva: study of ploidy, HPV, p53, and pRb. *Int J Gynecol Pathol*, 18:191–197 [doi:10.1097/00004347-199907000-00001](https://doi.org/10.1097/00004347-199907000-00001). [PMID:12090585](https://pubmed.ncbi.nlm.nih.gov/12090585/)
- Levi JE, Rahal P, Sarkis AS, Villa LL (1998). Human papillomavirus DNA and p53 status in penile carcinomas. *Int J Cancer*, 76:779–783 [doi:10.1002/\(SICI\)1097-0215\(19980610\)76:6<779::AID-IJCI>3.0.CO;2-V](https://doi.org/10.1002/(SICI)1097-0215(19980610)76:6<779::AID-IJCI>3.0.CO;2-V). [PMID:9626339](https://pubmed.ncbi.nlm.nih.gov/9626339/)
- Liang XH, Lewis J, Foote R *et al.* (2008). Prevalence and significance of human papillomavirus in oral tongue cancer: the Mayo Clinic experience. *J Oral Maxillofac Surg*, 66:1875–1880 [doi:10.1016/j.joms.2008.04.009](https://doi.org/10.1016/j.joms.2008.04.009). [PMID:18718395](https://pubmed.ncbi.nlm.nih.gov/18718395/)
- Löning T, Riviere A, Henke RP *et al.* (1988). Penile/anal condylomas and squamous cell cancer. A HPV DNA hybridization study. *Virchows Arch A Pathol Anat Histopathol*, 413:491–498. [PMID:2849229](https://pubmed.ncbi.nlm.nih.gov/2849229/)
- Lont AP, Kroon BK, Horenblas S *et al.* (2006). Presence of high-risk human papillomavirus DNA in penile carcinoma predicts favorable outcome in survival. *Int J Cancer*, 119:1078–1081 [doi:10.1002/ijc.21961](https://doi.org/10.1002/ijc.21961). [PMID:16570278](https://pubmed.ncbi.nlm.nih.gov/16570278/)
- Lu XM, Monnier-Benoit S, Mo LZ *et al.* (2008). Human papillomavirus in esophageal squamous cell carcinoma of the high-risk Kazakh ethnic group in Xinjiang, China. *Eur J Surg Oncol*, 34:765–770. [PMID:18160246](https://pubmed.ncbi.nlm.nih.gov/18160246/)
- Luo CW, Roan CH, Liu CJ (2007). Human papillomaviruses in oral squamous cell carcinoma and pre-cancerous lesions detected by PCR-based gene-chip array. *Int J Oral Maxillofac Surg*, 36:153–158 [doi:10.1016/j.ijom.2006.09.005](https://doi.org/10.1016/j.ijom.2006.09.005). [PMID:17110084](https://pubmed.ncbi.nlm.nih.gov/17110084/)
- Madeleine MM, Daling JR, Carter JJ *et al.* (1997). Cofactors with human papillomavirus in a population-based study of vulvar cancer. *J Natl Cancer Inst*, 89:1516–1523 [doi:10.1093/jnci/89.20.1516](https://doi.org/10.1093/jnci/89.20.1516). [PMID:9337348](https://pubmed.ncbi.nlm.nih.gov/9337348/)
- Maden C, Sherman KJ, Beckmann AM *et al.* (1993). History of circumcision, medical conditions, and sexual activity and risk of penile cancer. *J Natl Cancer Inst*, 85:19–24 [doi:10.1093/jnci/85.1.19](https://doi.org/10.1093/jnci/85.1.19). [PMID:8380060](https://pubmed.ncbi.nlm.nih.gov/8380060/)
- Madsen BS, Jensen HL, van den Brule AJ *et al.* (2008). Risk factors for invasive squamous cell carcinoma of the vulva and vagina—population-based case-control study in Denmark. *Int J Cancer*, 122:2827–2834 [doi:10.1002/ijc.23446](https://doi.org/10.1002/ijc.23446). [PMID:18348142](https://pubmed.ncbi.nlm.nih.gov/18348142/)

- Masini C, Fuchs PG, Gabrielli F *et al.* (2003). Evidence for the association of human papillomavirus infection and cutaneous squamous cell carcinoma in immunocompetent individuals. *Arch Dermatol*, 139:890–894 [doi:10.1001/archderm.139.7.890](https://doi.org/10.1001/archderm.139.7.890). PMID:12873884
- McCance DJ, Kalache A, Ashdown K *et al.* (1986). Human papillomavirus types 16 and 18 in carcinomas of the penis from Brazil. *Int J Cancer*, 37:55–59 [doi:10.1002/ijc.2910370110](https://doi.org/10.1002/ijc.2910370110). PMID:3000954
- Milde-Langosch K, Albrecht K, Joram S *et al.* (1995). Presence and persistence of HPV infection and p53 mutation in cancer of the cervix uteri and the vulva. *Int J Cancer*, 63:639–645 [doi:10.1002/ijc.2910630507](https://doi.org/10.1002/ijc.2910630507). PMID:7591279
- Minucci D, Cinel A, Insacco E, Oselladore M (1995). Epidemiological aspects of vaginal intraepithelial neoplasia (VAIN). *Clin Exp Obstet Gynecol*, 22:36–42. PMID:7736640
- Monk BJ, Burger RA, Lin F *et al.* (1995). Prognostic significance of human papillomavirus DNA in vulvar carcinoma. *Obstet Gynecol*, 85:709–715 [doi:10.1016/0029-7844\(95\)00045-S](https://doi.org/10.1016/0029-7844(95)00045-S). PMID:7724101
- Moonen PM, Bakkers JM, Kiemeny LA *et al.* (2007). Human papilloma virus DNA and p53 mutation analysis on bladder washes in relation to clinical outcome of bladder cancer. *Eur Urol*, 52:464–468 [doi:10.1016/j.eururo.2006.11.017](https://doi.org/10.1016/j.eururo.2006.11.017). PMID:17116360
- Nadji SA, Mokhtari-Azad T, Mahmoodi M *et al.* (2007). Relationship between lung cancer and human papillomavirus in north of Iran, Mazandaran province. *Cancer Lett*, 248:41–46 [doi:10.1016/j.canlet.2006.05.016](https://doi.org/10.1016/j.canlet.2006.05.016). PMID:16814459
- Nagano H, Yoshikawa H, Kawana T *et al.* (1996). Association of multiple human papillomavirus types with vulvar neoplasias. *J Obstet Gynaecol Res*, 22:1–8. PMID:8624885
- Newton R, Bousarghin L, Ziegler J *et al.*; Uganda Kaposi's Sarcoma Study Group (2004). Human papillomaviruses and cancer in Uganda. *Eur J Cancer Prev*, 13:113–118 [doi:10.1097/00008469-200404000-00004](https://doi.org/10.1097/00008469-200404000-00004). PMID:15100577
- Noffsinger AE, Hui YZ, Suzuk L *et al.* (1995). The relationship of human papillomavirus to proliferation and ploidy in carcinoma of the anus. *Cancer*, 75:958–967 [doi:10.1002/1097-0142\(19950215\)75:4<958::AID-CNCR2820750410>3.0.CO;2-1](https://doi.org/10.1002/1097-0142(19950215)75:4<958::AID-CNCR2820750410>3.0.CO;2-1). PMID:7842416
- Ogunbiyi OA, Scholefield JH, Robertson G *et al.* (1994). Anal human papillomavirus infection and squamous neoplasia in patients with invasive vulvar cancer. *Obstet Gynecol*, 83:212–216. PMID:8290182
- Osakabe M, Hayashi M, Katayama Y *et al.* (2007). Characteristics of vulvar squamous cell carcinoma in Japanese women. *Pathol Int*, 57:322–327 [doi:10.1111/j.1440-1827.2007.02103.x](https://doi.org/10.1111/j.1440-1827.2007.02103.x). PMID:17539962
- Palefsky JM, Holly EA, Gonzales J *et al.* (1991). Detection of human papillomavirus DNA in anal intraepithelial neoplasia and anal cancer. *Cancer Res*, 51:1014–1019. PMID:1846314
- Pantelis A, Pantelis D, Ruemmele P *et al.* (2007). p53 Codon 72 polymorphism, loss of heterozygosity and high-risk human papillomavirus infection in a low-incidence German esophageal squamous cell carcinoma patient cohort. *Oncol Rep*, 17:1243–1248. PMID:17390072
- Pascual A, Pariente M, Godínez JM *et al.* (2007). High prevalence of human papillomavirus 16 in penile carcinoma. *Histol Histopathol*, 22:177–183. PMID:17149690
- Perceau G, Derancourt C, Clavel C *et al.* (2003). Lichen sclerosus is frequently present in penile squamous cell carcinomas but is not always associated with oncogenic human papillomavirus. *Br J Dermatol*, 148:934–938 [doi:10.1046/j.1365-2133.2003.05326.x](https://doi.org/10.1046/j.1365-2133.2003.05326.x). PMID:12786823
- Petry KU, Köchel H, Bode U *et al.* (1996). Human papillomavirus is associated with the frequent detection of warty and basaloid high-grade neoplasia of the vulva and cervical neoplasia among immunocompromised women. *Gynecol Oncol*, 60:30–34 [doi:10.1006/gyno.1996.0007](https://doi.org/10.1006/gyno.1996.0007). PMID:8557224
- Picconi MA, Eiján AM, Distéfano AL *et al.* (2000). Human papillomavirus (HPV) DNA in penile carcinomas in Argentina: analysis of primary tumors and lymph nodes. *J Med Virol*, 61:65–69. PMID:10745234 [doi:10.1002/\(SICI\)1096-9071\(200005\)61:1<65::AID-JMV10>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1096-9071(200005)61:1<65::AID-JMV10>3.0.CO;2-Z)
- Pinto AP, Schlecht NF, Pintos J *et al.* (2004). Prognostic significance of lymph node variables and human papillomavirus DNA in invasive vulvar carcinoma. *Gynecol Oncol*, 92:856–865 [doi:10.1016/j.ygyno.2003.11.052](https://doi.org/10.1016/j.ygyno.2003.11.052). PMID:14984953
- Pinto AP, Signorello LB, Crum CP *et al.* (1999). Squamous cell carcinoma of the vulva in Brazil: prognostic importance of host and viral variables. *Gynecol Oncol*, 74:61–67 [doi:10.1006/gyno.1999.5458](https://doi.org/10.1006/gyno.1999.5458). PMID:10385552

- Poletti PA, Halfon A, Marti MC (1998). Papillomavirus and anal carcinoma. *Int J Colorectal Dis*, 13:108–111 [doi:10.1007/s003840050145](https://doi.org/10.1007/s003840050145). PMID:9638498
- Prétet JL, Jacquard AC, Saunier M *et al.*; EDiTH study group (2008). Human papillomavirus genotype distribution in low-grade squamous intraepithelial lesions in France and comparison with CIN2/3 and invasive cervical cancer: the EDiTH III study. *Gynecol Oncol*, 110:179–184 [doi:10.1016/j.ygyno.2008.04.012](https://doi.org/10.1016/j.ygyno.2008.04.012). PMID:18514800
- Protzel C, Knoedel J, Zimmermann U *et al.* (2007). Expression of proliferation marker Ki67 correlates to occurrence of metastasis and prognosis, histological subtypes and HPV DNA detection in penile carcinomas. *Histol Histopathol*, 22:1197–1204. PMID:17647192
- Prowse DM, Ktori EN, Chandrasekaran D *et al.* (2008). Human papillomavirus-associated increase in p16INK4A expression in penile lichen sclerosus and squamous cell carcinoma. *Br J Dermatol*, 158:261–265 [doi:10.1111/j.1365-2133.2007.08305.x](https://doi.org/10.1111/j.1365-2133.2007.08305.x). PMID:18047520
- Ramanujam PS, Venkatesh KS, Co Barnett T *et al.* (1996). Study of human papillomavirus infection in patients with anal squamous carcinoma. *Dis Colon Rectum*, 39:37–39 [doi:10.1007/BF02048266](https://doi.org/10.1007/BF02048266). PMID:8601354
- Rosenblatt KA, Carter JJ, Iwasaki LM *et al.* (2003). Serologic evidence of human papillomavirus 16 and 18 infections and risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev*, 12:763–768. PMID:12917208
- Rubin MA, Kleter B, Zhou M *et al.* (2001). Detection and typing of human papillomavirus DNA in penile carcinoma: evidence for multiple independent pathways of penile carcinogenesis. *Am J Pathol*, 159:1211–1218 [doi:10.1016/S0002-9440\(10\)62506-0](https://doi.org/10.1016/S0002-9440(10)62506-0). PMID:11583947
- Rufforny I, Wilkinson EJ, Liu C *et al.* (2005). Human papillomavirus infection and p16(INK4a) protein expression in vulvar intraepithelial neoplasia and invasive squamous cell carcinoma. *J Low Genit Tract Dis*, 9:108–113 [doi:10.1097/00128360-200504000-00007](https://doi.org/10.1097/00128360-200504000-00007). PMID:15870532
- Sagerman PM, Choi YJ, Hu Y, Niedt GW (1996). Human papilloma virus, vulvar dystrophy, and vulvar carcinoma: differential expression of human papillomavirus and vulvar dystrophy in the presence and absence of squamous cell carcinoma of the vulva. *Gynecol Oncol*, 61:328–332 [doi:10.1006/gyno.1996.0151](https://doi.org/10.1006/gyno.1996.0151). PMID:8641610
- Salazar EL, Mercado E, Calzada L (2005). Human papillomavirus hvp-16 DNA as an epitheliotropic virus that induces hyperproliferation in squamous penile tissue. *Arch Androl*, 51:327–334 [doi:10.1080/014850190923396](https://doi.org/10.1080/014850190923396). PMID:16036641
- Sarkar FH, Miles BJ, Plieth DH, Crissman JD (1992). Detection of human papillomavirus in squamous neoplasm of the penis. *J Urol*, 147:389–392. PMID:1310122
- Scholefield JH, Kerr IB, Shepherd NA *et al.* (1991). Human papillomavirus type 16 DNA in anal cancers from six different countries. *Gut*, 32:674–676 [doi:10.1136/gut.32.6.674](https://doi.org/10.1136/gut.32.6.674). PMID:1648029
- Scholefield JH, Sonnex C, Talbot IC *et al.* (1989). Anal and cervical intraepithelial neoplasia: possible parallel. *Lancet*, 2:765–769 [doi:10.1016/S0140-6736\(89\)90830-1](https://doi.org/10.1016/S0140-6736(89)90830-1). PMID:2571011
- Senba M, Kumatori A, Fujita S *et al.* (2006). The prevalence of human papillomavirus genotypes in penile cancers from northern Thailand. *J Med Virol*, 78:1341–1346 [doi:10.1002/jmv.20703](https://doi.org/10.1002/jmv.20703). PMID:16927292
- Shroyer KR, Brookes CG, Markham NE, Shroyer AL (1995). Detection of human papillomavirus in anorectal squamous cell carcinoma. Correlation with basaloid pattern of differentiation. *Am J Clin Pathol*, 104:299–305. PMID:7677119
- Shuyama K, Castillo A, Aguayo F *et al.* (2007). Human papillomavirus in high- and low-risk areas of oesophageal squamous cell carcinoma in China. *Br J Cancer*, 96:1554–1559 [doi:10.1038/sj.bjc.6603765](https://doi.org/10.1038/sj.bjc.6603765). PMID:17453003
- Smith JS, Lindsay L, Hoots B *et al.* (2007). Human papillomavirus type distribution in invasive cervical cancer and high-grade cervical lesions: a meta-analysis update. *Int J Cancer*, 121:621–632 [doi:10.1002/ijc.22527](https://doi.org/10.1002/ijc.22527). PMID:17405118
- Souto Damin AP, Guedes Frazzon AP, de Carvalho Damin D *et al.* (2006). Detection of human papillomavirus DNA in squamous cell carcinoma of the esophagus by auto-nested PCR. *Dis Esophagus*, 19:64–68 [doi:10.1111/j.1442-2050.2006.00541.x](https://doi.org/10.1111/j.1442-2050.2006.00541.x). PMID:16643171
- Srodon M, Stoler MH, Baber GB, Kurman RJ (2006). The distribution of low and high-risk HPV types in vulvar and vaginal intraepithelial neoplasia (VIN and VaIN). *Am J Surg Pathol*, 30:1513–1518 [doi:10.1097/01.pas.0000213291.96401.48](https://doi.org/10.1097/01.pas.0000213291.96401.48). PMID:17122506

- Struijk L, Bouwes Bavinck JN, Wanningen P *et al.* (2003). Presence of human papillomavirus DNA in plucked eyebrow hairs is associated with a history of cutaneous squamous cell carcinoma. *J Invest Dermatol*, 121:1531–1535 [doi:10.1046/j.1523-1747.2003.12632.x](https://doi.org/10.1046/j.1523-1747.2003.12632.x). PMID:14675206
- Struijk L, Hall L, van der Meijden E *et al.* (2006). Markers of cutaneous human papillomavirus infection in individuals with tumor-free skin, actinic keratoses, and squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev*, 15:529–535 [doi:10.1158/1055-9965.EPI-05-0747](https://doi.org/10.1158/1055-9965.EPI-05-0747). PMID:16537712
- Sugase M, Matsukura T (1997). Distinct manifestations of human papillomaviruses in the vagina. *Int J Cancer*, 72:412–415 [doi:10.1002/\(SICI\)1097-0215\(19970729\)72:3<412::AID-IJC7>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1097-0215(19970729)72:3<412::AID-IJC7>3.0.CO;2-S). PMID:9247283
- Sun Y, Hildesheim A, Brinton LA *et al.* (1996). Human papillomavirus-specific serologic response in vulvar neoplasia. *Gynecol Oncol*, 63:200–203 [doi:10.1006/gyno.1996.0306](https://doi.org/10.1006/gyno.1996.0306). PMID:8910627
- Sutcliffe S, Giovannucci E, Gaydos CA *et al.* (2007). Plasma antibodies against Chlamydia trachomatis, human papillomavirus, and human herpesvirus type 8 in relation to prostate cancer: a prospective study. *Cancer Epidemiol Biomarkers Prev*, 16:1573–1580 [doi:10.1158/1055-9965.EPI-07-0134](https://doi.org/10.1158/1055-9965.EPI-07-0134). PMID:17684131
- Sutton BC, Allen RA, Moore WE, Dunn ST (2008). Distribution of human papillomavirus genotypes in invasive squamous carcinoma of the vulva. *Mod Pathol*, 21:345–354 [doi:10.1038/modpathol.3801010](https://doi.org/10.1038/modpathol.3801010). PMID:18192967
- Suzuki H, Sato N, Kodama T *et al.* (1994). Detection of human papillomavirus DNA and state of p53 gene in Japanese penile cancer. *Jpn J Clin Oncol*, 24:1–6. PMID:8126914
- Tachezy R, Jirasek T, Salakova M *et al.* (2007). Human papillomavirus infection and tumours of the anal canal: correlation of histology, PCR detection in paraffin sections and serology. *APMIS*, 115:195–203 [doi:10.1111/j.1600-0463.2007.apm_526.x](https://doi.org/10.1111/j.1600-0463.2007.apm_526.x). PMID:17367464
- Tachezy R, Klozar J, Saláková M *et al.* (2005). HPV and other risk factors of oral cavity/oropharyngeal cancer in the Czech Republic. *Oral Dis*, 11:181–185 [doi:10.1111/j.1601-0825.2005.01112.x](https://doi.org/10.1111/j.1601-0825.2005.01112.x). PMID:15888110
- Termorshuizen F, Feltkamp MC, Struijk L *et al.* (2004). Sunlight exposure and (sero)prevalence of epidermodysplasia verruciformis-associated human papillomavirus. *J Invest Dermatol*, 122:1456–1462 [doi:10.1111/j.0022-202X.2004.22617.x](https://doi.org/10.1111/j.0022-202X.2004.22617.x). PMID:15175037
- Todd RW, Roberts S, Mann CH *et al.* (2004). Human papillomavirus (HPV) type 16-specific CD8+ T cell responses in women with high grade vulvar intraepithelial neoplasia. *Int J Cancer*, 108:857–862 [doi:10.1002/ijc.11645](https://doi.org/10.1002/ijc.11645). PMID:14712488
- Tornesello ML, Duraturo ML, Guida V *et al.* (2008). Analysis of TP53 codon 72 polymorphism in HPV-positive and HPV-negative penile carcinoma. *Cancer Lett*, 269:159–164 [doi:10.1016/j.canlet.2008.04.027](https://doi.org/10.1016/j.canlet.2008.04.027). PMID:18513854
- Trimble CL, Hildesheim A, Brinton LA *et al.* (1996). Heterogeneous etiology of squamous carcinoma of the vulva. *Obstet Gynecol*, 87:59–64 [doi:10.1016/0029-7844\(95\)00351-7](https://doi.org/10.1016/0029-7844(95)00351-7). PMID:8532267
- van Beurden M, ten Kate FJ, Smits HL *et al.* (1995). Multifocal vulvar intraepithelial neoplasia grade III and multicentric lower genital tract neoplasia is associated with transcriptionally active human papillomavirus. *Cancer*, 75:2879–2884 [doi:10.1002/1097-0142\(19950615\)75:12<2879::AID-CNCR2820751214>3.0.CO;2-W](https://doi.org/10.1002/1097-0142(19950615)75:12<2879::AID-CNCR2820751214>3.0.CO;2-W). PMID:7773937
- van Beurden M, ten Kate FW, Tjong-A-Hung SP *et al.* (1998). Human papillomavirus DNA in multicentric vulvar intraepithelial neoplasia. *Int J Gynecol Pathol*, 17:12–16 [doi:10.1097/00004347-199801000-00003](https://doi.org/10.1097/00004347-199801000-00003). PMID:9475186
- van der Avoort IAM, Shirango H, Hoevenaars BM *et al.* (2006). Vulvar squamous cell carcinoma is a multifactorial disease following two separate and independent pathways. *Int J Gynecol Pathol*, 25:22–29 [doi:10.1097/01.pgp.0000177646.38266.6a](https://doi.org/10.1097/01.pgp.0000177646.38266.6a). PMID:16306780
- van Seters M, ten Kate FJ, van Beurden M *et al.* (2007). In the absence of (early) invasive carcinoma, vulvar intraepithelial neoplasia associated with lichen sclerosus is mainly of undifferentiated type: new insights in histology and aetiology. *J Clin Pathol*, 60:504–509 [doi:10.1136/jcp.2005.031989](https://doi.org/10.1136/jcp.2005.031989). PMID:16714399
- Varma VA, Sanchez-Lanier M, Unger ER *et al.* (1991). Association of human papillomavirus with penile carcinoma: a study using polymerase chain reaction and in situ hybridization. *Hum Pathol*, 22:908–913 [doi:10.1016/0046-8177\(91\)90181-N](https://doi.org/10.1016/0046-8177(91)90181-N). PMID:1655618

- Villa LL, Lopes A (1986). Human papillomavirus DNA sequences in penile carcinomas in Brazil. *Int J Cancer*, 37:853–855 [doi:10.1002/ijc.2910370610](https://doi.org/10.1002/ijc.2910370610). [PMID:3011685](https://pubmed.ncbi.nlm.nih.gov/3011685/)
- Vincent-Salomon A, de la Rochefordière A, Salmon R *et al.* (1996). Frequent association of human papillomavirus 16 and 18 DNA with anal squamous cell and basaloid carcinoma. *Mod Pathol*, 9:614–620. [PMID:8782197](https://pubmed.ncbi.nlm.nih.gov/8782197/)
- Waterboer T, Abeni D, Sampogna F *et al.* (2008). Serological association of beta and gamma human papillomaviruses with squamous cell carcinoma of the skin. *Br J Dermatol*, 159:457–459 [doi:10.1111/j.1365-2133.2008.08621.x](https://doi.org/10.1111/j.1365-2133.2008.08621.x). [PMID:18503604](https://pubmed.ncbi.nlm.nih.gov/18503604/)
- Wentzensen N, Schiffman M, Dunn ST *et al.* (2009). Grading the severity of cervical neoplasia based on combined histopathology, cytopathology, and HPV genotype distribution among 1,700 women referred to colposcopy in Oklahoma. *Int J Cancer*, 124:964–969 [doi:10.1002/ijc.23969](https://doi.org/10.1002/ijc.23969). [PMID:19030188](https://pubmed.ncbi.nlm.nih.gov/19030188/)
- Wiener JS, Effert PJ, Humphrey PA *et al.* (1992). Prevalence of human papillomavirus types 16 and 18 in squamous-cell carcinoma of the penis: a retrospective analysis of primary and metastatic lesions by differential polymerase chain reaction. *Int J Cancer*, 50:694–701 [doi:10.1002/ijc.2910500505](https://doi.org/10.1002/ijc.2910500505). [PMID:1312062](https://pubmed.ncbi.nlm.nih.gov/1312062/)
- Williams GR, Lu QL, Love SB *et al.* (1996). Properties of HPV-positive and HPV-negative anal carcinomas. *J Pathol*, 180:378–382. [PMID:9014857](https://pubmed.ncbi.nlm.nih.gov/9014857/) [doi:10.1002/\(SICI\)1096-9896\(199612\)180:4<378::AID-PATH697>3.0.CO;2-H](https://doi.org/10.1002/(SICI)1096-9896(199612)180:4<378::AID-PATH697>3.0.CO;2-H)
- Yanagawa N, Osakabe M, Hayashi M *et al.* (2008). Frequent epigenetic silencing of the FHIT gene in penile squamous cell carcinomas. *Virchows Arch*, 452:377–382 [doi:10.1007/s00428-008-0597-6](https://doi.org/10.1007/s00428-008-0597-6). [PMID:18299890](https://pubmed.ncbi.nlm.nih.gov/18299890/)
- Zaki SR, Judd R, Coffield LM *et al.* (1992). Human papillomavirus infection and anal carcinoma. Retrospective analysis by in situ hybridization and the polymerase chain reaction. *Am J Pathol*, 140:1345–1355. [PMID:1318640](https://pubmed.ncbi.nlm.nih.gov/1318640/)