

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Wong (1987), USA	4602 male chemical workers from 7 plants occupationally exposed to benzene for at least 6 months 1947-1975, and 3074 workers from the same plants not exposed to benzene	Jobs with exposure to benzene classified as continuous and intermittent exposure. Tasks classified by occupational hygienists, with level of exposure assigned from exposure measurements.	KC 189	Benzene Continually exposed <180 ppm-months 180-719 ppm-months ≥ 720 ppm-months	2 2 0	(SMR) 1.5 2.4 -		
Dagg <i>et al.</i> , (1992), USA	Cohort study of 14074 employees of two petroleum refineries who worked at least a year 1950-1980, followed through 1986.	Only first and last jobs coded. No exposure determination performed	KC 189	None	200	(SMR) 0.8 (0.6-0.9)		Non-significant elevated risks for <15 years employment in the two refineries, but reduced risk for employment > 15 years.

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Tsai <i>et al.</i> , (1993), USA	Cohort 4585 workers at two refinery and petrochemical plants who worked more than 6 months before or after January 1 1973 and followed through 1989.	No exposure determination performed	KC 189	None	8	(SMR) 1.3 (0.6-2.6)		
Wong <i>et al.</i> (1993), USA	18135 distribution employees with potential exposure to gasoline for at least 1 year between 1946-1985, followed through June 1989.	8hr TWA total hydrocarbon exposure in ppm estimated for job categories based on industrial hygiene data and expert judgment. Cumulative exposure defined as the sum of products of TWA exposure and duration of exposure of each job in an employee's work history.	KC 189	Land-based employees exposed to gasoline	12	(SMR) 0.7 (0.3-1.1)		
				Marine based employees exposed to gasoline	12	0.8 (0.5-1.4)		

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Honda <i>et al.</i> , (1995), USA	Cohort of 9796 white males who had worked at a petroleum manufacturing plant for 6 months 1942-1989 and followed to January 1 1990.	No exposure determination performed	KC 189	None	23	(SMR) 1.0 (0.6-1.4)		
Collingwood <i>et al.</i> , (1996), USA	Cohort of 4855 refinery workers employed for a minimum of 1 year 1946-1979, asnd followed through 1987.	No exposure determination performed	KC 189		4	(SMR) 0.5 (0.1-1.2)		
Satin <i>et al.</i> , (1996), USA	Cohort of 17,844 petroleum refinery workers, employed 1937-1983, followed through 1987	No exposure determination performed	KC 189		40	(SMR) 1.3 (0.9-1.7)		Reference Texas general population

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Tsai <i>et al.</i> , (1996), USA	9720 employees who had worked at least 3 months 1948-1989 in a petrochemical refinery and chemical plant complex	Workers assigned to refinery (5019) or chemical plant (3988) according to longest duration job. No benzene exposure determination performed	KC 189	Refinery Chemical plant	7 4	(SMR) 0.8 (0.3-1.8) 0.9 (0.2-2.2)		
Lynge <i>et al.</i> , (1997), Denmark, Norway, Sweden, Finland	Cohort of gasoline service station workers (16524 male, 2445 female), followed through 20 years by linkage to national cancer registries	Service station occupation identified from the 1970 censuses of each country	KC 189	Males Females	53 4	(SIR) 1.3 (1.0-1.7) 1.2 (0.3-3.0)		Estimated average level of exposure to benzene below 1 mg/m ³ The SIR for renal pelvis cancer in males was 2.0 (95% CI 1.0-3.7) based on 10 cases (no cases in females).

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Pukkala (1998), Finland	Cohort of 7512 men and 1942 women employed in one group including oil refinery, and chemical manufacturing; follow-up by linkage with the Finnish cancer registry 1971-1994.	No estimate of exposure to benzene	KC 189	Employed at least 3 months during 1967-1982	26	(SIR) 2.0 (1.3-2.9)		
Consonni <i>et al.</i> , (1999), Italy	Cohort of 1583 workers employed 1949-1982 in an oil refinery and followed through May 1991.	No estimate of exposure to benzene	KC 189		3	(SMR) 1.1 (0.2-3.2)		

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Divine <i>et al.</i> (1999a) USA	White men (24604) employees who worked at selected refinery, petrochemical, and research establishments at least one day between 1947-1977; employed at these for a cumulative total >5 years; still employed on the end date of the study	Complete work history of all jobs held at the participating factories	KC 189	Employed: Ever before 1950 1950 and after	55 43 12	(SMR) 0.9 (0.7-1.2) 0.9 (0.7-1.3) 0.9 (0.5-1.6)		No estimates of exposure to benzene
Wong <i>et al.</i> , (2001a) USA	Cohort of 7543 petroleum refinery workers (91% male) employed for at least 1 year, 1945-1996, followed to 1996	No exposure determination performed	KC 189	Male workers Female workers	16 0	(SMR) 1.0 (0.6-1.6) -		No estimates of exposure to benzene

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Wong <i>et al.</i> , (2001b) USA	Cohort of 3328 petroleum refinery workers employed for at least 1 year, 1959-1997, followed to 1997	No exposure determination performed	KC 189	All workers	3	(SMR) 0.6 (0.1-1.8)		No estimates of exposure to benzene

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Sorahan <i>et al.</i> , (2002), United Kingdom	Cohorts of 29630 oil refinery and 16480 petroleum distribution workers, first employed after 1 January 1946 who had worked for at least a year, followed 1951-1998 by linkage with national files	No exposure determination performed	KC 189	Refinery workers Distribution workers	66 45	(SMR) 1.1 (0.8-1.4) 1.2 (0.9-1.6)		
Lewis <i>et al.</i> , (2003), Canada	Cohort of 17230 males and 8062 females employed for at least a year of a petroleum company hired between 1964 and 1994 linked to national mortality and cancer incidence databases to 1994.	Estimates of similar exposure groups based upon job title and location by industrial hygienists	KC 189	Males Females	15 2	(SIR) 1.0 (0.6-1.6) [0.6]		No specific estimates of benzene exposure

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Tsai <i>et al.</i> , (2003), USA	Cohort of 3579 male employees of a chemical and refinery facility during 1973-1999 followed 1979-1999	No exposure determination performed	KC 189	All male workers	6	(SMR) 1.3 (0.5-2.8)		
Bloemen <i>et al.</i> , (2004), USA	Workers exposed to benzene at a chemical plant; 2266 workers (94% male) Followed for up to 30 years.	Job-titles were assigned to exposure categories by an industrial hygienist, based on industrial hygiene measurements.	KC 189	Benzene exposed workers	8	(SMR) 1.2 (0.5-2.3)		

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Huebner <i>et al.</i> , (2004), USA	All employees of two refinery/petrochemical plants (Baton Rouge 6941 males, Baytown 6241 males) active in 1970 or hired between 1970-1982 with at least 1 month of employment at the facility and followed through 1997	No exposure determination performed	KC 189	Hired before 1950 Baton Rouge Baytown Hired in 1950 or later Baton Rouge Baytown	17 13 4 2	(SMR) 1.6 (0.9-2.5) 1.3 (0.7-2.2) 1.2 (0.3-3.0) 0.7 (0.0-2.4)		

Table 2.17. Cohort studies of benzene and kidney cancer

Reference, location	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Sorahan <i>et al.</i> , (2005), United Kingdom	Cohort of 5130 males and 384 females followed for mortality, 1968-2002, and incidence 1971-2001 by linkage to national files.	Occupational exposure to benzene 1966/67 or earlier; as reported by 233 employers	KC 189	All benzene exposed workers	14	(SMR) 1.0 (0.6-1.7)		
					16	(SRR) 0.9 (0.5-1.5)		
Gun <i>et al.</i> (2006), Australia	Australian petroleum workers cohort, 16547 men and 1356 women, followed 1981-1999	None	KC 189		45	(SIR) 1.2 (0.8-1.5)		