

6. References

- Abdallah, F. (2003) Tobacco, how does it taste? Contribution of blending and processing to sensory quality and chemistry of cigarette smoke. *Recent Adv. Tob. Sci.*, **29**, 19–46
- Abernathy, T.J. & Bertrand, L.D. (1992) The prevalence of smokeless tobacco and cigarette use among sixth, seventh and eighth grade students: A longitudinal investigation. *Can. J. public Health*, **83**, 15–18
- Accortt, N.A., Waterbor, J.W., Beall, C. & Howard, G. (2002) Chronic disease mortality in a cohort of smokeless tobacco users. *Am. J. Epidemiol.*, **156**, 730–737
- Adams, J.D., Owens-Tucciarone, P. & Hoffmann, D. (1987) Tobacco-specific *N*-nitrosamines in dry snuff. *Food chem. Toxicol.*, **25**, 245–246
- Adlaf, E.M. & Smart, R.G. (1988) Smokeless tobacco use among Ontario students. *J. Drug Educ.*, **18**, 353–357
- Agarwal, S., Mathur, M., Srivastava, A. & Ralhan, R. (1999) MDM2/p53 co-expression in oral pre-malignant and malignant lesions: Potential prognostic implications. *Oral Oncol.*, **35**, 209–216
- Agrawal, P., Chansoriya, M. & Kaul, K.K. (1983) Effect of tobacco chewing by mothers on placental morphology. *Indian Ped.*, **20**, 561–565
- Ahijevych, K. & Garrett, B.E. (2004) Menthol pharmacology and its potential impact on cigarette smoking behavior. *Nicotine Tob. Res.*, **6** (Suppl. 1), 17–28
- Ahmed, H.G., Idris, A.M. & Ibrahim S.O. (2003) Study of oral epithelial atypia among Sudanese tobacco users by exfoliative cytology. *Anticancer Res.*, **23**, 1943–1949
- Aleksandrova, H.M. (1970) [Role of nass in the development of pathological changes in the mouth.] In: *Epidemiology of Malignant Tumours*, Alma-Ata, Nauka Publishing House, pp. 243–246 (in Russian)

- Alguacil, J. & Silverman, D.T. (2004) Smokeless and other noncigarette tobacco use and pancreatic cancer: A case-control study based on direct interviews. *Cancer Epidemiol. Biomarkers Prev.*, **13**, 55–58
- Al-Idrissi, H.Y. (1990) Head and neck cancer in Saudi Arabia: Retrospective analysis of 65 patients. *J. int. med. Res.*, **18**, 515–519
- Allard, W.F., DeVol, E.B. & Te, O.B. (1999) Smokeless tobacco (shamma) and oral cancer in Saudi Arabia. *Community Dent. oral Epidemiol.*, **27**, 398–405
- Allen, S.S., Hatsukami, D., Gorsline, J. & the Transdermal Nicotine Study Group (1994) Cholesterol changes in smoking cessation using the transdermal nicotine system. *Prev. Med.*, **23**, 190–196
- Allen, S.S., Hatsukami, D., Jensen, J., Grillo, M. & Bliss, R. (1995) Effects of treatment on cardiovascular risk among smokeless tobacco users. *Prev. Med.*, **24**, 357–362
- Amer, M., Bull, C.A., Daouk, M.N., McArthur, P.D., Lundmark, G.J. & El Senoussi, M. (1985) Shamma usage and oral cancer in Saudi Arabia. *Ann. Saudi Med.*, **5**, 135–140
- American Medical Association (1986) Health effects of smokeless tobacco. Council on Scientific Affairs. *J. Am. med. Assoc.*, **255**, 1038–1044
- American Psychiatric Association (1980) *Diagnostic and Statistical Manual of Mental Disorders (DSM-III)*, Washington DC, p. 176
- American Psychiatric Association (1987) *Diagnostic and Statistical Manual of Mental Disorders (DSM-III-R)*, 3rd Ed. rev., Washington DC
- American Psychiatric Association (1994) *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, 4th Ed., Washington DC
- Ammigan, N., Nair, U.J. & Bhide, S.V. (1989a) Effect of masher, a pyrolysed tobacco product on carcinogen metabolizing enzymes. *Indian J. exp. Biol.*, **27**, 692–694
- Ammigan, N., Nair, U.J., Amonkar, A.J. & Bhide, S.V. (1989b) Modulations in the biotransformation of tobacco extract and *N'*-nitrosonornicotine under differential dietary protein status. *J. biochem. Toxicol.*, **4**, 7–13
- Ammigan, N., Nair, U.J., Amonkar, A.J. & Bhide, S.V. (1990a) Effect of tobacco extract and *N'*-nitrosonornicotine on the carcinogen metabolising enzymes under different dietary vitamin B status. *Cancer Lett.*, **52**, 153–159
- Ammigan, N., Nagabhushan, M., Nair, U.J., Amonkar, A.J. & Bhide, S.V. (1990b) Effect of nutritional status on mutagenicity of urine excreted by rats treated with standard/experimental carcinogens. *Indian J. exp. Biol.*, **28**, 711–713
- Ammigan, N., Nair, U.J. & Bhide, S.V. (1990c) Modulation of masher- and benzo[*a*]pyrene-inducible carcinogen-metabolizing enzymes by dietary vitamin A. *Biochem. Med. metab. Biol.*, **44**, 181–191
- Ammigan, N., Nair, U.J., Lalitha, V.S. & Bhide, S.V. (1991) Carcinogenicity studies of masher: Pyrolysed tobacco product, in vitamin-A-deficient Sprague Dawley rats. *J. Cancer Res. clin. Oncol.*, **117**, 50–54
- Andersen, R.A., Burton, H.R., Fleming, P.D. & Hamilton-Kemp, T.R. (1989) Effect of storage conditions on nitrosated, acetylated, and oxidized pyridine alkaloid derivatives in smokeless tobacco products. *Cancer Res.*, **49**, 5895–5900
- Andersson, G. & Warfvinge, G. (2003) The influence of pH and nicotine concentration in oral moist snuff on mucosal changes and salivary pH in Swedish snuff users. *Swed. dent. J.*, **27**, 67–75

- Andersson, G., Axéll, T. & Larsson, A. (1989) Histologic changes associated with the use of loose and portion-bag packed Swedish moist snuff: A comparative study. *J. oral Pathol. Med.*, **18**, 491–497
- Andersson, G., Axéll, T. & Larsson, A. (1990) Impact of consumption factors on soft tissue changes in Swedish moist snuff users: A histologic study. *J. oral Pathol. Med.*, **19**, 453–458
- Andersson, G., Björnberg, G. & Curvall, M. (1994) Oral mucosal changes and nicotine disposition in users of Swedish smokeless tobacco products: A comparative study. *J. oral Pathol. Med.*, **23**, 161–167
- Andersson, G., Vala, E.K. & Curvall, M. (1997) The influence of cigarette consumption and smoking machine yields of tar and nicotine on the nicotine uptake and oral mucosal lesions in smokers. *J. oral Pathol. Med.*, **26**, 117–123
- Archard, H.O. & Tarpley, T.M. (1972) Clinicopathologic and histochemical characterisation of submucosal deposits in snuff dipper's keratosis. *J. oral Pathol.*, **1**, 3–11
- Armitage, A.K. & Turner, D.M. (1970) Absorption of nicotine in cigarette and cigar smoke. *Nature*, **226**, 1231–1232
- Arora, M., Aghi, M. & Reddy, K.S. (2001) *Global Youth Tobacco Survey—Delhi Report* [http://www.cdc.gov/tobacco/global/GYTS/reports/india_newDelhi_2001_searo.htm], accessed 21 June 2004
- Ary, D.V. (1989) Use of smokeless tobacco among male adolescents: Concurrent and prospective relationships. *Natl Cancer Inst. Monogr.*, **8**, 49–55
- Ary, D.V., Lichtenstein, E. & Severson, H.H. (1987) Smokeless tobacco use among male adolescents: Patterns, correlates, predictors, and the use of other drugs. *Prev. Med.*, **16**, 385–401
- Ary, D.V., Lichtenstein, E., Severson, H., Weissman, W. & Seeley, J.R. (1989) An in-depth analysis of male adolescent smokeless tobacco users: Interviews with users and their fathers. *J. behavior. Med.*, **12**, 449–467
- Asal, N.R., Risser, D.R., Kadamani, S., Geyer, J.R., Lee, E.T. & Cherng, N. (1988) Risk factors in renal cell carcinoma: I. Methodology, demographics, tobacco, beverage use, and obesity. *Cancer Detect. Prev.*, **11**, 359–377
- Asplund, K., Nasic, S., Janlert, U. & Stegmayr, B. (2003) Smokeless tobacco as a possible risk factor for stroke in men: A nested case-control study. *Stroke*, **34**, 1754–1759
- Attvall, S., Fowelin, J., Lager, I., Von Schenk, H. & Smith, U. (1993) Smoking induces insulin resistance — A potential link with the insulin resistance syndrome. *J. intern. Med.*, **233**, 327–332
- Autrup, J.L., Hansen, C. & Autrup, H. (1992) Detection of tobacco smoke carcinogen — DNA adducts in cultured rat buccal mucosa cells following exposure to ethanol and total cigarette smoke condensate or chewing tobacco. *Chem.-biol. Interact.*, **85**, 141–150
- Axéll, T.E. (1993) Oral mucosal changes related to smokeless tobacco usage: Research findings in Scandinavia. *Eur. J. Cancer, Part B, Oral Oncol.*, **29**, 299–302
- Axéll, T., Mörnstad, H. & Sundström, B. (1976) The relation of the clinical picture to histopathology of snuff dipper's lesions in a Swedish population. *J. oral Pathol.*, **5**, 229–236
- Axéll, T., Holmstrup, P., Kramer, I.R.A., Pindborg, J.J. & Shear, M. (1984) International seminar on oral leukoplakia and associated lesions related to tobacco habits. *Community Dent. oral Epidemiol.*, **12**, 146–154
- Axelsson, T., Jansson, P.-A., Smith, U. & Eliasson, B. (2001) Nicotine infusion acutely impairs insulin sensitivity in type 2 diabetic patients but not in healthy subjects. *J. intern. Med.*, **249**, 539–544

- Axton, W.F. (1975) *Tobacco and Kentucky*, Lexington, University Press of Kentucky, pp. 8, 25, 58–59
- Ayo-Yusuf, O.A., Swart, T.J.P. & Ayo-Yusuf, I.J. (2000) Prevalence and pattern of snuff dipping in a rural South African population. *S. Afr. dent. J.*, **55**, 610–614
- Ayo-Yusuf, O.A., Swart, T.J.P. & Pickworth, W.B. (2004) Nicotine delivery capabilities of smokeless tobacco products and implications for control of tobacco dependence in South Africa. *Tob. Control*, **13**, 186–189
- Babekir, A.R.E., El Fahal, A.H. & Alla, A.H. (1989) Oesophageal cancer in Sudan. *Trop. Doct.*, **19**, 33–34
- Bagchi, M., Bagchi, D., Hassoun, E.A. & Stohs, S.J. (1994) Smokeless tobacco induced increases in hepatic lipid peroxidation, DNA damage and excretion of urinary lipid metabolites. *Int. J. exp. Pathol.*, **75**, 197–202
- Bagchi, D., Hassoun, E.A., Bagchi, M. & Stohs, S.J. (1995) Protective effects of free radical scavengers and antioxidants against smokeless tobacco extract (STE)-induced oxidative stress in macrophage J774A.1 cell cultures. *Arch. environ. Contam. Toxicol.*, **29**, 424–428
- Bagchi, M., Bagchi, D. & Stohs, S.J. (1996) *In vitro* effects of a smokeless tobacco extract on the production of reactive oxygen species by human oral epidermal cells and rat hepatic mitochondria and microsomes, and peritoneal macrophages. *Arch. environ. Contam. Toxicol.*, **30**, 418–422
- Bagchi, M., Kuszynski, C., Patterson, E.B., Tang, L., Bagchi, D., Saani, T. & Stohs, S.J. (1997) *In vitro* free radical production in human oral keratinocytes induced by smokeless tobacco extract. *In Vitro Toxicol.*, **10**, 263–274
- Bagchi, M., Bagchi, D., Hassoun, E.A. & Stohs, S.J. (1998) Subchronic effects of smokeless tobacco extract (STE) on hepatic lipid peroxidation, DNA damage and excretion of urinary metabolites in rats. *Toxicology*, **127**, 29–38
- Bagchi, M., Balmoori, J., Bagchi, D., Ray, S.D., Kuszynski, C. & Stohs, S.J. (1999) Smokeless tobacco, oxidative stress, apoptosis, and antioxidants in human oral keratinocytes. *Free Radic. Biol. Med.*, **26**, 992–1000
- Bagchi, M., Kuszynski, C.A., Balmoori, J., Joshi, S.S., Stohs, S.J. & Bagchi, D. (2001) Protective effects of antioxidants against smokeless tobacco-induced oxidative stress and modulation of *Bcl-2* and *p53* genes in human oral keratinocytes. *Free Radic. Res.*, **35**, 181–194
- Bagchi, M., Balmoori, J., Bagchi, D., Stohs, S.J., Chakrabarti, J. & Das, D.K. (2002) Role of reactive oxygen species in the development of cytotoxicity with various forms of chewing tobacco and pan masala. *Toxicology*, **179**, 247–255
- Bagwe, A.N. & Bhisey, R.A. (1993) Occupational exposure to tobacco and resultant genotoxicity in bidi industry workers. *Mutat. Res.*, **299**, 103–109
- Bagwe, A.N., Ramchandani, A.G. & Bhisey, R.A. (1994) Skin-tumour-promoting activity of processed bidi tobacco in hairless S/RV Cri-ba mice. *J. Cancer Res. clin. Oncol.*, **120**, 485–489
- Bahrke, M.S. & Baur, T.S. (1988) Tobacco use and performance on the U.S. Army physical fitness test. *Mil. Med.*, **153**, 229–235
- Baker, R.R. & Smith, G. (2003) Toxicological aspects of tobacco flavor ingredients. *Recent Adv. Tob. Sci.*, **29**, 47–76
- Ballweg, J.A. & Bray, R.M. (1989) Smoking and tobacco use by U.S. military personnel. *Mil. Med.*, **154**, 165–168

- Balster, R.L. & Bigelow, G.E. (2003) Guidelines and methodological reviews concerning drug abuse liability assessment. *Drug Alcohol Depend.*, **70**, S13–S40
- Banerjee, A., Pakrashi, A., Chatterjee, S., Ghosh, S. & Dutta, S.K. (1993) Semen characteristics of tobacco users in India. *Arch. Androl.*, **30**, 35–40
- Baral, R.N., Patnaik, S. & Das, B.R. (1998) Co-overexpression of p53 and c-myc proteins linked with advanced stages of betel and tobacco related oral squamous cell carcinomas from eastern India. *Eur. J. oral Sci.*, **106**, 907–913
- Bates, C., Fagerström, K., Jarvis, M.J., Kunze, M., McNeill, A. & Ramström, L. (2003) European Union policy on smokeless tobacco: A statement in favour of evidence based regulation for public health. *Tob. Control*, **12**, 360–367
- Beck, J.D., Koch, G.G. & Offenbacher, S. (1995) Incidence of attachment loss over 3 years in older adults — New and progressing lesions. *Community Dent. oral Epidemiol.*, **23**, 291–296
- Benowitz, N.L. (2003) Cigarette smoking and cardiovascular disease: Pathophysiology and implications for treatment. *Prog. cardiovasc. Dis.*, **46**, 91–111
- Benowitz, N.L. & Gourlay, S.G. (1997) Cardiovascular toxicity of nicotine: Implications for nicotine replacement therapy. *J. Am. Coll. Cardiol.*, **29**, 1422–1431
- Benowitz, N.L. & Henningfield, J.E. (1994) Establishing a nicotine threshold for addiction — The implications for tobacco regulation. *New Engl. J. Med.*, **331**, 123–125
- Benowitz, N.L., Jacob, P., III, Jones, R.T. & Rosenberg, J. (1982) Interindividual variability in the metabolism and cardiovascular effects of nicotine in man. *J. Pharmacol. exp. Ther.*, **221**, 368–372
- Benowitz, N.L., Kuyt, F., Jacob, P., III, Jones, R.T. & Osman, A.-L. (1983) Cotinine disposition and effects. *Clin. Pharmacol. Ther.*, **34**, 604–611
- Benowitz, N.L., Porchet, H., Sheiner, L. & Jacob, P., III (1988) Nicotine absorption and cardiovascular effects with smokeless tobacco use: Comparison with cigarettes and nicotine gum. *Clin. Pharmacol. Ther.*, **44**, 23–28
- Benowitz, N.L., Jacob, P., III & Yu, L. (1989) Daily use of smokeless tobacco: Systemic effects. *Ann. intern. Med.*, **111**, 112–116
- Benowitz, N.L., Jacob, P., III, Fong, I. & Gupta, S. (1994) Nicotine metabolic profile in man: Comparison of cigarette smoking and transdermal nicotine. *J. Pharmacol. exp. Ther.*, **268**, 296–303
- Benowitz, N.L., Hansson, A. & Jacob, P., III (2002) Cardiovascular effects of nasal and transdermal nicotine and cigarette smoking. *Hypertension*, **39**, 1107–1112
- Benowitz, N.L., Herrera, B. & Jacob, P., III (2004) Mentholated cigarette smoking inhibits nicotine metabolism. *J. Pharmacol. exp. Ther.*, **310**, 1208–1215
- Bernzweig, E., Payne, J.B., Reinhardt, R.A., Dyer, J.K. & Patil, K.D. (1998) Nicotine and smokeless tobacco effects on gingival and peripheral blood mononuclear cells. *J. clin. Periodontol.*, **25**, 246–252
- Bhide, S.V., Murdia, U.S. & Nair, J. (1984a) Polycyclic aromatic hydrocarbon profiles of pyrolysed tobacco products commonly used in India. *Cancer Lett.*, **24**, 89–94
- Bhide, S.V., Shah, A.S., Nair, J. & Nagarajrao, D. (1984b) Epidemiological and experimental studies on tobacco related oral cancer in India. In: O'Neill, I.K., von Borstel, R.C., Miller, C.T., Long, J. & Bartsch, H., eds, *N-Nitroso Compounds: Occurrence, Biological Effects and Relevance to Human Cancer* (IARC Scientific Publications No. 57), Lyon, pp. 851–857
- Bhide, S.V., Nair, U.J., Nair, J., Spiegelhalder, B. & Preussmann, R. (1986) *N*-Nitrosamines in the saliva of tobacco chewers or masheri users. *Food chem. Toxicol.*, **24**, 293–297

- Bhide, S.V., Nair, J., Maru, G.B., Nair, U.J., Kameshwar Rao, B.V., Chakraborty, M.K. & Brunnemann, K.D. (1987a) Tobacco-specific *N*-nitrosamines [TNSA] in green mature and processed tobacco leaves from India. *Beit. Tabakforsch.*, **14**, 29–32
- Bhide, S.V., Kulkarni, J., Nair, U.J., Spiegelhalter, B. & Preussmann, R. (1987b) Mutagenicity and carcinogenicity of *masheri*, a pyrolysed tobacco product, and its content of tobacco-specific nitrosamines. In: Bartsch, H., O'Neill, I.K. & Schulte-Herman, R., eds, *Relevance of N-Nitroso Compounds to Human Cancer: Exposures and Mechanisms* (IARC Scientific Publications No. 84), Lyon, pp. 460–462
- Bhide, S.V., Ammigan, N., Nair, U.J. & Lalitha, V.S. (1991) Carcinogenicity studies of tobacco extract in vitamin A-deficient Sprague-Dawley rats. *Cancer Res.*, **51**, 3018–3023
- Bhisey, R.A. & Govekar, R.B. (1991) Biological monitoring of bidi rollers with respect to genotoxic hazards of occupational tobacco exposure. *Mutat. Res.*, **261**, 139–147
- Bhisey, R.A., Bagwe, A.N., Mahimkar, M.B. & Buch, S.C. (1999) Biological monitoring of bidi industry workers occupationally exposed to tobacco. *Toxicol. Lett.*, **108**, 259–265
- Bhonsle, R.B., Murti, P.R., Gupta, P.C. & Mehta, F.S. (1976) Reverse dhumti smoking in Goa. An epidemiologic study of 5449 villagers for oral precancerous lesions. *Indian J. Cancer*, **13**, 301–305
- Bhonsle, R.B., Murti, P.R. & Gupta, P.C. (1992) Tobacco habits in India. In: Gupta, P.C., Hamner, J.E., III. & Murti, P.R., eds, *Control of Tobacco-related Cancers and Other Diseases* (Proceedings of an International Symposium January 15–19, 1990, TIFR, Bombay), Oxford, Oxford University Press, pp. 25–46
- Blackford, K.A., Bailey, P.H. & Coutu-Wakulczyk, G.M. (1994) Tobacco use in northeastern Ontario teenagers: Prevalence of use and associated factors. *Can. J. public Health*, **85**, 89–92
- Blomqvist, G., Hirsch, J.-M. & Alberius, P. (1991) Association between development of lower lip cancer and tobacco habits. *J. oral maxillofac. Surg.*, **49**, 1044–1047
- Blot, W.J., Winn, D.M. & Fraumeni, J.F. (1983) Oral cancer and mouthwash. *J. natl Cancer Inst.*, **70**, 251–253
- Blot, W.J., McLaughlin, J.K., Winn, D.M., Austin, D.F., Greenberg, R.S., Preston-Martin, S., Bernstein, L., Schoenberg, J.B., Stemhagen, A. & Fraumeni, J.F., Jr (1988) Smoking and drinking in relation to oral and pharyngeal cancer. *Cancer Res.*, **48**, 3282–3287
- Bock, F.G., Moore, G.E. & Crouch, S.K. (1964) Tumor-promoting activity of extracts of unburned tobacco. *Science*, **145**, 831–833
- Bock, F.G., Shamberger, R.J. & Myers, H.K. (1965) Tumour-promoting agents in unburned cigarette tobacco. *Nature*, **208**, 584–585
- Boffetta, P., Aagnes, B., Weiderpass, E. & Andersen, A. (2005) Smokeless tobacco use and risk of cancer of the pancreas and other organs. *Int. J. Cancer*, **114**, 992–995
- Bolinder, G. & de Faire, U. (1998) Ambulatory 24-h blood pressure monitoring in healthy, middle-aged smokeless tobacco users, smokers, and nontobacco users. *Am. J. Hypertens.*, **11**, 1153–1161
- Bolinder, G.M., Ahlberg, B.O. & Lindell, J.H. (1992) Use of smokeless tobacco: Blood pressure elevation and other health hazards found in a large-scale population survey. *J. intern. Med.*, **232**, 327–334
- Bolinder, G., Alfredsson, L., Englund, A. & de Faire, U. (1994) Smokeless tobacco use and increased cardiovascular mortality among Swedish construction workers. *Am. J. public Health*, **84**, 399–404

- Bolinder, G., N ren, A., de Faire, U. & Wahren, J. (1997a) Smokeless tobacco use and atherosclerosis: An ultrasonographic investigation of carotid intima media thickness in healthy middle-aged men. *Atherosclerosis*, **132**, 95–103
- Bolinder, G., Nor n, A., Wahren, J. & de Faire, U. (1997b) Long-term use of smokeless tobacco and physical performance in middle-aged men. *Eur. J. clin. Invest.*, **27**, 427–433
- Boyle, R.G., Jensen, J., Hatsukami, D.K. & Severson, H.H. (1995) Measuring dependence in smokeless tobacco users. *Addict. Behav.*, **20**, 443–450
- Braverman, M.T., Svendsen, T., Lund, K.E. & Aar , L.E. (2001) Tobacco use by early adolescents in Norway. *Eur. J. public Health*, **11**, 218–224
- Brinton, L.A., Blot, W.J., Becker, J.A., Winn, D.M., Browder, J.P., Farmer, J.C., Jr & Fraumeni, J.F., Jr (1984) A case–control study of cancers of the nasal cavity and paranasal sinuses. *Am. J. Epidemiol.*, **119**, 896–906
- Brooks, J.E. (1952) *The Mighty Leaf. Tobacco Through the Centuries*, Boston, MA, Little, Brown & Co., pp. 109, 213, 277
- Brown, L.M., Blot, W.J., Schuman, S.H., Smith, V.M., Ershow, A.G., Marks, R.D. & Fraumeni, J.F., Jr (1988) Environmental factors and high risk of esophageal cancer among men in coastal South Carolina. *J. natl Cancer Inst.*, **80**, 1620–1625
- Brown, L.M., Everett, G.D., Gibson, R., Burmeister, L.F., Schuman, L.M. & Blair, A. (1992a) Smoking and risk of non-Hodgkin’s lymphoma and multiple myeloma. *Cancer Causes Control*, **3**, 49–55
- Brown, L.M., Gibson, R., Blair, A., Burmeister, L.F., Schuman, L.M., Cantor, K.P. & Fraumeni, J.F., Jr (1992b) Smoking and risk of leukemia. *Am. J. Epidemiol.*, **135**, 763–768
- Brunnemann, K.D. & Hoffmann, D. (1991) Decreased concentrations of *N*-nitrosodiethanolamine and *N*-nitrosomorpholine in commercial tobacco products. *J. agric. Food Chem.*, **39**, 207–208
- Brunnemann, K.D. & Hoffmann, D. (1992) Chemical composition of smokeless tobacco products. In: *Smokeless Tobacco or Health. An International Perspective* (Smoking and Tobacco Control Monograph No. 2; NIH Publ. No. 93-3461), Bethesda, MD, National Cancer Institute
- Brunnemann, K.D., Genoble, L. & Hoffmann, D. (1985) *N*-Nitrosamines in chewing tobacco: An international comparison. *J. agric. Food Chem.*, **33**, 1178–1181
- Brunnemann, K.D., Genoble, L. & Hoffmann, D. (1987a) Identification and analysis of a new tobacco-specific *N*-nitrosamine, 4-(methylnitrosamino)-4-(3-pyridyl)-1-butanol. *Carcinogenesis*, **8**, 465–469
- Brunnemann, K.D., Hornby, A.P. & Stich, H.F. (1987b) Tobacco-specific nitrosamines in the saliva of Inuit snuff dippers in the northwest territories of Canada. *Cancer Lett.*, **37**, 7–16
- Brunnemann, K.D., Rivenson, A., Adams, J.D., Hecht, S.S. & Hoffmann, D. (1987c) A study of snuff carcinogenesis. In: Bartsch, H., O’Neill, I.K. & Schulte-Herman, R., eds, *Relevance of N-Nitroso Compounds to Human Cancer: Exposures and Mechanisms* (IARC Scientific Publications No. 84), Lyon, pp. 456–459
- Brunnemann, K.D., Qi, J. & Hoffmann, D. (2002) Chemical profile of two types of oral snuff tobacco. *Food. chem. Toxicol.*, **40**, 1699–1703
- Brunnemann, K.D., Qi, J. & Hoffmann, D. (2004) Levels of TSNA in oral moist snuff in the past 30 years and today (Abstract No. 38), In: *58th Tobacco Science Research Conference Proceedings*, Winston-Salem, NC, Tobacco Science Research
- Buch, S.C., Notani, P.N. & Bhisey, R.A. (2002) Polymorphism at *GSTM1*, *GSTM3* and *GSTT1* gene loci and susceptibility to oral cancer in an Indian population. *Carcinogenesis*, **23**, 803–807

- Bundgaard, T., Wildt, J., Frydenberg, M., Elbrønd, O. & Nielsen, J.E. (1995) Case-control study of squamous cell cancer of the oral cavity in Denmark. *Cancer Causes Control*, **6**, 57–67
- Burch, J.D., Rohan, T.E., Howe, G.R., Risch, H.A., Hill, G.B., Steele, R. & Miller, A.B. (1989) Risk of bladder cancer by source and type of tobacco exposure: A case-control study. *Int. J. Cancer*, **44**, 622–628
- Burton, H.R., Bush, L.P & Hamilton, J.L (1983) Effect of curing on the chemical composition of burley tobacco. *Recent Adv. Tob. Sci.*, **9**, 91–153
- Burton, H.R., Bush, L.P., & Djordjevic, M.V. (1989a) Influence of temperature and humidity on the accumulation of tobacco-specific nitrosamines in stored burley tobacco. *J. agric. Food Chem.*, **37**, 1372–1377
- Burton, H.R., Childs, G.H., Jr., Andersen, R.A. & Fleming, P.D. (1989b) Changes in chemical composition of burley tobacco during senescence and curing. 3. Tobacco-specific nitrosamines. *J. agric. Food Chem.*, **37**, 426–430
- Burton, H.R., Dye, N.K. & Bush, L.P. (1994) Relationship between tobacco-specific nitrosamines and nitrite from different air-cured tobacco varieties. *J. agric. Food Chem.*, **42**, 2007–2011
- Bush, L.P., Cui, M., Shi, H., Burton, H.R., Fannin, F.F., Lei, L. & Dye, N. (2001) Formation of tobacco-specific nitrosamines in air-cured tobacco. *Recent Adv. Tob. Sci.*, **27**, 23–46
- Byrd, G.D., Chang, K.-M., Greene, J.M. & deBethizy, J.D. (1992) Evidence for urinary excretion of glucuronide conjugates of nicotine, cotinine, and *trans*-3'-hydroxycotinine in smokers. *Drug Metab. Dispos.*, **20**, 192–197
- Calmels, S., Ohshima, H., Henry, Y. & Bartsch, H. (1996) Characterization of bacterial cytochrome cd₁-nitrite reductase as one enzyme responsible for catalysis of nitrosation of secondary amines. *Carcinogenesis*, **17**, 533–536
- Carmella, S.G., Kagan, S.S., Kagan, M., Foiles, P.G., Palladino, G., Quart, A.M., Quart, E. & Hecht, S.S. (1990) Mass spectrometric analysis of tobacco-specific nitrosamine hemoglobin adducts in snuff dippers, smokers, and nonsmokers. *Cancer Res.*, **50**, 5438–5445
- Carmella, S.G., Borukhova, A., Akerkar, S.A. & Hecht, S.S. (1997) Analysis of human urine for pyridine-*N*-oxide metabolites of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, a tobacco-specific lung carcinogen. *Cancer Epidemiol. Biomarkers Prevent.*, **6**, 113–120
- Carmella, S.G., Le, K.-A., Upadhyaya, P. & Hecht, S.S. (2002) Analysis of *N*- and *O*-glucuronides of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in human urine. *Chem. Res. Toxicol.*, **15**, 545–550
- Carossa, S., Pera, P., Doglio, P., Lombardo, S., Colagrande, P., Brussino, L., Rolla, G. & Bucca, C. (2001) Oral nitric oxide during plaque deposition. *Eur. J. clin. Invest.*, **31**, 876–879
- Carter, B.L., Long, T.Y. & Cinciripini, P.M. (2004) A meta-analytic review of the CYP2A6 genotype and smoking behavior. *Nicotine Tob. Res.*, **6**, 221–227
- Castrucci, B.C., Gerlach, K.K., Kaufman, N.J. & Orleans, C.T. (2004) Tobacco use and cessation behavior among adolescents participating in organized sports. *Am. J. Health Behav.*, **28**, 63–71
- CDC (Centers for Disease Control and Prevention) (1999a) Determination of nicotine, pH, and moisture content of six U.S. commercial moist snuff products — Florida, January–February 1999. *Mortal. Morbid. Wkly Rep.*, **48**, 398–401
- CDC (Centers for Disease Control and Prevention) (1999b) Notice regarding requirement for annual submission of the quantity of nicotine contained in smokeless tobacco products manu-

- factured, imported, or packaged in the United States. Centers for Disease Control and Prevention (CDC), Department of Health and Human Services. Notice. *Fed. Regist.*, **64**, 14085–14096
- Chacko, M. & Gupta, R.C. (1988) Evaluation of DNA damage in the oral mucosa of tobacco users and non-users by ³²P-adduct assay. *Carcinogenesis*, **9**, 2309–2313
- Chakradeo, P.P., Nair, J. & Bhide, S.V. (1994) Endogenous formation of *N*-nitrosoproline and other *N*-nitrosamino acids in tobacco users. *Cancer Lett.*, **86**, 187–194
- Chakravarti, N., Mathur, M., Bahadur, S., Shukla, N.K., Rochette-Egly, C. & Ralhan, R. (2001) Expression of RAR α and RAR β in human oral potentially malignant and neoplastic lesions. *Int. J. Cancer*, **91**, 27–31
- Chakravarti, N., Mathur, M., Bahadur, S., Shukla, N.K. & Ralhan, R. (2003) Retinoic acid receptor-alpha as a prognostic indicator in oral squamous cell carcinoma. *Int. J. Cancer*, **103**, 544–549
- Chakravorty, B., Ahmed, A. & Buchanan, R.J. (2000) Midproject findings from a study of the National Collegiate Athletic Association's policy on smokeless tobacco use. *Subst. Use Misuse*, **35**, 1431–1441
- Chamberlain, W.J. & Chortyk, O.T. (1992) Effects of curing and fertilization on nitrosamine formation in bright and burley tobacco. *Beitr. Tabakforsch.*, **15**, 87–92
- Chamberlain, W.J., Schlotzhauer, W.S. & Chortyk, O.T. (1988) Chemical composition of non-smoking tobacco products. *J. agric. Food Chem.*, **36**, 48–50
- Chan, G., Boyle, J.O., Yang, E.K., Zhang, F., Sacks, P.G., Shah, J.P., Edelstein, D., Soslow, R.A., Koki, A.T., Woerner, B.M., Masferrer, J.L. & Dannenberg, A.J. (1999) Cyclooxygenase-2 expression is up-regulated in squamous cell carcinoma of the head and neck. *Cancer Res.*, **59**, 991–994
- Chandra, A. (1962) Different habits and their relation with cancer cheek. *Chittaranjan Cancer Hosp. Calcutta natl Cancer Res. Centre Bull.*, 33–36
- Chang, T., Chowdhry, S., Budhu, P. & Kew, R.R. (1998) Smokeless tobacco extracts activate complement *in vitro*: A potential pathogenic mechanism for initiating inflammation of the oral mucosa. *Clin. Immunol. Immunopathol.*, **87**, 223–229
- Chao, A., Thun, M.J., Henley, S.J., Jacobs, E.J., McCullough, M.L. & Calle, E.E. (2002) Cigarette smoking, use of other tobacco products and stomach cancer mortality in US adults: The Cancer Prevention Study II. *Int. J. Cancer*, **101**, 380–389
- Chapman, S. & Wakefield, M. (2001) Tobacco control advocacy in Australia: Reflections on 30 years of progress. *Health Educ Behav.*, **28**, 274–289
- Chaudhry, K. (2001) *Prevalence of Tobacco Use in Karnataka and Uttar Pradesh in India 2001*, Indian Council of Medical Research
- Chen, S.-Y. (1989) Effects of smokeless tobacco on the buccal mucosa of HMT rats. *J. oral. Pathol. Med.*, **18**, 108–112
- Chisick, M.C., Poindexter, F.R. & York, A.K. (1998) Comparing tobacco use among incoming recruits and military personnel on active duty in the United States. *Tob. Control*, **7**, 236–240
- Chow, W.-H., McLaughlin, J.K., Menck, H.R. & Mack, T.M. (1994) Risk factors for extrahepatic bile duct cancers: Los Angeles County, California (USA). *Cancer Causes Control*, **5**, 267–272
- Christen, A.G. & Glover, E.D. (1981) Smokeless tobacco: Seduction of youth. *World Smoking Health*, **6**, 20–24
- Christen, A.G., Armstrong, W.R. & McDaniel, R.K. (1979) Intraoral leukoplakia, abrasion, periodontal breakdown and tooth loss in a snuff dipper. *J. Am. dent. Assoc.*, **98**, 584–586

- Christen, A.G., Swanson, B.Z., Glover, E.D., Henderson, A.H. (1982) Smokeless tobacco: The folklore and social history of snuffing, sneezing, dipping, and chewing. *J. Am. dent. Assoc.*, **105**, 821–829
- Ciolino, L.A., McCauley, H.A., Fraser, D.B. & Wolnik, K.A. (2001) The relative buffering capacities of saliva and moist snuff: Implications for nicotine absorption. *J. anal. Toxicol.*, **25**, 15–25
- Clark, P.I., Natanblut, S.L., Schmitt, C.L., Wolters, C. & Iachan, R. (2000) Factors associated with tobacco sales to minors: Lessons learned from the FDA compliance checks. *J. Am. med. Assoc.*, **284**, 729–734
- Colborn, J.W., Cummings, K.M. & Michalek, A.M. (1989) Correlates of adolescents' use of smokeless tobacco. *Health Educ. Q.*, **16**, 91–100
- Coleman, W.M., III & Perfetti, T.A. (1997) The roles of amino acids and sugars in the production of volatile materials in microwave heated tobacco dust suspensions. *Beit. Tabakforsch.*, **17**, 75–95
- Connolly, G. (1995) The marketing of nicotine addiction by one oral snuff manufacturer. *Tob. Control*, **4**, 73–79
- Connolly, G.N. (2001) *Establishing Tolerance Limits for Tobacco Specific Nitrosamines (TSNAs) in Oral Snuff under the Massachusetts Hazardous Substance Act Mg L c 94 s-*, Massachusetts Department of Public Health
- Connolly, G.N., Winn, D.M., Hecht, S.S., Henningfield, J.E., Walker, B., Jr & Hoffmann, D. (1986) The reemergence of smokeless tobacco. *New Engl. J. Med.*, **314**, 1020–1027
- Coogan, P.F., Geller, A. & Adams, M. (2000) Prevalence and correlates of smokeless tobacco use in a sample of Connecticut students. *J. Adolesc.*, **23**, 129–135
- Cooper, J., Ellison, J.A. & Walsh, M.M. (2003) Spit (smokeless)-tobacco use by baseball players entering the professional ranks. *J. Athl. Train.*, **38**, 126–132
- Creath, C.J., Shelton, W.O., Wright, J.T., Bradley, D.H., Feinstein, R.A. & Wisniewski, J.F. (1988) The prevalence of smokeless tobacco use among adolescent male athletes. *J. Am. dent. Assoc.*, **116**, 43–48
- Creath, C.J., Cutter, G., Bradley, D.H. & Wright, J.T. (1991) Oral leukoplakia and adolescent smokeless tobacco use. *Oral Surg. oral Med. oral Pathol.*, **72**, 35–41
- Critchley, J.A. & Unal, B. (2003) Health effects associated with smokeless tobacco: A systematic review. *Thorax*, **58**, 435–443
- Cummings, K.M., Michalek, A.M., Carl, W., Wood, R. & Haley, N.J. (1989) Use of smokeless tobacco in a group of professional baseball players. *J. behav. Med.*, **12**, 559–567
- Curtis, M.M. (1935) *The Book of Snuff and Snuff Boxes*, New York, Bramhall House, pp. 20–27, 67
- Curvall, M., Romert, L., Norlén, E. & Enzell, C.R. (1987) Mutagen levels in urine from snuff users, cigarette smokers and non tobacco users — A comparison. *Mutat. Res.*, **188**, 105–110
- Dahl, B.L., Stølen, S.O. & Øilo, G. (1989) Abrasives in snuff? *Acta odontol. scand.*, **47**, 239–243
- D'Alessandro, A., Benowitz, N.L., Muzi, G., Eisner, M.D., Filiberto, S., Fantozzi, P., Montanari, L. & Abbritti, G. (2001) Systemic nicotine exposure in tobacco harvesters. *Arch. environ. Health*, **56**, 257–263
- Daniels, T.E., Hansen, L.S., Greenspan, J.S., Grady, D.G., Hauck, W.W., Greene, J.C. & Ernster, V.L. (1992) Histopathology of smokeless tobacco lesions in professional baseball players. *Oral Surg. oral Med. oral Pathol.*, **73**, 720–725

- Dannenberg, A.J., Altorki, N.K., Boyle, J.O., Dang, C., Howe, L.R., Weksler, B.B. & Subbaramaiah, K. (2001) Cyclo-oxygenase 2: A pharmacological target for the prevention of cancer. *Lancet Oncol.*, **2**, 544–551
- Das, R.K. & Dash, B.C. (1992) Genotoxicity of ‘gudakhu’, a tobacco preparation. II. In habitual users. *Food chem. Toxicol.*, **30**, 1045–1049
- Davis, T.C., Arnold, C., Nandy, I., Bocchini, J.A., Gottlieb, A., George, R.B. & Berkel, H. (1997) Tobacco use among male high school athletes. *J. adolesc. Health*, **21**, 97–101
- Demirci, C.S., Miller, D.R., Baillargeon, J., Moyer, M.P. & Jenson, H.B. (2000) Smokeless tobacco extracts modulate exogenous gene expression in early passage cultured human oral epithelial cells: An *in vitro* system to study chemical and viral enhancer/promoter interaction. *J. pharmacol. toxicol. Meth.*, **44**, 527–532
- Dent, C.W., Sussman, S., Johnson, C.A., Hansen, W.B. & Flay, B.R. (1987) Adolescent smokeless tobacco incidence: Relations with other drugs and psychosocial variables. *Prev. Med.*, **16**, 422–431
- Department of Agriculture (2003) *Tobacco Situation and Outlook Yearbook (Report No. TBS-2003)*, Washington, DC, Market and Trade Economics Division, Economic Research Service
- Department of Agriculture (2006) *Tobacco Situation and Outlook Yearbook (Report No. TBS-2006)*, Washington, DC, Market and Trade Economics Division, Economic Research Service
- Department of Commerce (2004) U.S. Census Bureau, Population Division; Census Data for Public Health Research, CDC WONDER On-line Database, accessed 22 July 2004
- DHHS (Department of Health and Human Services) (1983) *The Health Consequences of Smoking: Cardiovascular Disease — A Report of the Surgeon General* (DHHS Publ. No. (CDC) 84-50204), Washington DC, US Government Printing Office
- DHHS (Department of Health and Human Services) (1986) *The Health Consequences of Using Smokeless Tobacco* (NIH Publication No. 86-2874), Washington DC, Public Health Service
- DHHS (Department of Health and Human Services) (1988) *The Health Consequences of Smoking: Nicotine Addiction. A Report of the Surgeon General* (DHHS Publ. No. (CDC) 88-8406), Washington DC, US Government Printing Office
- DHHS (Department of Health and Human Services) (1989) *Reducing the Health Consequences of Smoking: 25 years of progress* (DHHS Publication No. (CDC) 89-8411), Rockville, MD, Centers for Disease Control, Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health
- DHHS (Department of Health and Human Services) (1990) *The Health Benefits of Smoking Cessation: A report of the Surgeon General* (DHHS Publ. No. (CDC) 90-8416), Rockville, MD
- DHHS (Department of Health and Human Services) (1995) Food and Drug Administration: Analysis regarding the food and drug administration’s jurisdiction over nicotine-containing cigarettes and smokeless tobacco products. *Federal Register*, **60**, 41453
- DHHS (Department of Health and Human Services) (1996) Food and Drug Administration: Analysis regarding the food and drug administration’s jurisdiction over nicotine-containing cigarettes and smokeless tobacco products: Reopening of the comment period as to specific documents. *Federal Register*, **61**, 11419
- DHHS (Department of Health and Human Services) (2004a) Centers for Disease Control and Prevention, National Center for Health Statistics, Bridged-Race Population Estimates, United States, 1990–2002, with 5-year Age Groups, on CDC WONDER On-line Database, 22 July 2004

- DHHS (Department of Health and Human Services) (2004b) *The Health Consequences of Smoking: A Report of the Surgeon General*, Atlanta, GA, US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health [http://www.cdc.gov/tobacco/sgr/sgr_2004/chapters.htm]
- Dikshit, R.K., Buch, J.G. & Mansuri, S.M. (1987) Effect of tobacco consumption on semen quality of a population of hypofertile males. *Fertil. Steril.*, **48**, 334–336
- Directorate of Health and Social Affairs (2001) *Tobakksbruk i Ungdomsskolen, November 2000* [Tobacco use among schoolchildren, November 2000], unpublished data received from Department of Tobacco Control (in Norwegian)
- Directorate of Health and Social Affairs (2003) *Tall om Tobakk 1973–2001* [Tobacco in figures 1973–2001] (in Norwegian)
- Directorate of Health and Social Affairs (2005) *Forbruk av Tobakk* [Tobacco consumption] [available at http://www.shdir.no/tobakk/statistikk/forbruk_av_tobakk; accessed 25.08.2006]
- Directorate of Health and Social Affairs (2006a) *Bruk av Snus i 2005* [Use of snuff in 2005] [available at http://Consumption of snuff in 2005/bruk_av_snus/]
- Directorate of Health and Social Affairs (2006b) *Røyking blant Unge* [Smoking among youth] [available at http://Smoking among the youth/r_yking_blant_unge/]
- Directorate of Health and Social Affairs (2006c) *Tobakksbruk i Ungdomsskolen, Januar 2006* [Tobacco use among schoolchildren, January 2006], unpublished data received from Department of Tobacco Control (in Norwegian)
- Directorate of Tobacco Development (1976–98) *Tobacco in India. A Handbook of Statistics (Various Issues)*, Ministry of Agriculture, Government of India
- Djordjevic, M.V., Gay, S.L., Bush, L.P., & Chaplin, J.F. (1989a) Tobacco-specific nitrosamine accumulation and distribution in flue-cured tobacco alkaloid isolines. *J. agric. Food Chem.*, **37**, 752–756
- Djordjevic, M.V., Brunnemann, K.D. & Hoffmann, D. (1989b) Identification and analysis of a nicotine-derived *N*-nitrosamino acid and other nitrosamino acids in tobacco. *Carcinogenesis*, **10**, 1725–1731
- Djordjevic, M.V., Fan, J., Bush, L.P., Brunnemann, K.D. & Hoffmann, D. (1993a) Effects of storage conditions on levels of tobacco-specific *N*-nitrosamines and *N*-nitrosamino acids in U.S. moist snuff. *J. agric. Food Chem.*, **41**, 1790–1794
- Djordjevic, M.V., Brunnemann, K.D. & Hoffmann, D. (1993b) The need for regulation of carcinogenic *N*-nitrosamines in oral snuff. *Food chem. Toxicol.*, **31**, 497–501
- Djordjevic, M.V., Fan, J., Krzeminski, J., Brunnemann, K.D. & Hoffmann, D. (1994) Characterization of *N*-nitrosoamino acids in tobacco products and assessment of their carcinogenic potential. In: Loeppky, R.N. & Michejda, C.J., eds, *Nitrosamines and Related N-Nitroso Compounds, Chemistry and Biochemistry* (ACS Symposium Series No. 553), Washington DC, American Chemical Society, pp. 367–368
- Djordjevic, M.V., Hoffmann, D., Glynn, T. & Connolly, G.N. (1995) US commercial brands of moist snuff, 1994. I. Assessment of nicotine, moisture, and pH. *Tob. Control*, **4**, 62–66
- Dokko, H., Min, P.S., Cherrick, H.M. & Park, N.-H. (1991) Effect of smokeless tobacco and tobacco-related chemical carcinogens on survival of ultraviolet light-inactivated herpes simplex virus. *Oral Surg. oral Med. oral Pathol.*, **71**, 464–468

- D'Orlando, K.J. & Fox, B.S. (2004) Tolerability and pharmacokinetics of single and repeated doses of nicotine with the straw, a novel nicotine replacement product. *Nicotine Tob. Res.*, **6**, 63–70
- Du, X., Squier, C.A. Kremer, M.J. & Wertz, P.W. (2000) Penetration of N-nitrosornicotine (NNN) across oral mucosa in the presence of ethanol and nicotine. *J. oral Pathol. Med.*, **29**, 80–85
- Dunham, L.J. & Herrold, K.M. (1962) Failure to produce tumors in the hamster cheek pouch by exposure to ingredients of betel quid; histopathologic changes in the pouch and other organs by exposure to known carcinogens. *J. natl. Cancer Inst.*, **29**, 1047–1051
- Dunham, L.J., Muir, C.S. & Hamner, J.E., III (1966) Epithelial atypia in hamster cheek pouches treated repeatedly with calcium hydroxide. *Br. J. Cancer*, **20**, 588–593
- Dunham, L.J., Snell, K.C. & Stewart, H.L. (1975) Argyrophilic carcinoids in two Syrian hamsters (*Mesocricetus auratus*). *J. natl. Cancer Inst.*, **54**, 507–509
- Dunn, B.P. & Stich, H.F. (1986) ³²P-Postlabelling analysis of aromatic DNA adducts in human oral mucosal cells. *Carcinogenesis*, **7**, 1115–1120
- van Duser, B.L. & Raven, P.B. (1992) The effects of oral smokeless tobacco on the cardiorespiratory response to exercise. *Med. Sci. Sports Exerc.*, **24**, 389–395
- Ebbert, J.O., Rowland, L.C., Montori, V.M., Vickers, K.S., Erwin, P.J. & Dale, L.C. (2003) Treatments for spit tobacco use: A quantitative systematic review. *Addiction*, **98**, 569–583
- Ebbert, J.O., Dale, L.C., Nirelli, L.M., Schroeder, D.R., Moyer, T.P. & Hurt, R.D. (2004) Cotinine as a biomarker of systemic nicotine exposure in spit tobacco users. *Addict. Behav.*, **29**, 349–355
- Eckfeldt, A., Hugoson, A., Bergendal, T. & Helkimo, M. (1990) An individual tooth wear index and an analysis of factors correlated to incisal and occlusal wear in an adult Swedish population. *Acta odontol. scand.*, **48**, 343–349
- Elbeshir, E.I., Abeen, H.A., Idris, A.M. & Abbas, K. (1989) Snuff dipping and oral cancer in Sudan: A retrospective study. *Br. J. oral maxillofac. Surg.*, **27**, 243–248
- Eliasson, B. (2003) Cigarette smoking and diabetes. *Prog. cardiovasc. Dis.*, **45**, 405–413
- Eliasson, M., Lundblad, D. & Hägg, E. (1991) Cardiovascular risk factors in young snuff-users and cigarette smokers. *J. intern. Med.*, **230**, 17–22
- Eliasson, M., Asplund, K., Evrin, P.-E. & Lundblad, D. (1995) Relationship of cigarette smoking and snuff dipping to plasma fibrinogen, fibrinolytic variables and serum insulin. The Northern Sweden MONICA study. *Atherosclerosis*, **113**, 41–53
- Eliasson, B., Taskinen, M.-R. & Smith, U. (1996) Long-term use of nicotine gum is associated with hyperinsulinemia and insulin resistance. *Circulation*, **94**, 878–881
- Eliasson, M., Asplund, K., Nasic, S. & Rodu, B. (2004) Influence of smoking and snus on the prevalence and incidence of type 2 diabetes amongst men: The northern Sweden MONICA study. *J. intern. Med.*, **256**, 101–110
- England, L.J., Levine, R.J., Mills, J.L., Klebanoff, M.A., Yu, K.F. & Cnattingius, S. (2003) Adverse pregnancy outcomes in snuff users. *Am. J. Obstet. Gynecol.*, **189**, 939–943
- Ernster, V.L., Grady, D.G., Greene, J.C., Walsh, M., Robertson, P., Daniels, T.E., Benowitz, N., Siegel, D., Gerbert, B. & Hauck, W.W. (1990) Smokeless tobacco use and health effects among baseball players. *J. Am. med. Assoc.*, **264**, 218–224
- European Parliament and Council (2001) Directive 2001/37/EC of the European Parliament and of the Council of 5 June 2001 on the approximation of the laws, regulations and administrative provisions of the Member States concerning the manufacture, presentation and sale of tobacco products. *Off. J. Eur. Commun.*, **L194**, 26–35

- Evstifeeva, T.V & Zaridze, D.G. (1992) Nass use, cigarette smoking, alcohol consumption and risk of oral and oesophageal precancer. *Eur. J. Cancer, Part B, Oral Oncol.*, **82**, 29–35
- Fagerström, K.-O. (1978) Measuring degree of physical dependence to tobacco smoking with reference to individualization of treatment. *Addict. Behav.*, **3**, 235–241
- Fakhfakh, R., Hsairi, M., Maalej, M., Achour, N. & Nacef, T. (2002) Tobacco use in Tunisia: Behaviour and awareness. *Bull. World Health Org.*, **80**, 350–356
- Fant, R.V., Henningfield, J.E., Nelson, R.A. & Pickworth, W.B. (1999) Pharmacokinetics and pharmacodynamics of moist snuff in humans. *Tob. Control*, **8**, 387–392
- Farrow, D.C. & Davis, S. (1990) Risk of pancreatic cancer in relation to medical history and the use of tobacco, alcohol and coffee. *Int. J. Cancer*, **45**, 816–820
- Federal Trade Commission (2003) *Smokeless Tobacco Report for the Years 2000 and 2001*, Washington DC [available at <http://www.ftc.gov/os/2003/08/2k2k1smokeless.pdf>; accessed 22 July 2004]
- Feinstein, A.R., Heinemann, L.A.J., Dalessio, D., Fox, J.M., Goldstein, J., Haag, G., Ladewig, D. & O'Brien, C.P. (2000) Do caffeine-containing analgesics promote dependence? A review and evaluation. *Clin. Pharmacol. Ther.*, **68**, 457–467
- Feyerabend, C., Ings, R.M. & Russell, M.A.H. (1985) Nicotine pharmacokinetics and its application to intake from smoking. *Br. J. clin. Pharmacol.*, **19**, 239–247
- Fiore, M.C., Novotny, T.E., Pierce, J.P., Giovino, G.A., Hatziandreu, E.J., Newcomb, P.A., Surawicz, T.S. & Davis, R.M. (1990) Methods used to quit smoking in the United States. Do cessation programs help? *J. Am. med. Assoc.*, **263**, 2760–2765
- Fiore, M.C., Bailey, W.C., Cohen, S.J., Dorfman, S.F., Goldstein, M.G., Gritz, E.R., Heyman, R.B., Jaén, C.R., Kottke, T.E., Lando, H.A., Mecklenburg, R.E., Mullen, P.D., Nett, L.M., Robinson, L., Stitzer, M.L., Tommasello, A.C., Villejo, L. & Wewers, M.E. (2000) *Treating Tobacco Use and Dependence (Quick Reference Guide for Clinicians)*, Rockville, MD, US Department of Health and Human Service, Public Health Service
- Fischer, S., Spiegelhalter, B., & Preussmann, R. (1989) Preformed tobacco-specific nitrosamines in tobacco — Role of nitrate and influence of tobacco type. *Carcinogenesis*, **10**, 1511–1517
- Fishman, J.A., Allison, H., Knowles, S.B., Fishburn, B.A., Woollery, T.A., Marx, W.T., Shelton, D.M., Husten, C.G. & Eriksen, M.P. (1999) State laws on tobacco control — United States, 1998. *MMWR CDC Surveill. Summ.*, **48**, 21–62
- Food and Drug Administration (1995) Regulations restricting sale and distribution of cigarettes and smokeless tobacco products to protect children and adolescents; final rule. *Fed. Reg.*, **60**, 41314–75, 41453–787
- Food and Drug Administration (1996) Regulations restricting sale and distribution of cigarettes and smokeless tobacco products to protect children and adolescents; final rule. *Fed. Reg.*, **61**, 44395–618
- Forey, B., Hamling, J., Lee, P. & Wald, N. (2002) *International Smoking Statistics: A Collection of Historical Data from 30 Economically Developed Countries*, 2nd Ed., London, Oxford University Press
- Forgas, L.B., Meyer, D.M. & Cohen, M.E. (1996) Tobacco use habits of naval personnel during Desert Storm. *Mil. Med.*, **161**, 165–168
- Foulds, J., Ramstrom, L., Burke, M. & Fagerström, K. (2003) Effect of smokeless tobacco (snus) on smoking and public health in Sweden. *Tob. Control*, **12**, 349–359

- Fox, J.W., Gyorfi, T. & Dragulev B. (1995) Effects of smokeless tobacco extract on HT 1080 cell adhesion, pp125^{FAK} phosphorylation and apoptosis. *In Vitro Toxicol.*, **8**, 339–347
- Franco, E.L., Kowalski, L.P., Oliveira, B.V., Curado, M.P., Pereira, R.N., Silva, M.E., Fava, A.S. & Torloni, H. (1989) Risk factors for oral cancer in Brazil: A case-control study. *Int. J. Cancer*, **43**, 992–1000
- Furie, M.B., Raffanello, J.A., Gergel, E.I., Lisinski, T.J. & Horb, L.D. (2000) Extracts of smokeless tobacco induce pro-inflammatory changes in cultured human vascular endothelial cells. *Immunopharmacology*, **47**, 13–23
- Galanti, M.R., Wickholm, S. & Gilljam, H. (2001a) Between harm and dangers. Oral snuff use, cigarette smoking and problem behaviours in a survey of Swedish male adolescents. *Eur. J. public Health*, **11**, 340–345
- Galanti, M.R., Rosendahl, I., Post, A. & Gilljam, H. (2001b) Early gender differences in adolescent tobacco use — The experience of a Swedish cohort. *Scand. J. public Health*, **29**, 314–317
- Galvin, R.J., Ramp, W.K. & Lenz, L.G. (1988) Smokeless tobacco contains a nonnicotine inhibitor of bone metabolism. *Toxicol. appl. Pharmacol.*, **95**, 292–300
- Galvin, R.J.S., Ramp, W.K. & Lenz, L.G. (1991) Comparison of the effects of smokeless tobacco extract with the effects of prolyl hydroxylase inhibitors on collagenous and noncollagenous protein synthesis by osteoblasts. *Connect. Tissue Res.*, **26**, 217–229
- Galvin, R.J.S., Ramp, W.K., Lenz, L.G. & Pierce, W.M., Jr (1992) Smokeless tobacco contains an inhibitor of prolyl hydroxylase activity. *Toxicol. Lett.*, **62**, 301–310
- Gan, C.-Y. (1995) Smokeless tobacco use among rural Kadazan women in Sabah, Malaysia. *S.E. Asian J. trop. Med. public Health*, **26**, 291–296
- Gan, C.-Y. (1998) Tobacco usage among rural Bajaus in Sabah, Malaysia. *S.E. Asian J. trop. Med. public Health*, **29**, 643–648
- Garner, W.W. (1951) *The Production of Tobacco*, New York, The Blakiston Company, pp. 477–478
- Ghose, U.R. & Parida, B.B. (1995) Cytological study of exfoliated buccal mucosa cells of the tribes in Orissa State (India) with high risk of oral cancer. *Indian J. Cancer*, **32**, 95–99
- Ghosh, S.K., Parikh, J.R., Gokani, V.N., Rao, N.M. & Doktor, P.B. (1985) Occupational health problems among tobacco processing workers: A preliminary study. *Arch. environ. Health*, **40**, 318–321
- Gijare, P.S., Rao, K.V.K. & Bhide, S.V. (1989) Effects of tobacco-specific nitrosamines and snuff extract on cell proliferation and activities of ornithine decarboxylase and aryl hydrocarbon hydroxylase in mouse tongue primary epithelial cell cultures. *J. Cancer Res. clin. Oncol.*, **115**, 558–563
- Gijare, P.S., Rao, K.V.K. & Bhide, S.V. (1990a) Modulatory effects of snuff, retinoic acid and β -carotene on DMBA-induced hamster cheek pouch carcinogenesis in relation to keratin expression. *Nutr. Cancer*, **14**, 253–259
- Gijare, P.S., Rao, K.V.K. & Bhide, S.V. (1990b) Inhibitory effects of snuff extract on ornithine decarboxylase and aryl hydrocarbon hydroxylase activities in relation to cell proliferation of mouse tongue epithelial cells. *Indian J. exp. Biol.*, **28**, 1012–1016
- Gillison, M.L., Koch, W.M., Capone, R.B., Spafford, M., Westra, W.H., Wu, L., Zahurak, M.L., Daniel, R.W., Viglione, M., Symer, D.E., Shah, K.V. & Sidransky, D. (2000) Evidence for a causal association between human papillomavirus and a subset of head and neck cancers. *J. natl Cancer Inst.*, **92**, 709–720

- Gilljam, H. & Galanti, M.R. (2003) Role of *snus* (oral moist snuff) in smoking cessation and smoking reduction in Sweden. *Addiction*, **98**, 1183–1189
- Giovino, G.A. (1999) Epidemiology of tobacco use among US adolescents. *Nicotine Tob. Res.*, **1**, S31–S40
- Giovino, G.A., Schooley, M.W., Zhu, B.P., Chrismon, J.H., Tomar, S.L., Peddicord, J.P., Merritt, R.K., Husten, C.G. & Eriksen, M.P. (1994) Surveillance for selected tobacco-use behaviors — United States, 1900–1994. *MMWR CDC Surveill. Summ.*, **43**, 1–43
- Gire, J.T. & Eissenberg, T. (2000) Placebo control study of acute smokeless tobacco abstinence in young adult men. *Psychol. addict. Behav.*, **14**, 356–366
- Global Youth Tobacco Survey Collaborative Group (2002) Tobacco use among youth: A cross country comparison. *Tob. Control*, **11**, 252–270
- Global Youth Tobacco Survey Collaborating Group (2003) Differences in worldwide tobacco use by gender: Findings from the Global Youth Tobacco Survey. *J. School Health*, **73**, 207–215
- Glover, E.D., Laflin, M. & Edwards, S.W. (1989) Age of initiation and switching patterns between smokeless tobacco and cigarettes among college students in the United States. *Am. J. public Health*, **79**, 207–208
- Glover, E.D., Wang, M.Q. & Glover, P.N. (1994) Development of a high school smokeless tobacco cessation manual. *Health Values*, **18**, 28–33
- Goodman, M.T., Morgenstern, H. & Wynder, E.L. (1986) A case–control study of factors affecting the development of renal cell cancer. *Am. J. Epidemiol.*, **124**, 926–941
- Gothia (2004) [<http://www.gothiatek.com/templates>; accessed 01 July 2004]
- Gothoskar, S.V., Sant, S.M. & Ranadive, K.J. (1975) Effect of tobacco and lime on oral mucosa of rats fed on vitamin B deficient diet. *Indian J. Cancer*, **12**, 424–429
- Gottsegen, J.J. (1940) *A Study of Its Consumption in the United States*, New York, Pitman, pp. 3
- Goud, M.L., Mohapatra, S.C., Mohapatra, P., Gaur, S.D., Pant, G.C. & Knanna, M.N. (1990) Epidemiological correlates between consumption of Indian chewing tobacco and oral cancer. *Eur. J. Epidemiol.*, **6**, 219–222
- Govekar, R.B. & Bhisey, R.A. (1992) Elevated urinary thioether excretion among bidi rollers exposed occupationally to processed tobacco. *Int. Arch. occup. environ. Health*, **64**, 101–104
- Govekar, R.B. & Bhisey, R.A. (1993) Mutagenic activity in urine samples from female tobacco habitues. *Cancer Lett.*, **69**, 75–80
- Government of Andhra Pradesh (2002) *Gazette No. 32*, Department of Health, Medical and Family Welfare
- Government of Bihar (2003) *Notification Vide 15/KH.A-24-6/02-387*, Department of Health
- Government of Goa (1997) *The Goa Prohibition of Smoking and Spitting Act, 1997* (Notification 7-21-97/LA), Saligao, Department of Science, Technology & Environment
- Government of Goa (2003) *The Gazette, Extraordinary No. 2 (DFDA/PFAA-121/98-/533)*, Directorate of Food & Drugs Administration
- Government of India (2001) *D.O.F. No. 334/1/2001-TRU*, Ministry of Finance, Department of Revenue [available at <http://indiabudget.nic.in/ub2001-02/cen/dojstru.pdf>; accessed 21 June 2004]
- Government of India (2003) *The Cigarettes and Other Tobacco Products (Prohibition of Advertisement and Regulation of Trade and Commerce, Production, Supply and Distribution) Act, 2003* (Gazette No. 34), Ministry of Law and Justice, 19 May 2003

- Government of Kerala (1999) *The Kerala Gazette No. 9*, Thiruvananthapuram, Department of Health Services
- Government of Maharashtra (2002) *Food (Health) Authority, Maharashtra State under Prevention of Food Adulteration Act, 1954* (No. PFA/Notifications/574-2002/7; The Maharashtra Gazette, No. 37/1954), Commissioner, Food and Drug Administration
- Government of Rajasthan (1950) *The Rajasthan Prevention of Juvenile Smoking Act, 1950*, 1 May 1950
- Government of Tamil Nadu (2001) *Gazette No. 780*, Director of Public Health & Preventive Medicine
- Government of Tamil Nadu (2003) *The Tamil Nadu Prohibition of Smoking and Spitting Act 2002*, Health and Family Welfare Department, 14 May 2003
- Government of West Bengal (2001) *The West Bengal Prohibition of Smoking and Spitting and Protection of Health of Non Smokers and Minors Bill, 2001* (Kolkata Gazette, Extraordinary No. WB (Part 4)/2001/SAR-37), 7 December 2001
- Grady, D., Greene, J., Daniels, T.E., Ernster, V.L., Robertson, P.B., Hauck, W., Greenspan, D., Greenspan, J. & Silverman, S., Jr (1990) Oral mucosal lesions found in smokeless tobacco users. *J. Am. dent. Assoc.*, **121**, 117–123
- Grady, D., Ernster, V.L., Stillman, L., Greene, J., Daniels, T.E. & Silverman, S., Jr (1991) Short term changes a surprise with smokeless tobacco oral lesions. *J. Am. dent. Assoc.*, **122**, 62–64
- Granberry, M.C., Smith, E.S., III, Troillet, R.D. & Eidt, J.F. (2003) Forearm endothelial response in smokeless tobacco users compared with cigarette smokers and nonusers of tobacco. *Pharmacotherapy*, **23**, 974–978
- Granero, R., Sánchez, M., Homes, G. & López, O. (2003) *Smokeless Tobacco Used Among Lara State, Venezuelan School Students. Global Youth Tobacco Survey (Gyts) (Abstract)*, 3rd Virtual Congress on Cardiology, Sept. 1–Dec. 30, 2003 [available at <http://www.fac.org.ar/tcvc/llave/tl071/htm>; accessed 8/12/2004]
- Grasser, J.A. & Childers, E. (1997) Prevalence of smokeless tobacco use and clinical oral leukoplakia in a military population. *Mil. Med.*, **162**, 401–404
- Grasso, P. & Mann, A.H. (1998) Smokeless tobacco and oral cancer: An assessment of evidence derived from laboratory animals. *Food chem. Toxicol.*, **36**, 1015–1029
- Green, G.A., Uryasz, F.D., Petr, T.A. & Bray, C.D. (2001) NCAA study of substance use and abuse habits of college student-athletes. *Clin. J. Sport Med.*, **11**, 51–56
- Greene, J.C., Ernster, V.L., Grady, D.G., Robertson, P.B., Walsh, M.M., Stillman, L.A. (1992) Oral mucosal lesions: Clinical findings in relation to smokeless tobacco use among US baseball players. In: *Smokeless Tobacco or Health: An International Perspective*, Vol. 2, Bethesda, MD, National Institutes of Health, pp. 41–50
- Greene, J.C., Walsh, M.M. & Letendre, M.A. (1998) Prevalence of spit tobacco use across studies of professional baseball players. *J. Calif. dent. Assoc.*, **26**, 358–364
- Greenland, S., Satterfield, M.H. & Lanes, S.F. (1998) A meta-analysis to assess the incidence of adverse effects associated with the transdermal nicotine patch. *Drug Saf.*, **18**, 297–308
- Greer, R.O. & Poulson, T.C. (1983) Oral tissue alterations associated with the use of smokeless tobacco by teen-agers. Part I. Clinical findings. *Oral Surg. oral Med. oral Pathol.*, **56**, 275–284

- Greer, R.O., Poulson, T.C., Boone, M.E., Lindenmuth, J.E. & Crosby, L. (1986) Smokeless tobacco associated oral changes in juvenile, adult and geriatric patients: Clinical and histomorphologic features. *Gerodontology*, **2**, 87–98
- Gritz, E.R., Baer-Weiss, V., Benowitz, N.L., Van Vunakis, H. & Jarvik, M.E. (1981) Plasma nicotine and cotinine concentrations in habitual smokeless tobacco users. *Clin. Pharmacol. Ther.*, **30**, 201–209
- Grunbaum, J.A., Kann, L., Kinchen, S., Ross, J., Hawkins, J., Lowry, R., Harris, W.A., McManus, T., Chyen, D. & Collins, J. (2004) Youth risk behavior surveillance — United States, 2003. *MMWR CDC Surveill. Summ.*, **53**, 1–96
- Gupta, P.C. (1992) Smokeless tobacco use in India. In: *Smokeless Tobacco or Health — An International Perspective* (Smoking and Tobacco Control Monograph 2; NIH Publication No. 92–3461), Bethesda, MD, National Institutes of Health, pp. 19–25
- Gupta, P.C. (1996) Survey of sociodemographic characteristics of tobacco use among 99 598 individuals in Bombay, India using handheld computers. *Tob. Control*, **5**, 114–120
- Gupta, P.C. (2000) Oral cancer and tobacco use in India: A new epidemic. In: *Tobacco: The Growing Epidemic. Proceedings of the 10th World Conference on Tobacco or Health, 24–28 August 1997, Beijing, China*
- Gupta, P.C. (2001) Tobacco Products Bill 2001: An aid to public health. *Natl med. J. India*, **14**, 131–134
- Gupta, P. (2004) *Laboratory Testing of Smokeless Tobacco Products. Final Report to the India Office of the WHO* (Allotment No.: SE IND TOB 001.RB.02), New Delhi
- Gupta, P.C. & Ray, C.S. (2002) Tobacco and youth in the South East Asian region. *Indian J. Cancer*, **39**, 5–33
- Gupta, P.C. & Ray, C.C. (2003) Smokeless tobacco and health in India and South Asia. *Respirology*, **8**, 419–431
- Gupta, I. & Sankar, D. (2004) Tobacco consumption in India: A new look using data from the National Sample Survey. *J. public Health Policy*, **24**, 233–245
- Gupta, P.C. & Sreevidya, S. (2004) Smokeless tobacco use, birth weight, and gestational age: Population based, prospective cohort study of 1217 women in Mumbai, India. *Br. med. J.* online: BMJ, doi:10.1136/bmj.38113.687882, EB
- Güven, A., Köksal, N., Büyükbese, M.A., Cetinkaya, A., Aksu, E. & Caglayan, C.E. (2003) Effects of using a different kind of smokeless tobacco on cardiac parameters: ‘Maras powder’. *Anadolu Kardiyol Derg*, **3**, 230–235
- Haddock, C.K., Vander Weg, M., DeBon, M., Klesges, R.C., Talcott, G.W., Lando, H. & Peterson, A. (2001) Evidence that smokeless tobacco use is a gateway for smoking initiation in young adult males. *Prev. Med.*, **32**, 262–267
- Hamazaki, Y. & Muraio, T. (1969) On the carcinogenic factors of tobacco leaf powder. I. Inhalation of tobacco leaf powder. *J. Karyopathol. esp. Tumor Tumovirus*, **12**, 163–178
- Hannam, S.E. (1997) Smokeless tobacco use among big ten wrestlers and selected associated factors. *Subst. Use Misuse*, **32**, 369–376
- Hannan, M.A., El-Yazigi, A., Paul, M., Gibson, D.P. & Phillips, R.L. (1986) Genotoxicity of ‘shamma’, a chewing material suspected of causing oral cancer in Saudi Arabia. *Mutat. Res.*, **169**, 41–46

- Hansson, L.-E., Baron, J., Nyrén, O., Bergström, R., Wolk, A. & Adami, H.-O. (1994) Tobacco, alcohol and the risk of gastric cancer. A population-based case-control study in Sweden. *Int. J. Cancer*, **57**, 26–31
- Harish, K. & Ravi, R. (1995) The role of tobacco in penile carcinoma. *Br. J. Urol.*, **75**, 375–377
- Harrison, D.F.N. (1964) Snuff — Its use and abuse. *Br. med. J.*, **ii**, 1649–1651
- Hartge, P., Hoover, R. & Kantor, A. (1985) Bladder cancer risk and pipes, cigars, and smokeless tobacco. *Cancer*, **55**, 901–906
- Hassoun, E.A., Bagchi, D., Bagchi, M. & Stohs, S.J. (1995) Effect of vitamin E succinate on smokeless tobacco-induced production of nitric oxide by rat peritoneal macrophages and J774A.1 macrophage cells in culture. *Free Radic. Biol. Med.*, **18**, 577–583
- Hatsukami, D.K. & Severson, H.H. (1999) Oral spit tobacco: Addiction, prevention and treatment. *Nicotine Tob. Res.*, **1**, 21–44
- Hatsukami, D.K., Gust, S.W. & Keenan, R.M. (1987) Physiologic and subjective changes from smokeless tobacco withdrawal. *Clin. Pharmacol. Ther.*, **41**, 103–107
- Hatsukami, D.K., Keenan, R.M. & Anton, D.J. (1988) Topographical features of smokeless tobacco use. *Psychopharmacology*, **96**, 428–429
- Hatsukami, D., Anton, D., Keenan, R. & Callies, A. (1992) Smokeless tobacco abstinence effects and nicotine gum dose. *Psychopharmacology*, **106**, 60–66
- Hatsukami, D., Jensen, J., Allen, S., Grillo, M. & Bliss, R. (1996) Effects of behavioral and pharmacological treatment on smokeless tobacco users. *J. consult. clin. Psychol.*, **64**, 153–161
- Hatsukami, D.K., Jensen, J., Boyle, R., Grillo, M. & Bliss, R. (1999) Characteristics of smokeless tobacco users seeking treatment. *Addict. Behav.*, **24**, 551–557
- Hatsukami, D.K., Lemmonds, C., Zhang, Y., Murphy, S.E., Le, C., Carmella, S.G. & Hecht, S.S. (2004) Evaluation of carcinogen exposure in people who used ‘reduced exposure’ tobacco products. *J. natl Cancer Inst.*, **96**, 844–852
- Hayes, R.B., Pottern, L.M., Swanson, G.M., Liff, J.M., Schoenberg, J.B., Greenberg, R.S., Schwartz, A.G., Brown, L.M., Silverman, D.T. & Hoover, R.N. (1994) Tobacco use and prostate cancer in blacks and whites in the United States. *Cancer Causes Control*, **5**, 221–226
- Hayes, J.D. & Pulford, D.J. (1995) The glutathione *S*-transferase supergene family: Regulation of GST* and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit. Rev. Biochem. mol. Biol.*, **30**, 445–600
- Health Canada (2000) *Tobacco Products Information Regulations, Regulation June 2000* (JUS-601413) [http://www.hc-sc.gc.ca/hl-vs/pubs/tobac-tabac/rc/index_e.html; accessed 22 July 2006]
- Health Canada (2001) *Regulations Amending the Tobacco Products Information Regulations, April 2001*, Ottawa, Tobacco Control Programme
- Heatherton, T.F., Kozlowski, L.T., Frecker, R.C. & Fagerström, K.-O. (1991) The Fagerström test for nicotine dependence: A revision of the Fagerström Tolerance Questionnaire. *Br. J. Addict.*, **86**, 1119–1127
- Hecht, S.S. (2002) Human urinary carcinogen metabolites: Biomarkers for investigating tobacco and cancer. *Carcinogenesis*, **23**, 907–922
- Hecht, S.S., Rivenson, A., Braley, J., DiBello, J., Adams, J.D. & Hoffmann, D. (1986) Induction of oral cavity tumors in F344 rats by tobacco-specific nitrosamines and snuff. *Cancer Res.*, **46**, 4162–4166

- Hecht, S.S., Carmella, S.G., Trushin, N., Foiles, P.G., Lin, D., Rubin, J.M. & Chung, F.-L. (1987) Investigations on the molecular dosimetry of tobacco-specific *N*-nitrosamines. In: Bartsch, H., O'Neill, I.K. & Schulte-Hermann, R., *The Relevance of N-Nitroso Compounds to Human Cancer: Exposures and Mechanisms* (IARC Scientific Publications No. 84), Lyon, pp. 423–429
- Hecht, S.S., Carmella, S.G. & Murphy, S.E. (1991) Hemoglobin adducts as biomarkers of exposure to and metabolic activation of carcinogenic tobacco-specific nitrosamines. *Biomed. environ. Sci.*, **4**, 93–103
- Hecht, S.S., Carmella, S.G. & Murphy, S.E. (1999a) Effects of watercress consumption on urinary metabolites of nicotine in smokers. *Cancer Epidemiol. Biomarkers Prev.*, **8**, 907–913
- Hecht, S.S., Carmella, S.G., Chen, M., Dor Koch, J.F., Miller, A.T., Murphy, S.E., Jensen, J.A., Zimmerman, C.L. & Hatsukami, D.K. (1999b) Quantitation of urinary metabolites of a tobacco-specific lung carcinogen after smoking cessation. *Cancer Res.*, **59**, 590–596
- Hecht, S.S., Carmella, S.G., Ye, M., Le, K.-A., Jensen, J.A., Zimmerman, C.L. & Hatsukami, D.K. (2002) Quantitation of metabolites of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone after cessation of smokeless tobacco use. *Cancer Res.*, **62**, 129–134
- Hecht, S.S., Murphy, S.E., Carmella, S.G., Li, S., Jensen, J., Le, C., Joseph, A.M. & Hatsukami, D.K. (2005) Similar uptake of lung carcinogens by smokers of regular, light, and ultralight cigarettes. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 693–698
- Heimann, R.K. (1960) *Tobacco and Americans*, New York, McGraw-Hill, pp. 8, 65, 118–119, 236
- Heineman, E.F., Hoar Zahm, S., McLaughlin, J.K. & Vaught, J.B. (1995) Increased risk of colorectal cancer among smokers: Results of a 26-year follow-up of US veterans and a review. *Int. J. Cancer*, **59**, 728–738
- Heinzel, P.A., Balaram, P. & Bernard, H.-U. (1996) Mutations and polymorphisms in the *p53*, *p21* and *p16* genes in oral carcinomas of Indian betel quid chewers. *Int. J. Cancer*, **68**, 420–423
- Helander, A. & Curvall, M. (1991) Comparison of blood aldehyde dehydrogenase activities in moist snuff users, cigarette smokers and nontobacco users. *Alcohol. clin. exp. Res.*, **15**, 1–6
- Henley, S.J., Thun, M.J., Connell, C. & Call, E.E. (2005) Two large prospective studies of mortality among men who use snuff or chewing tobacco (United States). *Cancer Causes Control*, **16**, 347–358
- Henningfield, J.E. & Fagerström, K.O. (2001) Swedish Match Company, Swedish snus and public health: A harm reduction experiment in progress? *Tob. Control*, **10**, 253–257
- Henningfield, J.E. & Fant, R.V. (1999) Tobacco use as drug addiction: The scientific foundation. *Nicotine Tob. Res.*, **1** (Suppl. 2), S31–S35
- Henningfield, J.E., Miyasato, K. & Jasinski, D.R. (1983) Cigarette smokers self-administer intravenous nicotine. *Pharmacol. biochem. Behav.*, **19**, 887–890
- Henningfield, J.E., Miyasato, K. & Jasinski, D.R. (1985) Abuse liability and pharmacodynamic characteristics of intravenous and inhaled nicotine. *J. Pharmacol. exp. Ther.*, **234**, 1–12
- Henningfield, J.E., Cohen, C. & Pickworth, W.B. (1993) Psychopharmacology of nicotine. In: Orleans, C.T. & Slade, J., eds, *Nicotine Addiction Principles and Management*, New York, Oxford University Press, pp. 24–45
- Henningfield, J.E., Radzius, A. & Cone, E.J. (1995) Estimation of available nicotine content of six smokeless tobacco products. *Tob. Control*, **4**, 57–61
- Henningfield, J.E., Fant, R.V. & Tomar, S.L. (1997) Smokeless tobacco: An addicting drug. *Adv. dent. Res.*, **11**, 330–335

- Henningfield, J.E., Benowitz, N.L., Slade, J., Houston, T.P., Davis, R.M. & Deitchman, S.D. for the Council on Scientific Affairs, American Medical Association (1998) Reducing the addictiveness of cigarettes. *Tob. Control*, **7**, 281–293
- Henningfield, J.E., Rose, C.A. & Giovino, G.A. (2002) Brave new world of tobacco disease prevention. Promoting dual tobacco-product use? *Am. J. prev. Med.*, **23**, 226–228
- Herrero, R., Castellsagué, X., Pawlita, M., Lissowska, J., Kee, F., Balaram, P., Rajkumar, T., Sridhar, H., Rose, B., Pintos, J., Fernández, L., Idris, A., Sánchez, M.J., Nieto, A., Talamini, R., Tavani, A., Bosch, F.X., Reidel, U., Snijders, P.J.F., Meijer, C.J.L.M., Viscidi, R., Muñoz, N. & Franceschi, S. for the IARC Multicenter Oral Cancer Study Group (2003) Human papillomavirus and oral cancer: The International Agency for Research on Cancer Multicenter Study. *J. natl Cancer Inst.*, **95**, 1772–1773
- Heuch, I., Kvåle, G., Jacobsen, B.K. & Bjelke, E. (1983) Use of alcohol, tobacco and coffee, and risk of pancreatic cancer. *Br. J. Cancer*, **48**, 637–643
- Hirsch, J.-M. & Johansson, S.L. (1983) Effect of long-term application of snuff on the oral mucosa — An experimental study in the rat. *J. oral Pathol.*, **12**, 187–198
- Hirsch, J.-M. & Thilander, H. (1981) Snuff-induced lesions of the oral mucosa — An experimental model in the rat. *J. oral Pathol.*, **10**, 342–353
- Hirsch, J.-M., Heyden, G. & Thilander, H. (1982) A clinical, histomorphological and histochemical study on snuff-induced lesions of varying severity. *J. oral Pathol.*, **11**, 387–398
- Hirsch, J.-M., Johansson, S.L. & Vahlne, A. (1984a) Effect of snuff and herpes simplex virus-1 on rat oral mucosa: Possible association with the development of squamous cell carcinoma. *J. oral Pathol.*, **13**, 52–62
- Hirsch, J.-M., Svennerholm, B. & Vahlne, A. (1984b) Inhibition of herpes simplex virus replication by tobacco extracts. *Cancer Res.*, **44**, 1991–1997
- Hirsch, J.M., Livian, G., Edward, S. & Noren, J.G. (1991) Tobacco habits among teenagers in the city of Göteborg, Sweden, and possible association with dental caries. *Swed. dent. J.*, **15**, 117–123
- Hirsch, J.-M., Hedner, J., Wernstedt, L., Lundberg, J. & Hedner, T. (1992) Hemodynamic effects of the use of oral snuff. *Clin. Pharmacol. Ther.*, **52**, 394–401
- Hoffmann, D. & Adams, J.D. (1981) Carcinogenic tobacco-specific *N*-nitrosamines in snuff and in the saliva of snuff dippers. *Cancer Res.*, **41**, 4305–4308
- Hoffmann, D. & Djordjevic, M.V. (1997) Chemical composition and carcinogenicity of smokeless tobacco. *Adv. dent. Res.*, **11**, 322–329
- Hoffmann, D., Harley, N.H., Fissene, I., Adams, J.D. & Brunnemann, K.D. (1986) Carcinogenic agents in snuff. *J. natl Cancer Inst.*, **76**, 435–437
- Hoffmann, D., Adams, J.D., Lisk, D., Fissenne, I. & Brunnemann, K.D. (1987) Toxic and carcinogenic agents in dry and moist snuff. *J. natl Cancer Inst.*, **79**, 1281–1286
- Hoffmann, D., Brunnemann, K.D. & Venitt, S. (1988) Carcinogenic nitrosamines in oral snuff. *Lancet*, **i**, 1232
- Hoffmann, D., Djordjevic, M.V. & Brunnemann, K.D. (1991) New brands of oral snuff. *Food chem. Toxicol.*, **29**, 65–68
- Hoffmann, D., Djordjevic, M.V., Fan, J., Zang, E., Glynn, T. & Connolly, G.N. (1995) Five leading U.S. commercial brands of moist snuff in 1994: Assessment of carcinogenic *N*-nitrosamines. *J. natl Cancer Inst.*, **87**, 1862–1869
- Hoffmann, D., Hoffmann, I. & El-Bayoumy, K. (2001) The less harmful cigarette: A controversial issue. A tribute to Ernst L. Wynder. *Chem. Res. Toxicol.*, **14**, 767–790

- Homburger, F. (1971) Mechanical irritation, polycyclic hydrocarbons, and snuff. Effects on facial skin, cheek pouch, and oral mucosa in Syrian hamsters. *Arch. Pathol.*, **91**, 411–417
- Homburger, F., Hsueh, S.-S., Russfield, A.B., Laird, C.W. & Van Dongen, C.G. (1976) Absence of carcinogenic effects of chronic feeding of snuff in inbred Syrian hamsters. *Toxicol. appl. Pharmacol.*, **35**, 515–521
- Hoover, J., McDermott, R. & Hartsfield, T. (1990) The prevalence of smokeless tobacco use in native children in northern Saskatchewan, Canada. *Can. J. public Health*, **81**, 350–352
- Hou-Jensen, K. (1964) On the occurrence of post nasal space tumours in Kenya. *Br. J. Cancer*, **18**, 58–68
- House of Representatives (1994) *Smokeless Tobacco Ingredient List as of April 4, 1994. U.S. House of Representatives Report to Subcommittee on Health and Environment, Committee on Energy and Commerce*, Washington DC, Patton, Boggs & Blow
- Howard, L.A., Miksys, S., Hoffmann, E., Mash, D. & Tyndale, R.F. (2003) Brain CYP2E1 is induced by nicotine and ethanol in rat and is higher in smokers and alcoholics. *Br. J. Pharmacol.*, **138**, 1376–1386
- Howe, G.R., Burch, J.D., Miller, A.B., Cook, G.M., Esteve, J., Morrison, B., Gordon, P., Chambers, L.W., Fodor, G. & Winsor, G.M. (1980) Tobacco use, occupation, coffee, various nutrients, and bladder cancer. *J. natl Cancer Inst.*, **64**, 701–713
- Hsairi, M., Achour, N., Zouari, B., Ben Romdhane, H., Achour, A., Maalej, M. & Nacef, T. (1993) Etiologic factors in primary bronchial carcinoma in Tunisia. *Tunis méd.*, **71**, 265–268
- Hsing, A.W., McLaughlin, J.K., Schuman, L.M., Bjelke, E., Gridley, G., Wacholder, S., Chien, H.T. & Blot, W.J. (1990) Diet, tobacco use, and fatal prostate cancer: Results from the Lutheran Brotherhood Cohort Study. *Cancer Res.*, **50**, 6836–6840
- Hsing, A.W., McLaughlin, J.K., Hrubec, Z., Blot, W.J. & Fraumeni, J.F., Jr (1991) Tobacco use and prostate cancer: 26-year follow-up of US veterans. *Am. J. Epidemiol.*, **133**, 437–441
- Huhtasaari, F., Asplund, K., Lundberg, V., Stegmayr, B. & Wester, P.O. (1992) Tobacco and myocardial infarction: Is snuff less dangerous than cigarettes? *Br. med. J.*, **305**, 1252–1256
- Huhtasaari, F., Lundberg, V., Eliasson, M., Janlert, U. & Asplund, K. (1999) Smokeless tobacco as a possible risk factor for myocardial infarction: A population-based study in middle-aged men. *J. Am. Coll. Cardiol.*, **34**, 1784–1790
- IARC (1985) *IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans*, Vol. 37, *Tobacco Habits Other than Smoking; Betel-quid and Areca-nut Chewing; and Some Related Nitrosamines*, Lyon
- IARC (1987) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Suppl. 7, *Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs Volumes 1 to 42*, Lyon, p. 68
- IARC (2004a) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 85, *Betel-quid and Areca-nut Chewing and Some Areca-nut Related Nitrosamines*, Lyon
- IARC (2004b) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 83, *Tobacco Smoke and Involuntary Smoking*, Lyon
- IARC (2007) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 90, *Human Papillomaviruses*, Lyon
- Ibrahim, E.M., Al Idrissi, H.Y., Magbool, G.M., Satti, M.B., Higazi, M.M. & Al Quorain, A. (1986) Oral cancer in Saudi Arabia: The role of Alqat and Alshammah. *Cancer Detect. Prev.*, **9**, 215–218

- Ibrahim, S.O., Johannessen, A.C., Idris, A.M., Hirsch, J.M., Vasstrand, E.N., Magnusson, B. & Nilsen, R. (1996) Immunohistochemical detection of p53 in non-malignant and malignant oral lesions associated with snuff dipping in the Sudan and Sweden. *Int. J. Cancer*, **68**, 749–753
- Ibrahim, S.O., Bertelsen, B., Kalvenes, M.B., Idris, A.M., Vasstrand, E.N., Nilsen, R. & Johannessen, A.C. (1998) Expression of keratin 13, 14 and 19 in oral squamous cell carcinomas from Sudanese snuff dippers: Lack of association with human papillomavirus infection. *APMIS*, **106**, 959–969
- Ibrahim, S.O., Vasstrand, E.N., Johannessen, A.C., Idris, A.M., Magnusson, B., Nilsen, R. & Lillehaug, J.R. (1999) Mutations of the *p53* gene in oral squamous-cell carcinomas from Sudanese dippers of nitrosamine-rich toombak and non-snuff-dippers from the Sudan and Scandinavia. *Int. J. Cancer*, **81**, 527–534
- Ibrahim, S.O., Lillehaug, J.R., Dolphine, O., Johnson, N.W., Warnakulasuriya, K.A.A.S. & Vasstrand, E.N. (2002) Mutations of the cell cycle arrest gene *p21^{WAF1}*, but not the metastasis-inducing gene S100A4, are frequent in oral squamous cell carcinomas from Sudanese *toombak* dippers and non-snuff-dippers from the Sudan, Scandinavia, USA and UK. *Anticancer Res.*, **22**, 1445–1451
- Idris, A.M., Nair, J., Ohshima, H., Friesen, M., Brouet, I., Faustman, E.M. & Bartsch, H. (1991) Unusually high levels of carcinogenic tobacco-specific nitrosamines in Sudan snuff (*toombak*). *Carcinogenesis*, **12**, 1115–1118
- Idris, A.M., Nair, J., Friesen, M., Ohshima, H., Brouet, I., Faustman, E.M. & Bartsch, H. (1992) Carcinogenic tobacco-specific nitrosamines are present at unusually high levels in the saliva of oral snuff users in Sudan. *Carcinogenesis*, **13**, 1001–1005
- Idris, A.M., Prokopczyk, B. & Hoffmann, D. (1994) *Toombak*: A major risk factor for cancer of the oral cavity in Sudan. *Prev. Med.*, **23**, 832–839
- Idris, A.M., Ahmed, H.M., Mukhtar B.I., Gadir, A.F. & el Beshir, E.I. (1995a) Descriptive epidemiology of oral neoplasms in Sudan, 1970–1985 and the role of *toombak*. *Int. J. Cancer*, **61**, 155–158
- Idris, A.M., Ahmed, H.M. & Malik, M.O.A. (1995b) *Toombak* dipping and cancer of the oral cavity in the Sudan: A case-control study. *Int. J. Cancer*, **63**, 477–480
- Idris, A.M., Warnakulasuriya, K.A.A.S., Ibrahim, Y.E., Nilsen, R., Cooper, D. & Johnson, N.W. (1996) *Toombak*-associated oral mucosa lesions in Sudanese show a low prevalence of epithelial dysplasia. *J. oral Pathol. Med.*, **25**, 239–244
- Idris, A.M., Ibrahim, S.O., Vasstrand, E.N., Johannessen, A.C., Lillehaug, J.R., Magnusson, B., Wallstrom, M., Hirsch, J.M. & Nilsen, R. (1998a) The Swedish snus and the Sudanese *toombak*: Are they different? *Oral Oncol.*, **34**, 558–566
- Idris, A.M., Ibrahim, Y.E., Warnakulasuriya, K.A.A.S., Cooper, D.J., Johnson, N.W. & Nilsen, R. (1998b) *Toombak* use and cigarette smoking in the Sudan: Estimates of prevalence in the Nile state. *Prev. Med.*, **27**, 597–603
- Idris, A.M., Warnakulasuriya, K.A.A.S., Ibrahim, Y.E., Hartley, R., Paterson, K., Patel, B., Nilsen, R. & Johnson, N.W. (1998c) Characterization of an amorphous deposit in the lamina propria in oral snuff users in the Sudan as collagen. *J. oral Pathol. Med.*, **27**, 157–162
- Institute of Medicine (2001) *Clearing the Smoke: Assessing the Science Base for Tobacco Harm Reduction*, Washington DC, National Academy Press [<http://www.nap.edu/books/0309072824/html/>]

- Islam, N. & Al-Khateeb, M. (1995) Challenges and opportunities for tobacco control in Islamic countries. A case-control study. *Eastern med. Health J.*, **1**, 230–234
- Jacob, P., III, Hatsukami, D., Severson, H., Hall, S., Yu, L. & Benowitz, N.L. (2002) Anabasine and anatabine as biomarkers for tobacco use during nicotine replacement therapy. *Cancer Epidemiol. Biomarkers Prev.*, **11**, 1668–1673
- Jacob, B.J., Straif, K., Thomas, G., Ramadas, K., Mathew, B., Zhang, Z.-F., Sankaranarayanan, R. & Hashibe, M. (2004) Betel quid without tobacco as a risk factor for oral precancers. *Oral Oncol.*, **40**, 697–704
- Jafarey, N.A., Mahmood, Z. & Zaidi, S.H.M. (1977) Habits and dietary pattern of cases of carcinoma of the oral cavity and oropharynx. *J. Pakistan med. Assoc.*, **27**, 340–343
- Jaffee, S.M. (2003) *Malawi's Tobacco Sector: Standing on One Strong Leg is Better Than on None* (Africa Region Working Paper Series No. 55) [<http://www.worldbank.org/afr/wps/index.htm>; accessed 01 August 2004]
- Jansson, T., Romert, L., Magnusson, J. & Jenssen, D. (1991) Genotoxicity testing of extracts of a Swedish moist oral snuff. *Mutat. Res.*, **261**, 101–115
- Jasinski, D.R., Johnson, R.E. & Henningfield, J.E. (1984) Abuse liability assessment in human subjects. *Trends pharmacol. Sci.*, **5**, 196–200
- Jenson, H.B., Heard, P. & Moyer, M.P. (1999a) Evaluation of the effect of smokeless tobacco purified products and extracts on latent Epstein-Barr virus. *Toxicology*, **133**, 35–42
- Jenson, H.B., Baillargeon, J., Heard, P. & Moyer, M.P. (1999b) Effects of smokeless tobacco and tumor promoters on cell population growth and apoptosis of B lymphocytes infected with Epstein-Barr virus types 1 and 2. *Toxicol. appl. Pharmacol.*, **160**, 171–182
- Johansson, S.L., Hirsch, J.M., Larsson, P.-A., Saidi, J. & Österdahl, B.-G. (1989) Snuff-induced carcinogenesis: Effect of snuff in rats initiated with 4-nitroquinoline *N*-oxide. *Cancer Res.*, **49**, 3063–3069
- Johansson, S.L., Saidi, J., Österdahl, B.-G. & Smith, R.A. (1991) Promoting effect of snuff in rats initiated by 4-nitroquinoline-*N*-oxide or 7,12-dimethylbenz(a)anthracene. *Cancer Res.*, **51**, 4388–4394
- Johnson, G.K., Poore, T.K., Squier, C.A., Wertz, P.W., Reinhardt, R.A. & Vincent, S.D. (1994) Prostaglandin E₂ and interleukin-1 levels in smokeless tobacco induced oral mucosal lesions. *J. Periodontol.*, **29**, 430–438
- Johnson, G.K., Poore, T.K., Payne, J.B. & Organ, C.C. (1996) Effect of smokeless tobacco extract on human gingival keratinocyte levels of prostaglandin E₂ and interleukin-1. *J. Periodontol.*, **67**, 116–124
- Johnson, G.K., Payne, J.B., Fili, J.M., Reinhardt, R.A., Organ, C.C. & Slager, S.L. (1998) Development of smokeless tobacco-induced oral mucosal lesions. *J. oral Pathol. Med.*, **27**, 388–394
- Johnston, L.D., O'Malley, P.M., Bachman, J.G. & Schulenberg, J.E. (2003) *Monitoring the Future — National Survey Results on Drug Use, 1975–2002*, Vol. I, *Secondary School Students* (NIH Publication No. 04-5507), Bethesda, MD, National Institute on Drug Abuse [available at http://monitoringthefuture.org/pubs/monographs/vol1_2002.pdf; accessed 23 July 2004]
- Jones, R.B. & Moberg, D.P. (1988) Correlates of smokeless tobacco use in a male adolescent population. *Am. J. public Health*, **78**, 61–63
- Joseph, A.M., Norman, S.M., Ferry, L.H., Prochazka, A.V., Westman, E.C., Steel, B.G., Sherman, S.E., Cleveland, M., Antonuccio, D.O., Hartman, N. & McGovern, P.G. (1996) The safety of

- transdermal nicotine as an aid to smoking cessation in patients with cardiac disease. *New Engl. J. Med.*, **335**, 1792–1798
- Jungell, P. & Malmström, M. (1985) Snuff-induced lesions in Finnish recruits. *Scand. J. dent. Res.*, **93**, 442–447
- Kabat, G.C., Chang, C.J. & Wynder, E.L. (1994) The role of tobacco, alcohol use, and body mass index in oral and pharyngeal cancer. *Int. J. Epidemiol.*, **23**, 1137–1144
- Kandarkar, S.V., Hasegkar, N.N. & Sirsat, S.M. (1981) Optical and ultrastructural pathology of vitamin A pretreated hamster cheek pouch exposed to lime (Ca(OH)₂) and tobacco over total life span. *Neoplasma*, **28**, 729–737
- Kann, L., Warren, C.W., Harris, W.A., Collins, J.L., Douglas, K.A., Collins, M.E., Williams, B.I., Ross, J.G. & Kolbe, L.J. (1995) Youth risk behavior surveillance — United States, 1993. *MMWR CDC Surveill. Summ.*, **44**, 1–55
- Kannan, K., Munirajan, A.K., Krishnamurthy, J., Bhuvaramurthy, V., Mohanprasad, B.K., Panishankar, K.H., Tsuchida, N. & Shanmugam, G. (1999) Low incidence of *p53* mutations in betel quid and tobacco chewing-associated oral squamous carcinoma from India. *Int. J. Oncol.*, **15**, 1133–1136
- Kannan, K., Munirajan, A.K., Bhuvaramurthy, V., Mohanprasad, B.K.C., Shankar, P., Tsuchida, N. & Shanmugam, G. (2000) *FHIT* gene mutations and single nucleotide polymorphism in Indian oral and cervical squamous cell carcinomas. *Oral Oncol.*, **36**, 189–193
- Kao, T.C., Schneider, S.J. & Hoffman, K.J. (2000) Co-occurrence of alcohol, smokeless tobacco, cigarette, and illicit drug use by lower ranking military personnel. *Addict. Behav.*, **25**, 253–262
- Kaplan, M., Carriker, L. & Waldron, I. (1990) Gender differences in tobacco use in Kenya. *Soc. Sci. Med.*, **30**, 305–310
- Karki, Y.B., Pant, K.D. & Pande, B.R. (2003) *A Study on the Economics of Tobacco in Nepal* (HNP Discussion Paper, Economics of Tobacco Control Paper No. 13), Geneva, WHO, Tobacco Free Initiative
- Karvonen, J.S., Rimpelä, A.H. & Rimpelä, M.K. (1993) Modernization and smoking —Regional differences in adolescent smoking in Finland in 1977–1987. *Scand. J. soc. Med.*, **21**, 188–196
- Kaugars, G.E., Mehailescu, W.L. & Gunsolley, J.C. (1989) Smokeless tobacco use and oral epithelial dysplasia. *Cancer*, **64**, 1527–1530
- Kaugars, G.E., Riley, W.T., Brandt, R.B., Burns, J.C. & Svirsky, J.A. (1992) The prevalence of oral lesions in smokeless tobacco users and an evaluation of risk factors. *Cancer*, **70**, 2579–2585
- Kaur, J. & Ralhan, R. (2003) Establishment and characterization of a cell line from smokeless tobacco associated oral squamous cell carcinoma. *Oral Oncol.*, **39**, 806–820
- Kaur, J., Srivastava, A. & Ralhan, R. (1994) Overexpression of p53 protein in betel- and tobacco-related human oral dysplasia and squamous-cell carcinoma in India. *Int. J. Cancer*, **58**, 340–345
- Kaur, J., Srivastava, A. & Ralhan, R. (1998) Prognostic significance of p53 protein overexpression in betel- and tobacco-related oral oncogenesis. *Int. J. Cancer*, **79**, 370–375
- Kayal, J.J., Trivedi, A.H., Dave, B.J., Nair, J., Nair, U.J., Bhide, S.V., Goswami, U.C. & Adhvaryu, S.G. (1993) Incidence of micronuclei in oral mucosa of users of tobacco products singly or in various combinations. *Mutagenesis*, **8**, 31–33
- Keen, P., De Moor, N.G., Shapiro, M.P., Cooper, R.L., Campbell, J.M. & Cohen, L. (1955) The aetiology of respiratory tract cancer in the South African Bantu. Part I. Clinical aspects. *Br. J. Cancer*, **9**, 528–533

- Keenan, R.M., Hatsukami, D.K. & Anton, D.J. (1989) The effects of short-term smokeless tobacco deprivation on performance. *Psychopharmacology*, **98**, 126–130
- Kenny, K.K., Quigley, N.C. & Regennitter, F.J. (1996) Survey of smokeless tobacco use in basic trainees and armor basic course officers. *Mil. Med.*, **161**, 37–42
- Kessler, D.A., Witt, A.M., Barnett, P.S., Zeller, M.R., Natanblut, S.L., Wilkenfeld, J.P., Lorraine, C.C., Thompson, L.J. & Schultz, W.B. (1996) The Food and Drug Administration's regulation of tobacco products. *New Engl. J. Med.*, **335**, 988–994
- Khurana, M., Sharma, D. & Khandelwal, P.D. (2000) Lipid profile in smokers and tobacco chewers — A comparative study. *J. Assoc. Physic. India*, **48**, 895–897
- Kilaru, S., Frangos, S.G., Chen, A.H., Gortler, D., Dhadwal, A.K., Aram, O. & Sumpio, B.E. (2001) Nicotine: A review of its role in atherosclerosis. *J. Am. Coll. Surg.*, **193**, 538–546
- Kiseleva, N.S., Milievskaia, I.L. & Caklin, A.V. (1976) Development of tumours in Syrian hamsters during prolonged experimental exposure to nass. *Bull. World Health Org.*, **54**, 597–605
- Kneller, R.W., McLaughlin, J.K., Bjelke, E., Schuman, L.M., Blot, W.J., Wacholder, S., Gridley, G., CoChien, H.T. & Fraumeni, J.F., Jr (1991) A cohort study of stomach cancer in a high-risk American population. *Cancer*, **68**, 672–678
- Kozlowski, L.T. (1981) The determinants of tobacco use: Cigarette smoking in the context of other forms of tobacco use. *Can. J. public Health*, **72**, 396–401
- Kozlowski, L.T., O'Connor, R.J., Edwards, B.Q. & Flaherty, B.P. (2003) Most smokeless tobacco use is not a causal gateway to cigarettes: Using order of product use to evaluate causation in a national US sample. *Addiction*, **98**, 1077–1085
- Kraft, P. & Svendsen, T. (1997) Tobacco use among young adults in Norway, 1973–95: Has the decrease levelled out? *Tob. Control*, **6**, 27–32
- Kreimer, A.R., Clifford, G.M., Boyle, P. & Franceschi, S. (2005) Human papillomavirus types in head and neck squamous cell carcinomas worldwide: A systematic review. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 467–475
- Kresty, L.A., Carmella, S.G., Borukhova, A., Akerkar, S.A., Gopalakrishnan, R., Harris, R.E., Stoner, G.D. & Hecht, S.S. (1996) Metabolites of a tobacco-specific nitrosamine, 4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone (NNK), in the urine of smokeless tobacco users: Relationship between urinary biomarkers and oral leukoplakia. *Cancer Epidemiol. Biomarkers Prev.*, **5**, 521–525
- Krishna, K. (1978) Tobacco chewing in pregnancy. *Br. J. Obstet. Gynaecol.*, **85**, 726–728
- Krishnamurthy, S. (1991) Strength of association of increased placental weight and smokeless tobacco use in pregnancy. *Indian J. Pediatr.*, **58**, 863–876
- Krishnamurthy, S., Ramaswamy, R., Trivedi, U. & Zachariah, V. (1997) Tobacco use in rural Indian children. *Indian Pediatr.*, **34**, 923–927
- Krishnamurthy, J., Kannan, K., Feng, J., Mohanprasad, B.K.C., Tsuchida, N. & Shanmugam, G. (2001) Mutational analysis of the candidate tumor suppressor gene *ING1* in Indian oral squamous cell carcinoma. *Oral Oncol.*, **37**, 222–224
- Ksir, C., Shank, M., Kraemer, W. & Noble, B. (1986) Effects of chewing tobacco on heart rate and blood pressure during exercise. *J. Sports Med.*, **26**, 384–389
- Kulkarni, V. & Saranath, D. (2004) Concurrent hypermethylation of multiple regulatory genes in chewing tobacco associated oral squamous cell carcinomas and adjacent normal tissues. *Oral Oncol.*, **40**, 145–153

- Kulkarni, J.R., Lalitha, V.S. & Bhide, S.V. (1988) Carcinogenicity studies of masher, a pyrolysed product of tobacco. *Carcinogenesis*, **9**, 2137–2138
- Kulkarni, J., Lalitha, V.S. & Bhide, S. (1989) Weak carcinogenic effect of masher, a pyrolysed tobacco product, in mouse skin tumor models. *J. Cancer Res. clin. Oncol.*, **115**, 166–169
- Kuttan, N.A.A., Rosin, M.P., Ambika, K., Priddy, R.W., Bhakthan, N.M.G. & Zhang, L. (1995) High prevalence of expression of p53 oncoprotein in oral carcinomas from India associated with betel and tobacco chewing. *Eur. J. Cancer, Part B, Oral Oncol.*, **31**, 169–173
- Kyaing, N.N. (2005) *Myanmar 2004, Global Youth Tobacco Survey (GYTS) — Fact Sheet* [available at http://www.cdc.gov/tobacco/global/GYTS/factsheets/2004/pdf/Myanmar_factsheet2004.pdf; accessed 01 June 2006]
- Lagergren, J., Bergström, R., Lindgren, A. & Nyrén, O. (2000) The role of tobacco, snuff and alcohol use in the aetiology of cancer of the oesophagus and gastric cardia. *Int. J. Cancer*, **85**, 340–346
- Lambe, M. (2004) Tobacco control, ethics and Swedish snus. *Acta oncol.*, **43**, 3–4
- Landmesser, U., Hornig, B. & Drexler, H. (2004) Endothelial function: A critical determinant in atherosclerosis? *Circulation*, **109** (Suppl. II), II-27–II-33
- Lanier, A.P., Bulkow, L.R., Novotny, T.E., Giovino, G.A. & Davis, R.M. (1990) Tobacco use and its consequences in northern populations. *Arctic med. Res.*, **49** (Suppl 2), 17–22
- Larsson, A., Axéll, T. & Andersson, G. (1991) Reversibility of snuff dippers' lesion in Swedish moist snuff users: A clinical and histologic follow-up study. *J. oral Pathol. Med.*, **20**, 258–264
- Larsson, P.-A., Hirsch, J.M., Gronowitz, J.S. & Vahlne, A. (1992) Inhibition of herpes simplex virus replication and protein synthesis by non-smoked tobacco, tobacco alkaloids and nitrosamines. *Arch. oral Biol.*, **37**, 969–978
- LaVoie, E.J., Tucciarone, P., Kagan, M., Adams, J.D. & Hoffmann, D. (1989) Analyses of steam distillates and aqueous extracts of smokeless tobacco. *J. agric. Food Chem.*, **37**, 154–157
- Lazarus, P., Garewal, H.S., Sciubba, J., Zwiebel, N., Calcagnotto, A., Fair, A., Schaefer, S. & Richie, J.P., Jr (1995) A low incidence of p53 mutations in pre-malignant lesions of the oral cavity from non-tobacco users. *Int. J. Cancer*, **60**, 458–463
- Lazarus, P., Stern, J., Zwiebel, N., Fair, A., Richie, J.P., Jr & Schantz, S. (1996a) Relationship between p53 mutation incidence in oral cavity squamous cell carcinomas and patient tobacco use. *Carcinogenesis*, **17**, 733–739
- Lazarus, P., Idris, A.M., Kim, J., Calcagnotto, A. & Hoffmann, D. (1996b) p53 Mutations in head and neck squamous cell carcinomas from Sudanese snuff (toombak) users. *Cancer Detect. Prev.*, **20**, 270–278
- Lee, S., Raker, T. & Chisick, M.C. (1994) Psychosocial factors influencing smokeless tobacco use by teen-age military dependents. *Mil. Med.*, **159**, 112–117
- Lee, J.J., Hong, W.K., Hittelman, W.N., Mao, L., Lotan, R., Shin, D.M., Benner, S.E., Xu, X.-C., Lee, J.S., Papadimitrakopoulou, V.M., Geyer, C., Perez, C., Martin, J.W., El-Naggar, A.K. & Lippman, S.M. (2000) Predicting cancer development in oral leukoplakia: Ten years of translational research. *Clin. Cancer Res.*, **6**, 1702–1710
- Lemmonds, C.A., Hecht, S.S., Jensen, J.A., Murphy, S.E., Carmella, S.G., Zhang, Y. & Hatsukami, D.K. (2005) Smokeless tobacco topography and toxin exposure. *Nicotine Tob. Res.*, **7**, 469–474
- Lenz, L.G., Ramp, W.K., Galvin, R.J.S. & Pierce, W.M., Jr (1992) Inhibition of cell metabolism by a smokeless tobacco extract: Tissue and species specificity. *Proc. Soc. exp. Biol. Med.*, **199**, 211–217

- Levenson-Gingiss, P. & Gottlieb, N.H. (1991) A comparison of smokeless tobacco and smoking practices of university varsity and intramural baseball players. *Addict. Behav.*, **16**, 335–340
- Levenson-Gingiss, P., Morrow, J.R., Jr & Dratt, L.M. (1989) Patterns of smokeless tobacco use among university athletes. *J. Am. Coll. Health*, **38**, 87–90
- Lewin, F., Norell, S.E., Johansson, H., Gustavsson, P., Wennerberg, J., Biörklund, A. & Rutqvist, L.E. (1998) Smoking tobacco, oral snuff, and alcohol in the etiology of squamous cell carcinoma of the head and neck: A population-based case-referent study in Sweden. *Cancer*, **82**, 1367–1375
- Li, X. & Bush, L.P. (2004) TSNA changes in cured powdered tobacco in short-time periods (Abstract No. 30). In: *58th Tobacco Science Research Conference Proceedings*, Winston-Salem, NC, Tobacco Science Research
- Lichtenstein, E., Severson, H.H., Friedman, L.S. & Ary, D.V. (1984) Chewing tobacco use by adolescents: Prevalence and relation to cigarette smoking. *Addict. Behav.*, **9**, 351–355
- Lindström, M., Isacson, S.-O. & The Malmö Shoulder-Neck Study Group (2002a) Smoking cessation among daily smokers, aged 45–69 years: A longitudinal study in Malmö, Sweden. *Addiction*, **97**, 205–215
- Lindström, M., Isacson, S.-O. & The Malmö Shoulder-Neck Study Group (2002b) Long term and transitional intermittent smokers: A longitudinal study. *Tob. Control*, **11**, 61–67
- Link, J.O., Kaugars, G.E. & Burns, J.C. (1992) Comparison of oral carcinomas in smokeless tobacco users and nonusers. *J. oral maxillofac. Surg.*, **50**, 452–455
- Little, S.J., Stevens, V.J., LaChance, P.A., Severson, H.H., Bartley, M.H., Lichtenstein, E. & Leben, J.R. (1992) Smokeless tobacco habits and oral mucosal lesions in dental patients. *J. public Health Dent.*, **52**, 269–276
- Liu, Y., Egyhazi, S., Hansson, J., Bhide, S.V., Kulkarni, P.S. & Grafström, R.C. (1997) O⁶-Methylguanine–DNA methyltransferase activity in human buccal mucosal tissue and cell cultures. Complex mixtures related to habitual use of tobacco and betel quid inhibit the activity *in vitro*. *Carcinogenesis*, **18**, 1889–1895
- Livingston, G.K., Reed, R.N., Olson, B.L. & Lockey, J.E. (1990) Induction of nuclear aberrations by smokeless tobacco in epithelial cells of human oral mucosa. *Environ. mol. Mutag.*, **15**, 136–144
- Loewe, W. (1981) [Snuff in Gothenburg]. In: *Svenska Tobaks AB 1981*, Stockholm, pp. 19–22 (in Swedish)
- Loro, L.L., Vintermyr, O.K., Ibrahim, S.O., Idris, A.M. & Johannessen, A.C. (2000) Apoptosis and expression of Bax and Bcl-2 in snuff- and non-snuff associated oral squamous cell carcinomas. *Anticancer Res.*, **20**, 2855–2860
- Luomanen, M., Tiitta, O., Heikinheimo, K., Heinaro, I. & Happonen, R.-P. (1997a) Effect of snuff on cytokeratin expression in oral vestibular sulcus epithelium. *J. oral Pathol. Med.*, **26**, 110–116
- Luomanen, M., Tiitta, O., Heikinheimo, K., Leimola-Virtanen, R., Heinaro, I. & Happonen, R.-P. (1997b) Effect of snuff and smoking on tenascin expression in oral mucosa. *J. oral Pathol. Med.*, **26**, 334–338
- Lyon, H., Poulsen, H.E. & Pindborg, J.J. (1964) Studies on oral leukoplakias: 6. Deposits of amyloid in the oral submucosa induced by prolonged use of snuff. *Acta pathol. microbiol. scand.*, **60**, 305–310

- Maden, C., Beckmann, A.M., Thomas, D.B., McKnight, B., Sherman, K.J., Ashley, R.L., Corey, L. & Daling, J.R. (1992) Human papillomaviruses, herpes simplex viruses, and the risk of oral cancer in men. *Am. J. Epidemiol.*, **135**, 1093–1102
- Mahale, A. & Saranath, D. (2000) Microsatellite alterations on chromosome 9 in chewing tobacco-induced oral squamous cell carcinomas from India. *Oral Oncol.*, **36**, 199–206
- Mahanta, J., Chetia, M., Hazarika, N.C., Narain, K. & Sharma, S.K. (1998) Toxicity of *tuibur*, a unique form of tobacco smoke extract in Mizoram, India. *Curr. Sci.*, **75**, 381–384
- Mahimkar, M.B., Buch, S.C., Samant, T.A., Kapoor, M.D. & Bhisey, R.A. (2001) Influence of smokeless tobacco exposure on detoxification status and chromosomal damage in male and female habitues. *Mutat. Res.*, **491**, 111–117
- Mahmood, Z. (1982) Smoking and chewing habits of people of Karachi — 1981. *J. Pakistan med. Assoc.*, **32**, 34–37
- Mahmood, Z., Jafarey, N.A., Samiuddin, M., Rais, A.K., Malik, S. & Qureshi, S.A. (1974) Dietary and other habits of the people of Karachi. *J. Pakistan med. Assoc.*, **24**, 222–229
- Mangipudy, R.S. & Vishwanatha, J.K. (1999) Role of nitric oxide in the induction of apoptosis by smokeless tobacco extract. *Mol. cell. Biochem.*, **200**, 51–57
- Mani, N.J. (1985) Preliminary report on prevalence of oral cancer and precancerous lesions among dental patients in Saudi Arabia. *Community Dent. oral Epidemiol.*, **13**, 247–248
- Marshall, J.R., Graham, S., Haughey, B.P., Shedd, D., O'Shea, R., Brasure, J., Wilkinson, G.S. & West, D. (1992) Smoking, alcohol, dentition and diet in the epidemiology of oral cancer. *Oral. Oncol. Eur. J. Cancer*, **28B**, 9–15
- Martin, G.C., Brown, J.P., Eifler, C.W. & Houston, G.D. (1999) Oral leukoplakia status six weeks after cessation of smokeless tobacco use. *J. Am. dent. Assoc.*, **130**, 945–954
- Mashberg, A., Boffetta, P., Winkelman, R. & Garfinkel, L. (1993) Tobacco smoking, alcohol drinking, and cancer of the oral cavity and oropharynx among U.S. veterans. *Cancer*, **72**, 1369–1375
- Masironi, R. (1992) WHO strategies to curb smokeless tobacco: A global perspective. In: *Smokeless Tobacco or Health: An International Perspective* (Smoking and Tobacco Control Monographs No. 2), Geneva, WHO, pp. 315–322
- Mavropoulos, A., Aars, H. & Brodin, P. (2001) The acute effects of smokeless tobacco (snuff) on gingival blood flow in man. *J. periodont. Res.*, **36**, 221–226
- Maxwell, J.C., Jr (1980) Chewing, snuff is growth segment. *Tob. Rep.*, **107**, 32–35
- Maxwell Tobacco Fact Book (2002) *Appendix 1. Manufacturing (Tables 12 and 14) Tobacco Reporter* (www.tobaccoreporter.com)
- Mazière, M., Comar, D., Marzano, C. & Berger, G. (1976) Nicotine-¹¹C: Synthesis and distribution kinetics in animals. *Eur. J. nucl. Med.*, **1**, 255–258
- McLaughlin, J.K., Lindblad, P., Mellemegaard, A., McCredie, M., Mandel, J.S., Schlehofer, B., Pommer, W. & Adami, H.-O. (1995) International renal-cell cancer study. I. Tobacco use. *Int. J. Cancer*, **60**, 194–198
- MDPH (Massachusetts Department of Public Health) (2004) *Smokeless Tobacco Data Base*, Boston, MA
- Meger, M., Megr-Kossien, I., Schuler-Metz, A., Janket, D. & Scherer, G. (2002) Simultaneous determination of nicotine and eight nicotine metabolites in urine of smokers using liquid chromatography–tandem mass spectrometry. *J. Chromatogr.*, **B778**, 251–261

- Mehta, F.S., Pindborg, J.J., Gupta, P.C. & Daftary, D.K. (1969) Epidemiologic and histologic study of oral cancer and leukoplakia among 50 915 villagers in India. *Cancer*, **24**, 832–849
- Mehta, F.S., Gupta, P.C., Daftary, D.K., Pindborg, J.J. & Choksi, S.K. (1972) An epidemiologic study of oral cancer and precancerous conditions among 101 761 villagers in Maharashtra, India. *Int. J. Cancer*, **10**, 134–141
- Melnick, M.J., Miller, K.E., Sabo, D.F., Farrell, M.P. & Barnes, G.M. (2001) Tobacco use among high school athletes and nonathletes: Results of the 1997 youth risk behavior survey. *Adolescence*, **36**, 727–747
- Mengesha, Y.A. & Bekele, A. (1998) Relative chronic effects of different occupational dusts on respiratory indices and health of workers in three Ethiopian factories. *Am. J. ind. Med.*, **34**, 373–380
- Merchant, A., Husain, S.S.M., Hosain, M., Fikree, F.F., Pitiphat, W., Siddiqui, A.R., Hayder, S.J., Haider, S.M., Ikram, M., Chuang, S.K. & Saeed, S.A. (2000) Paan without tobacco. An independent risk factor for oral cancer. *Int. J. Cancer*, **86**, 128–131
- Merne, M.E.T., Tiekso, J.T. & Syrj nen, S.M. (1998) Snuff use and smoking among senior high school students: Effects of a snuff sales ban. *Oral Dis.*, **4**, 207–212
- Merne, M., Heinaro, I., L hteenoja, H. & Syrj nen, S. (2002) Proliferation and differentiation markers in snuff-induced oral mucosal lesions. *J. oral Pathol. Med.*, **31**, 259–266
- Merne, M., Heikinheimo, K., Saloniemi, I. & Syrj nen, S. (2004) Effects of snuff extract on epithelial growth and differentiation *in vitro*. *Oral Oncol.*, **40**, 6–12
- Miksys, S., Lerman, C., Shields, P.G., Mash, D.C. & Tyndale, R.F. (2003) Smoking, alcoholism and genetic polymorphisms after CYP2B6 levels in human brain. *Neuropharmacology*, **45**, 122–132
- Milievskaja, I.L. & Kiseleva, N.S. (1976) Comparative study of the carcinogenic activities of nass and some chemical carcinogens when introduced into the buccal pouch of the Syrian hamster. *Bull. World Health Org.*, **54**, 607–614
- Millar, W.J. (1989) The use of chewing tobacco and snuff in Canada, 1986. *Can. J. public Health*, **80**, 131–135
- Millar, W.J. (1990) Smokeless tobacco use by youth in the Canadian Arctic. *Arctic med. Res.*, **49** (Suppl 2), 39–47
- MMWR (1994) *Reasons for Tobacco Use and Symptoms of Nicotine Withdrawal Among Adolescent and Young Adult Tobacco Users — United States, 1993*, Washington DC, US Department of Health and Human Services, Centers for Disease Control and Prevention, pp. 745–764
- Mody, J.K. & Ranadive, K.J. (1959) Biological study of tobacco in relation to oral cancer. *Indian J. med. Sci.*, **13**, 1023–1037
- Mookherjee, J.E. & Wilson, R.A. (1988) Tobacco constituents — Their importance in flavor and fragrance chemistry. *Recent Adv. Tob. Sci.*, **14**, 114–168
- Morse, R.M., Norvich, R.C. & Graf, J.A. (1977) Tobacco chewing: An unusual case of drug dependence. *Mayo Clin. Proc.*, **52**, 358–360
- Munirajan, A.K., Tutsumi-Ishii, Y., Mohanprasad, B.K.C., Hirano, Y., Munakata, N., Shanmugam, G. & Tsuchida, N. (1996) *p53* Gene mutations in oral carcinomas from India. *Int. J. Cancer*, **66**, 297–300
- Munirajan, A.K., Mohanprasad, B.K.C., Shanmugam, G. & Tsuchida, N. (1998) Detection of a rare point mutation at codon 59 and relatively high incidence of *H-ras* mutation in Indian oral cancer. *Int. J. Oncol.*, **13**, 971–974

- Murdia, U.S., Mehta, F.J. & Bhide, S.V. (1982) Nitrate reductase activity and nitrite levels in the saliva of habitual users of various tobacco products. *Food chem. Toxicol.*, **20**, 269–271
- Murphy, S.E., Carmella, S.G., Idris, A.M. & Hoffmann, D. (1994) Uptake and metabolism of carcinogenic levels of tobacco-specific nitrosamines by Sudanese snuff dippers. *Cancer Epidemiol. Biomarkers Prev.*, **3**, 423–428
- Murrah, V.A., Gilchrist, E.P. & Moyer, M.P. (1993) Morphologic and growth effects of tobacco associated chemical carcinogens and smokeless tobacco extracts on human oral epithelial cells in culture. *Oral Surg. oral Med. oral Pathol.*, **75**, 323–332
- Murray, D.M., Roche, L.M., Goldman, A.I. & Whitbeck, J. (1988) Smokeless tobacco use among ninth graders in a north-central metropolitan population: Cross-sectional and prospective associations with age, gender, race, family structure, and other drug use. *Prev. Med.*, **17**, 449–460
- Murray, R.P., Bailey, W.C., Daniels, K., Bjornson, W.M., Kurnow, K., Connett, J.E., Nides, M.A. & Kiley, J.P. (1996) Safety of nicotine polacrilex gum used by 3,094 participants in the Lung Health Study. *Chest*, **109**, 438–445
- Murti, P.R., Warnakulasuriya, K.A.A.S., Johnson, N.W., Bhonsle, R.B., Gupta, P.C., Daftary, D.K. & Mehta, F.S. (1998) p53 Expression in oral precancer as a marker for malignant potential. *J. oral Pathol. Med.*, **27**, 191–196
- Muscat, J.E. & Wynder, E.L. (1998) A case-control study of risk factors for major salivary gland cancer. *Otolaryngol. Head Neck Surg.*, **118**, 195–198
- Muscat, J.E., Hoffmann, D. & Wynder, E.L. (1995) The epidemiology of renal cell carcinoma. A second look. *Cancer*, **75**, 2552–2557
- Muscat, J.E., Richie, J.P., Jr, Thompson, S. & Wynder, E.L. (1996) Gender differences in smoking and risk for oral cancer. *Cancer Res.*, **56**, 5192–5197
- Muscat, J.E., Stellman, S.D., Hoffmann, D. & Wynder, E.L. (1997) Smoking and pancreatic cancer in men and women. *Cancer Epidemiol. Biomarkers Prev.*, **6**, 15–19
- Mustajbegovic, J., Zuskin, E., Schachter, E.N., Kern, J., Luburic-Milas, M. & Pucaric, J. (2003) Respiratory findings in tobacco workers. *Chest occup. environ. Lung Dis.*, **123**, 1740–1748
- Myers, M. (2003) Beware the rooster: Smokeless tobacco companies who claim they want to help. *Tob. Control*, **12**, 342
- Nagpal, J.K., Mishra, R. & Das, B.R. (2002a) Activation of *Stat-3* as one of the early events in tobacco chewing-mediated in oral carcinogenesis. *Cancer*, **94**, 2393–2400
- Nagpal, J.K., Patnaik, S. & Das, B.R. (2002b) Prevalence of high-risk human papilloma virus types and its association with *p53* codon 72 polymorphism in tobacco addicted oral squamous cell carcinoma (OSCC) patients of eastern India. *Int. J. Cancer*, **97**, 649–653
- Nair, U. & Bartsch, H. (2001) Metabolic polymorphisms as susceptibility markers for lung and oral cavity cancer. In: Miller, A.B., Bartsch, H., Boffetta, P., Dragsted, L. & Vainio, H., eds, *Biomarkers in Cancer Chemoprevention* (IARC Scientific Publications No. 154), Lyon, pp. 271–290
- Nair, J., Ohshima, H., Friesen, M., Croisy, A., Bhide, S.V. & Bartsch, H. (1985) Tobacco-specific and betel nut-specific N-nitroso compounds: Occurrence in saliva and urine of betel quid chewers and formation *in vitro* by nitrosation of betel quid. *Carcinogenesis*, **6**, 295–303
- Nair, J., Nair, U.J., Ohshima, H., Bhide, S.V. & Bartsch, H. (1987) Endogenous nitrosation in the oral cavity of chewers while chewing betel quid with or without tobacco. In: Bartsch, H., O'Neill, I.K. & Schulte-Herman, R., eds, *Relevance of N-Nitroso Compounds to Human Cancer: Exposures and Mechanisms* (IARC Scientific Publications No. 84), Lyon, pp. 465–469

- Nair, U.J., Pakhale, S.S., Spiegelhalter, B., Preussmann, R. & Bhide, S.V. (1987) Carcinogenic and cocarcinogenic constituents of Masherī, a pyrolysed tobacco product. *Indian J. Biochem. Biophys.*, **24**, 257–259
- Nair, U.J., Friesen, M., Richard, I., MacLennan, R., Thomas, S. & Bartsch, H. (1990) Effect of lime composition on the formation of reactive oxygen species from areca nut extract *in vitro*. *Carcinogenesis*, **11**, 2145–2148
- Nair, U.J., Ammigan, N., Nagabhushan, M., Amonkar, A.J. & Bhide, S.V. (1991a) Effect of vitamin A status of rats on metabolizing enzymes after exposure to tobacco extract or *N*'-nitrosornicotine. In: O'Neill, I.K., Chen, J. & Bartsch, H., eds, *Relevance to Human Cancer of N-Nitroso Compounds, Tobacco and Mycotoxins* (IARC Scientific Publications No. 105), Lyon, pp. 525–528
- Nair, U.J., Ammigan, N., Kayal, J.J. & Bhide, S.V. (1991b) Species differences in hepatic pulmonary and upper gastrointestinal tract biotransformation enzymes on long-term feeding of masherī — A pyrolyzed tobacco product. *Digest. Dis. Sci.*, **36**, 293–298
- Nair, U.J., Ammigan, N., Kulkarni, J.R. & Bhide, S.V. (1991c) Species differences in intestinal drug metabolising enzymes in mouse, rat and hamster and their inducibility by masherī, a pyrolysed tobacco product. *Indian J. exp. Biol.*, **29**, 256–258
- Nair, U., Obe, G., Nair, J., Maru, G.B., Bhide, S.V., Pieper, R. & Bartsch, H. (1991d) Evaluation of frequency of micronucleated oral mucosa cells as a marker for genotoxic damage in chewers of betel quid with or without tobacco. *Mutat. Res.*, **261**, 163–168
- Nair, U.J., Obe, G., Friesen, M., Goldberg, M.T. & Bartsch, H. (1992) Role of lime in the generation of reactive oxygen species from betel-quid ingredients. *Environ. Health Perspect.*, **98**, 203–205
- Nair, J., Ohshima, H., Nair, U.J. & Bartsch, H. (1996) Endogenous formation of nitrosamines and oxidative DNA-damaging agents in tobacco users. *Crit. Rev. Toxicol.*, **26**, 149–161
- Nair, U.J., Nair, J., Mathew, B. & Bartsch, H. (1999) Glutathione *S*-transferase M1 and T1 null genotypes as risk factors for oral leukoplakia in ethnic Indian betel quid/tobacco chewers. *Carcinogenesis*, **20**, 743–748
- Nair, U., Bartsch, H. & Nair, J. (2004) Alert for an epidemic of oral cancer due to use of the betel quid substitutes *gutkha* and *pan masala*: A review of agents and causative mechanisms. *Mutagenesis*, **19**, 251–262
- Nakajima, M., Kwon, J.T., Tanaka, N., Zenta, T., Yamamoto, Y., Yamamoto, H., Yamazaki, H., Yamamoto, T., Kuroiwa, Y. & Yokoi, T. (2001) Relationship between interindividual differences in nicotine metabolism and CYP2A6 genetic polymorphism in humans. *Clin. Pharmacol. Ther.*, **69**, 72–78
- Nanda, P.K. & Sharma, M.M. (1988) Immediate effect of tobacco chewing in the form of 'paan' on certain cardio-respiratory parameters. *Indian J. Physiol. Pharmacol.*, **32**, 105–113
- Nasr, M.M., Reepmeyer, J.C. & Tang, Y. (1998) In vitro study of nicotine release from smokeless tobacco. *J. Assoc. off. anal. Chem. int.*, **81**, 540–543
- Natanblut, S.L., Mital, M. & Zeller, M.R. (2001) The FDA's enforcement of age restrictions on the sale of cigarettes and smokeless tobacco. *J. public Health Manag. Pract.*, **7**, 1–10
- National Association of Attorneys General (1998) *Smokeless Tobacco Master Settlement Agreement* [available at: <http://www.naag.org/issues/tobacco/index.php?pubsec=0&sdpid=920>; accessed 24 August 2004]

- National Cancer Institute/Centers for Disease Control (2002) *Smokeless Tobacco Fact Sheets* [available at http://dcccps.nci.nih.gov/TCRB/stfact_sheet_combined10-23-02.pdf; accessed 22 June 2004]
- National Sample Survey Organisation (1998) *Consumption of Tobacco in India, 1993–94, Fifth Quinquennial Survey of Consumer Expenditure, NSS 50th Round July 1993–June 1994* (NSS Report No. 427), Department of Statistics, Government of India
- Nguyen, H., Lynch, B. & Dus, J. (2002) Review of Ariva — A compressed powdered tobacco product (Abstract). In: *Proceedings of the 3rd International Conference on Smokeless Tobacco, September 22–25, 2002*, Stockholm, Sweden
- Niphadkar, M.P., Contractor, Q.Q. & Bhisey, R.A. (1994) Mutagenic activity of gastric fluid from chewers of tobacco with lime. *Carcinogenesis*, **15**, 927–931
- Niphadkar, M.P., Bagwe, A.N. & Bhisey, R.A. (1996) Mutagenic potential of Indian tobacco products. *Mutagenesis*, **11**, 151–154
- Niraula, S.R. (2004) Tobacco use among women in Dharan, eastern Nepal. *J. Health Popul. Nutr.*, **22**, 68–74
- Nordgren, P. & Ramström, L. (1990) Moist snuff in Sweden — Tradition and evolution. *Br. J. Addict.*, **85**, 1107–1112
- Nugmanov, S.N. & Baimakanov, S.S. (1970) [The results of an epidemiological study of oropharyngeal tumours in Kazakhstan following the WHO project.] In: *Epidemiology of Malignant Tumours*, Alma-Ata, Nauka Publishing House, pp. 227–231 (in Russian)
- O'Connor, R.J., Flaherty, B.P., Quinio Edwards, B. & Kozlowski, L.T. (2003) Regular smokeless tobacco use is not a reliable predictor of smoking onset when psychosocial predictors are included in the model. *Nicotine Tob. Res.*, **5**, 535–543
- Offenbacher, S. & Weathers, D.R. (1985) Effects of smokeless tobacco on the periodontal, mucosal and caries status of adolescent males. *J. oral Pathol.*, **14**, 169–181
- Ohshima, H. & Bartsch, H. (1981) Quantitative estimation of endogenous nitrosation in humans by monitoring *N*-nitrosoproline excreted in the urine. *Cancer Res.*, **41**, 3658–3662
- Ohshima, H., Nair, J., Bourgade, M.-C., Friesen, M., Garren, L. & Bartsch, H. (1985) Identification and occurrence of two new *N*-nitrosamino acids in tobacco products: 3-(*N*-nitroso-*N*-methylamino) propionic acid and 4-(*N*-nitroso-*N*-methylamino) butyric acid. *Cancer Lett.*, **26**, 153–162
- Oldendorf, W.H. (1974) Lipid solubility and drug penetration in the blood–brain barrier. *Proc. Soc. exp. Biol. Med.*, **147**, 813–816
- Olds, R.S. (1988) Patterns and prevalence of smokeless tobacco use by high school seniors in New York. *J. School Health*, **58**, 374–378
- Österdahl, B.-G. & Slorach, S. (1988) Tobacco-specific *N*-nitrosamines in the saliva of habitual male snuff dippers. *Food Addit. Contam.*, **5**, 581–586
- Österdahl, B.-G., Jansson, C. & Paccou, A. (2004) Decreased levels of tobacco-specific *N*-nitrosamines in moist snuff on the Swedish market. *J. agric. Food Chem.*, **52**, 5085–5088
- Ozkul, Y., Donmez, H., Erenmemisoglu, A., Demirtas, H. & Imamoglu, N. (1997) Induction of micronuclei by smokeless tobacco on buccal mucosa cells of habitual users. *Mutagenesis*, **12**, 285–287
- Palladino, G., Adams, J.D., Brunnemann, K.D., Haley, N.J. & Hoffmann, D. (1986) Snuff-dipping in college students: A clinical profile. *Mil. Med.*, **151**, 342–346
- Pande, P., Mathur, M., Shukla, N.K. & Ralhan, R. (1998) pRb and p16 Protein alterations in human oral tumorigenesis. *Oral Oncol.*, **34**, 396–403

- Pande, P., Soni, S., Chakravarti, N., Mathur, M., Shukla, N.K. & Ralhan, R. (2001) Prognostic impact of Ets-1 overexpression in betel and tobacco related oral cancer. *Cancer Detect. Prev.*, **25**, 496–501
- Pandey, M.R. & Pathak, R.P. (2001) *Challenges of Tobacco Use Behavior in Central Development Region of Nepal: Global Youth Tobacco Survey, Nepal* [<http://www.cdc.gov/tobacco/global/gyts/reports/searo/2001/nepal01.html>]
- Park, N.-H., Sapp, J.P. & Herbosa, E.G. (1986) Oral cancer induced in hamsters with herpes simplex infection and simulated snuff dipping. *Oral Surg. oral Med. oral Pathol.*, **62**, 164–168
- Park, N.-H., Akoto-Amanfu, E. & Paik, D.I. (1988) Smokeless tobacco carcinogenesis: The role of viral and other factors. *Cancer J. Clin.*, **38**, 248–256
- Park, N.-H., Dokko, H., Li, S.-L. & Cherrick, H.M. (1991) Synergism of herpes simplex virus and tobacco-specific N'-nitrosamines in cell transformation. *J. oral maxillofac. Surg.*, **49**, 276–281
- Parsons, L.L., Smith, M.S., Hamilton, J.L. & MacKown, C.T. (1986) Nitrate reduction during curing and processing of burley tobacco. *Tob. Sci.*, **30**, 100–103
- Patja, K. & Vartianen, E. (2003) Tobacco epidemic in Finland. *Suomen Lääkärilehti (Finnish med. J.)*, **58**, 2959–2963
- Patnaik, S., Nayak, B.K. & Das, B.R. (1999) Genetic alteration of p53 in oral tumours. In: Varma, A., ed., *Oral Oncology, Vol. VI, Proceedings of the 6th International Congress on Cancer*, New Dehli, Macmillan, pp. 87–92
- Paudel, D. (2003) *Tobacco Use among Adolescent Students in Secondary Schools of Pokhara Sub Metropolitan City of Nepal*, MPH Thesis, Kathmandu, Tribhuvan University
- Paulson, R., Shanfeld, J., Sachs, L., Ismail, M. & Paulson, J. (1988) Effect of smokeless tobacco on the development of the CD-1 mouse fetus. *Teratog. Carcinog. Mutag.*, **8**, 81–93
- Paulson, R., Shanfeld, J., Sachs, L., Price, T. & Paulson, J. (1989) Effect of smokeless tobacco on the development of the CD-1 mouse fetus. *Teratology*, **40**, 483–494
- Paulson, R.B., Shanfeld, J., Prause, L., Iranpour, S. & Paulson, J.O. (1991) Pre- and post-conceptual tobacco effects on the CD-1 mouse fetus. *J. craniofac. genet. dev. Biol.*, **11**, 48–58
- Paulson, R.B., Shanfeld, J., Dean, J., Mullet, D., Fernandez, M. & Paulson, J.O. (1992) Alcohol and smokeless tobacco effects on the CD-1 mouse fetus. *J. craniofac. genet. dev. Biol.*, **12**, 107–117
- Paulson, R.B., Shanfeld, J., Vorhees, C.V., Sweazy, A., Gagni, S., Smith, A.R. & Paulson, J.O. (1993) Behavioral effects of prenatally administered smokeless tobacco on rat offspring. *Neurotoxicol. Teratol.*, **15**, 183–192
- Paulson, R.B., Shanfeld, J., Mullet, D., Cole, J. & Paulson, J.O. (1994a) Prenatal smokeless tobacco effects on the rat fetus. *J. craniofac. genet. dev. Biol.*, **14**, 16–25
- Paulson, R.B., Shanfeld, J., Vorhees, C.V., Cole, J., Sweazy, A. & Paulson, J.O. (1994b) Behavioral effects of smokeless tobacco on the neonate and young Sprague Dawley rat. *Teratology*, **49**, 293–305
- Payne, J.B., Johnson, G.K., Reinhardt, R.A. & Schmid, M. (1998) Histologic alterations following short-term smokeless tobacco exposure in humans. *J. periodont. Res.*, **33**, 274–279
- Peacock, E.E., Jr & Brawley, B.W. (1959) An evaluation of snuff and tobacco in the production of mouth cancer. *Plast. reconstr. Surg.*, **23**, 628–635
- Peacock, E.E., Jr, Greenberg, B.G. & Brawley, B.W. (1960) The effect of snuff and tobacco on the production of oral carcinoma: An experimental and epidemiological study. *Ann. Surg.*, **151**, 542–550

- Pednekar, M.S. & Gupta, P.C. (2004) Tobacco use among school students in Goa, India. *Indian J. public Health*, **48**, 147–152
- Peele, D.M., Danehower, D.A. & Goins, G.D. (1995) Chemical and biochemical changes during flue curing. *Recent Adv. Tob. Sci.*, **21**, 81–133
- Peele, D.M., Riddick, M.G. & Edwards, M.E. (2001) Formation of tobacco-specific nitrosamines in flue-cured tobacco. *Recent Adv. Tob. Sci.*, **27**, 3–12
- Peltzer, K. (1999) Smokeless tobacco use among urban white and black South Africans. *Psychol. Rep.*, **85**, 933–934
- Peltzer, K. (2003) Smokeless tobacco and cigarette use among black secondary school students in South Africa. *Subst. Use Misuse*, **38**, 1003–1016
- Persson, P.-G., Carlsson, S., Svanström, L., Östenson, C.-G., Efendic, S. & Grill, V. (2000) Cigarette smoking, oral moist snuff use and glucose intolerance. *J. intern. Med.*, **248**, 103–110
- Peterson, A.V., Marek, P.M. & Mann, S.L. (1989) Initiation and use of smokeless tobacco in relation to smoking. *Natl Cancer Inst. Monogr.*, **8**, 63–69
- Phukan, R.K., Ali, M.S., Chetia, C.K. & Mahanta, J. (2001) Betel nut and tobacco chewing; Potential risk factors of cancer of oesophagus in Assam, India. *Br. J. Cancer*, **85**, 661–667
- Pindborg, J.J. & Poulsen, H.E. (1962) Studies in oral leukoplakias. 1. The influence of snuff upon connective tissue of the oral mucosa. Preliminary report. *Acta pathol. microbiol. scand.*, **55**, 412–414
- Pindborg, J.J., Murti, P.R., Bhonsle, R.B. & Gupta, P.C. (1992) Global aspects of tobacco use and its implications for oral health. In: Gupta, P.C., Hamner, J.E., III. & Murti, P.R., eds, *Control of Tobacco-related Cancers and Other Diseases* (Proceedings of an International Symposium January 15–19, 1990, TIFR, Bombay), Oxford, Oxford University Press, pp. 13–23
- Pindborg, J.J., Reichart, P.A., Smith, C.J. & van der Waal, I. (1996) Histological typing of cancer and precancer of the oral mucosa, In: Sobin, L.H., ed., *International Histological Classification of Tumours*, 2nd Ed., Geneva, World Health Organization
- Poore, T.K., Johnson, G.K., Reinhardt, R.A. & Organ, C.C. (1995) The effects of smokeless tobacco on clinical parameters of inflammation and gingival crevicular fluid prostaglandin E₂, interleukin-1 α , and interleukin-1 β . *J. Periodontol.*, **66**, 177–183
- Prokopczyk, G., Adams, J.D., LaVoie, E.J. & Hoffmann, D. (1987) Effect of snuff and nicotine on DNA methylation by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*, **8**, 1395–1397
- Prokopczyk, B., Hoffmann, D., Cox, J.E., Djordjevic, M.V. & Brunnemann, K.D. (1992a) Supercritical fluid extraction in the determination of tobacco-specific *N*-nitrosamines in smokeless tobacco. *Chem. Res. Toxicol.*, **5**, 336–340
- Prokopczyk, B., Wu, M., Cox, J.E. & Hoffmann, D. (1992b) Bioavailability of tobacco-specific *N*-nitrosamines to the snuff dipper. *Carcinogenesis*, **13**, 863–866
- Prokopczyk, B., Wu, M., Cox, J.E., Amin, S., Desai, D., Idris, A.M. & Hoffmann, D. (1995) Improved methodology for the quantitative assessment of tobacco-specific *N*-nitrosamines in tobacco by supercritical fluid extraction. *J. agric. Food Chem.*, **43**, 916–922
- Putnam, S.D., Cerhan, J.R., Parker, A.S., Bianchi, G.D., Wallace, R.B., Cantor, K.P. & Lynch, C.F. (2000) Lifestyle and anthropometric risk factors for prostate cancer in a cohort of Iowa men. *Ann. Epidemiol.*, **10**, 361–369

- Rahman, M., Rahman, M., Flora, M.S., Rahman, M., Akter, S.F.U., Al Haque, N. & Kabir, I. (2001) *Community Based Primary Prevention of Non-communicable Diseases — An Intervention Study*, Mohakhali, Dhaka, National Institute of Preventive and Social Medicine
- Rahman, M., Rahman, M., Flora, M.S., Aktar, S.F.U., Hossain, S. & Mascie-Taylor, C.G.N. (2004) *Community Based Health Behaviour Surveillance in Urban and Rural Areas of Bangladesh — A Baseline Study*, Mohakhali, Dhaka, National Institute of Preventive and Social Medicine
- Ralhan, R., Nath, N., Agarwal, S., Mathur, M., Wasyluk, B. & Shukla, N.K. (1998) Circulating p53 antibodies as early markers of oral cancer: Correlation with p53 alterations. *Clin. Cancer Res.*, **4**, 2147–2152
- Ralhan, R., Agarwal, S., Nath, N., Mathur, M., Wasyluk, B. & Srivastava, A. (2001) Correlation between p53 gene mutations and circulating antibodies in betel- and tobacco-consuming North Indian population. *Oral Oncol.*, **37**, 243–250
- Ramaesh, T., Mendis, B.R.R.N., Ratnatunga, N. & Thattil, R.O. (1999) The effect of tobacco smoking and of betel chewing with tobacco on the buccal mucosa: A cytomorphometric analysis. *J. oral Pathol. Med.*, **28**, 385–388
- Ramström, L.M. (2000) Snuff — An alternative nicotine delivery system. In: Ferrence, R., Slade, J., Room, R. & Pope, M., eds, *Nicotine and Public Health*, Washington, DC, American Public Health Association, pp. 159–178
- Ranadive, K.J., Gothoskar, S.V. & Khanolkar, V.R. (1963) Experimental studies on the etiology of cancer types specific to India. a) Oral cancer; b) Kangri cancer. *Acta unio int. contr. cancer.*, **19**, 634–639
- Ranadive, K.J., Gothoskar, S.V., Rao, A.R., Tezabwalla, B.U. & Ambaye, R.Y. (1976) Experimental studies on betel nut and tobacco carcinogenicity. *Int. J. Cancer*, **17**, 469–476
- Randeria, J.D. (1972) Tobacco induced changes in the bladder, buccal and vaginal mucosae. *Indian J. med. Res.*, **60**, 694–698
- Rani, M., Bonu, S., Jha, P., Nguyen, S.N. & Jamjoum, L. (2003) Tobacco use in India: Prevalence and predictors of smoking and chewing in a national cross sectional household survey. *Tob. Control*, **12**, 4–11
- Rao, A.R. (1984) Modifying influences of betel quid ingredients on B(a)P-induced carcinogenesis in the buccal pouch of hamster. *Int. J. Cancer*, **33**, 581–586
- Ravi, D., Ramadas, K., Mathew, B.S., Nalinakumari, K.R., Nair, M.K. & Pillai, M.R. (1999) De novo programmed cell death in oral cancer. *Histopathology*, **34**, 241–249
- Reddy, K.S. & Gupta, P.C. (2004) *Report on Tobacco Control in India*, New Delhi, Ministry of Health and Family Welfare, Government of India
- Redmond, D.E. (1970) Tobacco and cancer: The first clinical report, 1761. *New Engl. J. Med.*, **282**, 18–23
- Richter, P. & Spierto, F.W. (2003) Surveillance of smokeless tobacco nicotine, pH, moisture, and unprotonated nicotine content. *Nicotine Tob. Res.*, **5**, 885–889
- Riley, W.T., Barenie, J.T. & Myers, D.R. (1989) Typology and correlates of smokeless tobacco use. *J. adolesc. Health Care*, **10**, 357–362
- Riley, W.T., Barenie, J.T., Woodard, C.E. & Mabe, P.A. (1996) Perceived smokeless tobacco addiction among adolescents. *Health Psychol.*, **15**, 289–292
- Ringel, J.S., Wasserman, J. & Pacula, R.L. (2000) *The Relationship Between Cigarette Use and Other Tobacco Products: Results from the 2000 National Youth Tobacco Survey*, Washington DC, American Legacy Foundation

- Robert, J.C. (1949) *The Story of Tobacco in America*, New York, Knopf, A.A.
- Roberts, D.L. (1988) Natural tobacco flavor. *Recent Adv. Tob. Sci.*, **14**, 49–81
- Roberts, D.M. (1997) Comparative cytology of the oral cavities of snuff users. *Acta cytol.*, **41**, 1008–1014
- Robertson, P.B., Walsh, M., Greene, J., Ernster, V., Grady, D. & Hauck, W. (1990) Periodontal effects associated with the use of smokeless tobacco. *J. Periodontol.*, **61**, 438–443
- Robinson, M., Houtsmuller, E., Moolchan, E.T. & Pickworth, W.B. (2000) Placebo cigarettes in smoking research. *Exp. clin. Psychopharmacol.*, **8**, 326–332
- Rodu, B., Stegmayr, B., Nasic, S. & Asplund, K. (2002) Impact of smokeless tobacco use on smoking in northern Sweden. *J. intern. Med.*, **252**, 398–404
- Rodu, B., Stegmayr, B., Nasic, S., Cole, P. & Asplund, K. (2003) Evolving patterns of tobacco use in northern Sweden. *J. intern. Med.*, **253**, 660–665
- Roed-Petersen, B. & Pindborg, J.J. (1973) A study of Danish snuff-induced oral leukoplakias. *J. oral Pathol.*, **2**, 301–313
- Rohani, M. & Agewall, S. (2004) Oral snuff impairs endothelial function in healthy snuff users. *J. intern. Med.*, **255**, 379–383
- Rothman, K.J. & Greenland, S. (1998) *Modern Epidemiology, 2nd Ed.*, Philadelphia, Lippincott, Williams & Wilkins
- Royal College of Physicians of London (2001) *Nicotine Addiction in Britain. A Report of the Tobacco Advisory Group of the Royal College of Physicians* [http://www.rcplondon.ac.uk/pubs/wp_nicotine_summary.htm]
- Rubinstein, I. (2000) Smokeless tobacco potentiates VIP-induced DNA synthesis and inactivates NEP 24.11 in oral keratinocytes. *Am. J. Physiol. Cell Physiol.*, **278**, C391–396
- Russell, M.A.H., Jarvis, M.J., Devitt, G. & Feyerabend, C. (1981) Nicotine intake by snuff users. *Br. med. J.*, **283**, 814–817
- Russell, M.A., Jarvis, M.J., West, R.J. & Feyerabend, C. (1985) Buccal absorption of nicotine from smokeless tobacco sachets. *Lancet*, **ii**, 1370
- Ryan, P. (1980) Snuff? Snus? At least, an alternative to smoke. *Smithsonian*, **10**, 176
- Salem, G., Juhl, R. & Schiødt, T. (1984) Oral malignant and premalignant changes in ‘shammah’-users from the Gizan region, Saudi Arabia. *Acta odontol. scand.*, **42**, 41–45
- Salonen, L., Axéll, T. & Helldén, L. (1990) Occurrence of oral mucosal lesions, the influence of tobacco habits and an estimate of treatment time in an adult Swedish population. *J. oral Pathol. Med.*, **19**, 170–176
- Sankaranarayanan, R., Duffy, S.W., Padmakumary, G., Day, N.E. & Padmanabhan, T.K. (1989a) Tobacco chewing, alcohol and nasal snuff in cancer of the gingiva in Kerala, India. *Br. J. Cancer*, **60**, 638–643
- Sankaranarayanan, R., Duffy, S.W., Day, N.E., Nair, M.K. & Padmakumary, G. (1989b) A case-control investigation of cancer of the oral tongue and the floor of the mouth in Southern India. *Int. J. Cancer*, **44**, 617–621
- Sankaranarayanan, R., Duffy, S.W., Padmakumary, G., Day, N.E. & Nair, M.K. (1990a) Risk factors for cancer of the buccal and labial mucosa in Kerala, southern India. *J. Epidemiol. Community Health*, **44**, 286–292
- Sankaranarayanan, R., Duffy, S.W., Nair, M.K., Padmakumary, G. & Day, N.E. (1990b) Tobacco and alcohol as risk factors in cancer of the larynx in Kerala, India. *Int. J. Cancer*, **45**, 879–882

- Sankaranarayanan, R., Duffy, S.W., Padmakumary, G., Muralidharan Nair, S., Day, N.E. & Padmanabhan, T.K. (1991) Risk factors for cancer of the oesophagus in Kerala, India. *Int. J. Cancer*, **49**, 485–489
- Sankaranarayanan, R., Mathew, B., Jacob, B.J., Thomas, G., Somanathan, T., Pisani, P., Pandey, M., Ramadas, K., Najeeb, K. & Abraham, E. for the Trivandrum Oral Cancer Screening Study Group (2000) Early findings from a community-based, cluster-randomized, controlled oral cancer screening trial in Kerala, India. *Cancer*, **88**, 664–673
- Saranath, D., Chang, S.E., Bhoite, L.T., Panchal., R.G., Kerr, I.B., Mehta, A.R., Johnson, N.W. & Deo, M.G. (1991) High frequency mutation in codons 12 and 61 of H-ras oncogene in chewing tobacco-related human oral carcinoma in India. *Br. J. Cancer*, **63**, 573–578
- Saranath, D., Panchal., R.G., Nair, R., Mehta, A.R., Sanghavi, V.D. & Deo, M.G. (1992) Amplification and overexpression of epidermal growth factor receptor gene in human oropharyngeal cancer. *Eur. J. Cancer*, **28B**, 139–143
- Saranath, D., Tandle, A.T., Teni, T.R., Dedhia, P.M., Borges, A.M., Parikh, D., Sanghavi, V. & Mehta, A.R. (1999) p53 Inactivation in chewing tobacco-induced oral cancers and leukoplakias from India. *Oral Oncol.*, **35**, 242–250
- Schäffler, G., Betz, C. & Richter, E. (1993) Mass spectrometric analysis of tobacco-specific hemoglobin adducts. *Environ. Health Perspect.*, **99**, 187–189
- Schei, E. (1992) Tobacco habits, lifestyle, and reactions to smoking restrictions among Norwegian military officers. *Mil. Med.*, **157**, 636–640
- Schei, E., Fønnebo, V. & Aarø, L.E. (1990) Use of smokeless tobacco among conscripts: A cross-sectional study of Norwegian army conscripts. *Prev. Med.*, **19**, 667–674
- Schildt, E.B., Eriksson, M., Hardell, L. & Magnuson, A. (1998) Oral snuff, smoking habits and alcohol consumption in relation to oral cancer in a Swedish case-control study. *Int. J. Cancer*, **77**, 341–346
- Schildt, E.B., Nylander, K., Eriksson, M., Hardell, L., Magnusson, A. & Roos, G. (2003) Expression of p53, PCNA, Ki-67 and bcl-2 in relation to risk factors in oral cancer — A molecular epidemiological study. *Int. J. Oncol.*, **22**, 861–868
- Schinke, S.P., Schilling, R.F., II, Gilchrist, L.D., Ashby, M.R. & Kitajima, E. (1987) Pacific north-west native American youth and smokeless tobacco use. *Int. J. Addict.*, **22**, 881–884
- Schmähl, D. (1965) [Investigation of chewing tobacco extracts for carcinogenic effects in rats.] *Arzneimittelforschung*, **15**, 704–705 (in German)
- Schmitterlöw, C.G., Hansson, E., Andersson, G., Applegren, L.-E. & Hoffmann, P.C. (1967) Distribution of nicotine in the central nervous system. *Ann. N.Y. Acad. Sci.*, **142**, 2–14
- Schroeder, K.L. & Chen, M.S. (1985) Smokeless tobacco and blood pressure. *New Engl. J. Med.*, **312**, 919
- Schroeder, J.C., Olshan, A.F., Baric, R., Dent, G.A., Weinberg, C.R., Yount, B., Cerhan, J.R., Lynch, C.F., Schuman, L.M., Tolbert, P.E., Rothman, N., Cantor, K.P. & Blair, A. (2002) A case-control study of tobacco use and other non-occupational risk factors for t(14;18) subtypes of non-Hodgkin's lymphoma (United States). *Cancer Causes Control*, **13**, 159–168
- Schwartz, S.M., Daling, J.R., Doody, D.R., Wipf, G.C., Carter, J.J., Madeleine, M.M., Mao, E.J., Fitzgibbons, E.D., Huang, S., Beckmann, A.M., McDougall, J.K. & Galloway, D.A. (1998) Oral cancer risk in relation to sexual history and evidence of human papillomavirus infection. *J. natl Cancer Inst.*, **90**, 1626–1636

- Sellers, E.M., Tyndale, R.F. & Fernandes, L.C. (2003) Decreasing smoking behaviour and risk through CYP2A6 inhibition. *Drug Discov. Today*, **8**, 487–493
- Severson, H.H. (1990) Psychosocial factors in the use of smokeless tobacco and their implications for P.L. 99-252. *J. public Health Dent.*, **50**, 90–97
- Severson, H.H. (2003) What have we learned from 20 years of research on smokeless tobacco cessation? *Am. J. med. Sci.*, **326**, 206–211
- Severson, H.H., Andrews, J.A., Lichtenstein, E., Gordon, J.S., Barckley, M.F. & Akers, L. (1998) Using the hygiene visit to deliver a tobacco cessation program: Results of a randomized clinical trial. *J. Am. dent. Assoc.*, **129**, 993–999
- Severson, H.H., Andrews, J.A., Lichtenstein, E., Gordon, J.S., Barckley, M. & Akers, L. (2000a) A self-help cessation program for smokeless tobacco users: Comparison of two interventions. *Nicotine Tob. Res.*, **2**, 363–370
- Severson, H.H., Akers, L., Andrews, J.A., Lichtenstein, E. & Jerome, A. (2000b) Evaluating two self-help interventions for smokeless tobacco cessation. *Addict. Behav.*, **25**, 465–470
- Shafey, O., Dolwick, S. & Guindon, G.E., eds (2003) *Tobacco Control Country Profile*, 2nd Ed., Atlanta, GA, American Cancer Society, WHO and International Union Against Cancer
- Shapiro, M.P., Keen, P., Cohen, L. & de Moor, N.G. (1955) Malignant disease in the Transvaal. III. Cancer of the respiratory tract. *S. Afr. med. J.*, **29**, 95–101
- Sharma, P.K., Lal, N. & Nagpaul, K.K. (1985) Study of trace amounts of U in snuff. *Health Phys.*, **48**, 811–812
- Sharma, A.K., Prokopczyk, B. & Hoffmann, D. (1991) Supercritical fluid extraction of moist snuff. *J. agric. Food Chem.*, **39**, 508–510
- Sheets, T.J. (1990) Pesticide residues on tobacco: Perception and realities. *Recent Adv. Tob. Sci.*, **17**, 33–69
- Shirnamé-Moré, L. (1991) Smokeless tobacco extracts mutate human cells. *Carcinogenesis*, **12**, 927–930
- Siegel, D., Benowitz, N., Ernster, V.L., Grady, D.G. & Hauck, W.W. (1992) Smokeless tobacco, cardiovascular risk factors, and nicotine and cotinine levels in professional baseball players. *Am. J. public Health*, **82**, 417–421
- Sikdar, N., Paul, R.R., Panda, C.K., Banerjee, S.K. & Roy, B. (2003) Loss of heterozygosity at *APC* and *MCC* genes of oral cancer and leukoplakia tissues from Indian tobacco chewers. *J. oral Pathol. Med.*, **32**, 450–454
- Sikdar, N., Paul, R.R. & Roy, B. (2004) Glutathione *S*-transferase *M3* (*A/A*) genotype as a risk factor for oral cancer and leukoplakia among Indian tobacco smokers. *Int. J. Cancer*, **109**, 95–101
- Singh, A. & Singh, S.P. (1998) Postnatal effect of smokeless tobacco on phytyc acid or the butylated hydroxyanisole-modulated hepatic detoxication system and antioxidant defense mechanism in suckling neonates and lactating mice. *Cancer Lett.*, **122**, 151–156
- Sinha, D.N. (2004) *Oral Tobacco Use and its Implications in Southeast Asia Region*, New Delhi, WHO South East Asia Regional Office
- Sinha, D.N. & Gupta, P.C. (2002a) *Report on Global Youth Tobacco Survey (GYTS) 2002, Uttar Pradesh, India* [http://www.cdc.gov/tobacco/global/GYTS/reports/india_UttarPradesh_2002; accessed 01 August 2004]

- Sinha, D.N. & Gupta, P.C. (2002b) *Report on Global Youth Tobacco Survey in Uttaranchal, India* [http://www.cdc.gov/tobacco/global/GYTS/reports/india_Uttaranchal_2002; accessed 01 August 2004]
- Sinha, D.N. & Gupta, P.C. (2004a) Tobacco use among school personnel in Orissa. *Indian J. public Health*, **48**, 123–127
- Sinha, D.N. & Gupta, P.C. (2004b) Tobacco use among students in Uttar Pradesh and Uttaranchal, India. *Indian J. public Health*, **48**, 132–137
- Sinha, D.N. & Roychowdhury, S. (2004) Tobacco control practices in 25 schools of West Bengal. *Indian J. public Health*, **48**, 128–131
- Sinha, D.N., Gupta, P.C., Pednekar, M.S., Jones, J.T. & Warren, C.W. (2002) Tobacco use among school personnel in Bihar, India. *Tob. Control*, **11**, 82–85
- Sinha, D.N., Gupta, P.C. & Pednekar, M.S. (2003a) Tobacco use in a rural area of Bihar, India. *Indian J. Community Med.*, **28**, 167–170
- Sinha, D.N., Gupta, P.C. & Pednekar, M.S. (2003b) Tobacco use among school personnel in eight north-eastern states of India. *Indian J. Cancer*, **40**, 3–14
- Sinha, D.N., Gupta, P.C. & Pednekar, M.S. (2003c) Tobacco use among students in the eight north-eastern states of India. *Indian J. Cancer*, **40**, 43–59
- Sinha, D.N., Gupta, P.C. & Pednekar, M. (2004a) Tobacco use among students in Bihar (India). *Indian J. public Health*, **48**, 111–117
- Sinha, D.N., Gupta, P.C. & Pednekar, M.S. (2004b) Use of tobacco products as dentifrice among adolescents in India: Questionnaire study. *Br. med. J.*, **328**, 323–324
- Sinusas, K., Coroso, J.G., Sopher, M.D. & Crabtree, B.F. (1992) Smokeless tobacco use and oral pathology in a professional baseball organization. *J. fam. Pract.*, **34**, 713–718
- Sisson, V.A. & Severson, R.F. (1990) Alkaloid composition of the *Nicotiana* species. *Beit. Tabakforsch.*, **14**, 327–340
- Slattery, M.L., Schumacher, M.C., West, D.W. & Robison, L.M. (1988) Smoking and bladder cancer. The modifying effect of cigarettes on other factors. *Cancer*, **61**, 402–408
- Smith, R.A., Lawson, T. & Johansson, S.L. (1997) Detection of DNA adducts by ³²P postlabeling following chronic exposure of rats to snuff. *Cancer Lett.*, **121**, 11–17
- Spangler, J.G. (2002) Corrections. *Ethn. Disease*, **12**, 158–159
- Spangler, J.G., Michielutte, R., Bell, R.A., Knick, S., Dignan, M.B. & Summerson, J.H. (2001a) Dual tobacco use among Native American adults in southeastern North Carolina. *Prev. Med.*, **32**, 521–528
- Spangler, J.G., Michielutte, R., Bell, R.A. & Dignan, M.B. (2001b) Association between smokeless tobacco use and breast cancer among Native-American women in North Carolina. *Ethn. Dis.*, **11**, 36–43
- Spence, J.C. & Gauvin, L. (1996) Drug and alcohol use by Canadian university athletes: A national survey. *J. Drug Educ.*, **26**, 275–287
- Spiegelhalter, B. & Fischer, S. (1991) Formation of tobacco-specific nitrosamines. *Crit. Rev. Toxicol.*, **21**, 241
- Spillane, J. & McAllister, W.B. (2003) Keeping the lid on: A century of drug regulation and control. *Drug Alcohol Depend.*, **70**, S5–S12
- Spitz, M.R., Fueger, J.J., Goepfert, H., Hong, W.K. & Newell, G.R. (1988) Squamous cell carcinoma of the upper aerodigestive tract. A case comparison analysis. *Cancer*, **61**, 203–208

- Spitz, M.R., Fueger, J.J., Halabi, S., Schantz, S.P., Sample, D. & Hsu, T.C. (1993) Mutagen sensitivity in upper aerodigestive tract cancer: A case-control analysis. *Cancer Epidemiol. Biomarkers Prev.*, **2**, 329–333
- Squier, C.A. (1986) Penetration of nicotine and nitrosornicotine across porcine oral mucosa. *J. appl. Toxicol.*, **6**, 123–128
- Squires, W.G., Brandon, T.A., Zinkgraf, S., Bonds, D., Hartung, G.H., Murray, T., Jackson, A.S. & Miller, R.R. (1984) Hemodynamic effects of oral smokeless tobacco in dogs and young adults. *Prev. Med.*, **13**, 195–206
- Stanfill, S.B., Brown, C.R., Yan, X.J., Watson, C.H. & Ashley, D.L. (2006) Quantification of flavor-related compounds in the unburned contents of bidi and clove cigarettes. *J. agric. Food Chem.*, **54**, 8580–8588
- Stegmayr, B., Johansson, I., Huhtasaari, F., Moser, U. & Asplund, K. (1993) Use of smokeless tobacco and cigarettes — Effects on plasma levels of antioxidant vitamins. *Int. J. Vit. nutr. Res.*, **63**, 195–200
- Stepanov, I. & Hecht, S.S. (2005) Tobacco-specific nitrosamines and their pyridine-*N*-glucuronides in the urine of smokers and smokeless tobacco users. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 885–891
- Stepanov, I., Hecht, S.S., Ramakrishnan, S. & Gupta, P.C. (2005) Tobacco-specific nitrosamines in smokeless tobacco products marketed in India. *Int. J. Cancer*, **116**, 16–19
- Stepanov, I., Jensen, J., Hatsukami, D. & Hecht, S.S. (2006) Tobacco-specific nitrosamines in new tobacco products. *Nicotine Tob. Res.*, **8**, 309–313
- Sterling, T.D., Rosenbaum, W.L. & Weinkam, J.J. (1992) Analysis of the relationship between smokeless tobacco and cancer based on data from the National Mortality Followback Survey. *J. clin. Epidemiol.*, **45**, 223–231
- Stevens, V.J., Severson, H., Lichtenstein, E., Little, S.J. & Leben, J. (1995) Making the most of a teachable moment: A smokeless-tobacco cessation intervention in the dental office. *Am. J. public Health*, **85**, 231–235
- Stewart, G.G. (1967) A history of medicinal use of tobacco 1492–1860. *Med. Hist.*, **11**, 228–268
- Steyn, K., Bradshaw, D., Norman, R., Laubscher, R. & Saloojee, Y. (2002) Tobacco use in South Africans during 1998. The first demographic and health survey. *J. cardiovasc. Risk*, **9**, 161–170
- Stich, H.F. & Anders, F. (1989) The involvement of reactive oxygen species in oral cancers of betel quid/tobacco chewers. *Mutat. Res.*, **214**, 47–61
- Stich, H.F., Curtis, J.R. & Parida, B.B. (1982a) Application of the micronucleus test to exfoliated cells of high cancer risk groups: Tobacco chewers. *Int. J. Cancer*, **30**, 553–559
- Stich, H.F., Stich, W. & Parida, B.B. (1982b) Elevated frequency of micronucleated cells in the buccal mucosa of individuals at high risk of oral cancer: Betel quid chewers. *Cancer Lett.*, **17**, 125–134
- Stich, H.F., Parida, B.B. & Brunnemann, K.D. (1992) Localized formation of micronuclei in the oral mucosa and tobacco-specific nitrosamines in the saliva of ‘reverse’ smokers, khaini-tobacco chewers and gudakhu users. *Int. J. Cancer*, **50**, 172–176
- Stockwell, H.G. & Lyman, G.H. (1986) Impact of smoking and smokeless tobacco on the risk of cancer of the head and neck. *Head Neck Surg.*, **9**, 104–110
- Stratton, K.S., Shetty, P., Wallace, R. & Bondurant, S., eds (2001) *Clearing the Smoke: Assessing the Science Base for Tobacco Harm Reduction*, Washington DC, National Academy Press

- Suadicani, P., Hein, H.O. & Gyntelberg, F. (1997) Mortality and morbidity of potentially misclassified smokers. *Int. J. Epidemiol.*, **26**, 321–327
- Sudbo, J., Bryne, M., Johannessen, A.C., Kildal, W., Danielsen, H.E. & Reith, A. (2001) Comparison of histological grading and large-scale genomic status (DNA ploidy) as prognostic tools in oral dysplasia. *J. Pathol.*, **194**, 303–310
- Summerlin, D.-J., Dunipace, A. & Potter, R. (1992) Histologic effects of smokeless tobacco and alcohol on the pouch mucosa and organs of the Syrian hamster. *J. oral Pathol. Med.*, **21**, 105–108
- Suri, K., Goldman, H.M. & Wells, H. (1971) Carcinogenic effect of a dimethyl sulphoxide extract of betel nut on the mucosa of the hamster buccal pouch. *Nature*, **230**, 383–384
- Sussman, S., Holt, L., Dent, C.W., Flay, B.R., Graham, J.W., Hansen, W.B. & Johnson, C.A. (1989) Activity involvement, risk-taking, demographic variables, and other drug use: Prediction of trying smokeless tobacco. *Natl Cancer Inst. Monogr.*, **8**, 57–62
- Suzuki, H., Gao, X.-P., Olopade, C.O., Jaffe, H.A., Pakhlevanians, S. & Rubinstein, I. (1996) Aqueous smokeless tobacco extracts impairs endothelium-dependent vasodilation in the oral mucosa. *J. appl. Physiol.*, **81**, 225–231
- Tandle, A.T., Sanghvi, V. & Saranath, D. (2001) Determination of p53 genotypes in oral cancer patients from India. *Br. J. Cancer*, **84**, 739–742
- Targher, G., Alberiche, M., Zenere, M.B., Bonadonna, R.C., Muggeo, M. & Bonora, E. (1997) Cigarette smoking and insulin resistance in patients with noninsulin-dependent diabetes mellitus. *J. clin. Endocrinol. Metab.*, **82**, 3619–3624
- Teni, T., Pawar, S., Sanghvi, V. & Saranath, D. (2002) Expression of bcl-2 and bax in chewing tobacco-induced oral cancers and oral lesions from India. *Pathol. oncol. Res.*, **8**, 109–114
- Thomas, G., Hashibe, M., Jacob, B.J., Ramadas, K., Mathew, B., Sankaranarayanan, R. & Zhang, Z.-F. (2003) Risk factors for multiple oral premalignant lesions. *Int. J. Cancer*, **107**, 185–291
- Tilashalski, K., Rodu, B. & Mayfield, C. (1994) Assessing the nicotine content of smokeless tobacco products. *J. Am. dent. Assoc.*, **125**, 590–592
- Tilashalski, K., Rodu, B. & Cole, P. (1998) A pilot study of smokeless tobacco in smoking cessation. *Am. J. Med.*, **104**, 456–458
- Tilashalski, K., Rodu, B. & Cole, P. (2003) Seven-year followup of smoking cessation with smokeless tobacco. In: *Summary Report, 3rd International Conference on Smokeless Tobacco: Advancing Science and Protecting Public Health. Stockholm, Sweden, 22–25 September 2002*, Rockville, MD, National Cancer Institute, Division of Cancer Control and Population Sciences, Behavioral Research Program, Tobacco Control Branch
- Tillgren, P., Haglund, B.J.A., Lundberg, M. & Romelsjö, A. (1996) The sociodemographic pattern of tobacco cessation in the 1980s: Results from a panel study of living condition surveys in Sweden. *J. Epidemiol. Community Health*, **50**, 625–630
- Tobacco and Health Research Institute (1987) *Smokeless Tobacco Research Products*, Lexington, KY, University of Kentucky
- Tobacco Board (1992) Annual Report 1992, Pretoria
- Tobacco Board (1994/95) Annual Report 1994/95, Pretoria
- Tobacco Board (2006) *Export of Tobacco Products from 1998–99*. Ministry of Commerce, Government of India [available at <http://indiantobacco.com/expertsoftobpro9697.htm>; accessed 01 August 2006]

- Tobacco Control Programme (2004) *Smokeless Tobacco Sales in Canada, 1989–2003* (unpublished data received from Tobacco Control Programme, Health Canada on 12 August 2004)
- Tobey, N.A., Schreiner, V.J., Readling, R.D. & Orlando, R.C. (1988) The acute effects of smokeless tobacco on transport and barrier function of buccal mucosa. *J. dent. Res.*, **67**, 1414–1421
- Tolbert, P.E., Shy, C.M. & Allen, J.W. (1991) Micronuclei and other nuclear anomalies in buccal smears: Field tests in snuff users. *Am. J. Epidemiol.*, **134**, 840–850
- Tolbert, P.E., Shy, C.M. & Allen, J.W. (1992) Micronuclei and other nuclear anomalies in buccal smears: Methods development. *Mutat. Res.*, **271**, 69–77
- Tomar, S.L. (2002) Snuff use and smoking in U.S. men. Implication for harm reduction. *Am. J. prev. Med.*, **23**, 143–149
- Tomar, S.L. (2003a) Trends and patterns of tobacco use in the United States. *Am. J. med. Sci.*, **326**, 248–254
- Tomar, S.L. (2003b) Is use of smokeless tobacco a risk factor for cigarette smoking? The U.S. experience. *Nicotine Tob. Res.*, **5**, 545–553
- Tomar, S.L. (2003c) Smokeless tobacco use is a significant predictor of smoking when appropriately modeled. *Nicotine Tob. Res.*, **5**, 571–573
- Tomar, S.L. (2007) Smokeless tobacco products in the United States and Sweden. In: *National Cancer Institute, Background Papers for the Scientific Advisory Committee on Tobacco Product Regulation* (Smoking and Tobacco Control Monograph 19), Geneva, World Health Organization
- Tomar, S.L. & Giovino, G.A. (1998) Incidence and predictors of smokeless tobacco use among US youth. *Am. J. public Health*, **88**, 20–26
- Tomar, S.L. & Henningfield, J.E. (1997) Review of the evidence that pH is a determinant of nicotine dosage from oral use of smokeless tobacco. *Tob. Control*, **6**, 219–225
- Tomar, S.L. & Loree, M. (2004) Errors in analyzing associations between use of smokeless tobacco and cigarettes. *Addiction*, **99**, 260–262
- Tomar, S.L. & Winn, D.M. (1999) Chewing tobacco use and dental caries among U.S. men. *J. Am. dent. Assoc.*, **130**, 1601–1610
- Tomar, S.L., Giovino, G.A. & Eriksen, M.P. (1995) Smokeless tobacco brand preference and brand switching among US adolescents and young adults. *Tob. Control*, **4**, 67–72
- Tomar, S.L., Winn, D.M., Swango, P.A., Giovino, G.A. & Kleinman, D.V. (1997) Oral mucosal smokeless tobacco lesions among adolescents in the United States. *J. dent. Res.*, **76**, 1277–1286
- Tomar, S.L., Connolly, G.N., Wilkenfeld, J. & Henningfield, J.E. (2003) Declining smoking in Sweden: Is Swedish Match getting the credit for Swedish tobacco control's efforts? *Tob. Control*, **12**, 368–371
- Tricker, A.R. (2003) Nicotine metabolism, human drug metabolism polymorphisms, and smoking behaviour. *Toxicology*, **183**, 151–173
- Tricker, A.R. & Preussmann, R. (1988) The occurrence of *N*-nitroso compounds in zarda tobacco. *Cancer Lett.*, **42**, 113–118
- Tricker, A.R. & Preussmann, R. (1989) The occurrence of *N*-nitroso compounds in kiwam tobacco. *Cancer Lett.*, **49**, 221–224
- Tricker, A.R. & Preussmann, R. (1991) Occurrence of and exposure to *N*-nitroso compounds in tobacco. In: O'Neill, I.K., Chen, J. & Bartsch, H., eds, *Relevance to Human Cancer of N-*

- Nitroso Compounds, Tobacco Smoke and Mycotoxins* (IARC Scientific Publications No. 105), Lyon, pp. 493–495
- Tricker, A.R., Haubner, R., Spiegelhalder, B. & Preussmann, R. (1988) The occurrence of tobacco-specific nitrosamines in oral tobacco products and their potential formation under simulated gastric conditions. *Food chem. Toxicol.*, **26**, 861–865
- Trivedi, A.H., Bakshi, S.R., Dave, B.J., Adhvaryu, S.G., Patel, D.D. & Balar, D.B. (1995) Elevated mutagen susceptibility in cultured lymphocytes of oral cancer patients. *Anticancer Res.*, **15**, 2589–2592
- Tso, T.C. (1990) *Production, Physiology, and Biochemistry of Tobacco Plant*, Beltsville, MD, Ideals, pp. 55–64, 431, 595–614
- Tso, T.C., Harley, N. & Alexander, L.T. (1966) Source of lead-210 and polonium-210 in tobacco. *Science*, **153**, 880–882
- Tucker, L.A. (1989) Use of smokeless tobacco, cigarette smoking, and hypercholesterolemia. *Am. J. public Health*, **79**, 1048–1050
- Turner, D.M., Armitage, A.K., Briant, R.H. & Dollery, C.T. (1975) Metabolism of nicotine by the isolated perfused dog lung. *Xenobiotica*, **5**, 539–551
- Tyldesley, W.R. (1971) Tobacco chewing in English coal miners. A preliminary report. *Br. J. oral Surg.*, **9**, 21–28
- Ugen, S. (2003) Bhutan: The world's most advanced tobacco control nation? *Tob. Control*, **12**, 431–433
- Uitti, J., Nordman, H., Huuskonen, M.S., Roto, P., Husman, K. & Reiman, M. (1998) Respiratory health of cigar factory workers. *Occup. environ. Med.*, **55**, 834–839
- United Nations Convention on Psychotropic Substances (1971) *1971 United Nations Conference for the Adoption of a Protocol on Psychotropic Substances, and the Schedules Annexed to the Convention*, New York
- United Nations Single Convention on Narcotic Drugs (1961) *International Narcotics Control Board, 1995–2005*, New York
- Vaidya, S.G., Vaidya, N.S. & Naik, U.D. (1992) Epidemiology of tobacco habits in Goa, India. In: Gupta, P.C., Hammer, J.E., III & Murti, P.R., eds, *Control of Tobacco-related Cancers and Other Diseases* (Proceedings of an International Symposium January 15–19, 1990, TIFR, Bombay), Oxford, Oxford University Press, pp. 25–46
- Van Duuren, B.L., Sivak, A., Segal, A., Orris, L. & Langseth, L. (1966) The tumor-promoting agents of tobacco leaf and tobacco smoke condensate. *J. natl. Cancer Inst.*, **37**, 519–526
- Verma, R.C., Chansoriya, M. & Kaul, K.K. (1983) Effect of tobacco chewing by mothers on fetal outcome. *Indian Pediatr.*, **20**, 105–111
- Vineis, P., Malats, N., Lang, M., d'Errico, A., Caporaso, N., Cuzick, J. & Boffetta, P., eds (1999) *Metabolic Polymorphisms and Susceptibility to Cancer* (IARC Scientific Publications No. 148), Lyon
- Vishwanatha, J.K., Swinney, R. & Banerjee, A.G. (2003) Modulation of annexin I and cyclooxygenase-2 in smokeless tobacco-induced inflammation and oral cancer. *Mol. cell. Biochem.*, **248**, 67–75
- Voges, E. (1984) The pleasures of tobacco — How it all began and the whole story. *Tob. J. int.*, **1**, 80–82
- Vogler, W.R., Lloyd, J.W. & Milmore, B.K. (1962) A retrospective study of etiological factors in cancer of the mouth, pharynx, and larynx. *Cancer*, **15**, 246–258

- Wahi, P.N. (1968) The epidemiology of oral and oropharyngeal cancer. A report of the study in Mainpuri district, Uttar Pradesh, India. *Bull. World Health Org.*, **38**, 495–521
- Wallenfeldt, K., Hulthe, J., Bokemark, L., Wikstrand, J. & Fagerberg, B. (2001) Carotid and femoral atherosclerosis, cardiovascular risk factors and C-reactive protein in relation to smokeless tobacco use or smoking in 58-year-old men. *J. intern. Med.*, **250**, 492–501
- Walsh, M.M., Hilton, J.F., Ernster, V.L., Masouredis, C.M. & Grady, D.G. (1994) Prevalence, patterns, and correlates of spit tobacco use in a college athlete population. *Addict. Behav.*, **19**, 411–427
- Walsh, M.M., Hilton, J.F., Masouredis, C.M., Gee, L., Chesney, M.A. & Ernster, V.L. (1999) Smokeless tobacco cessation intervention for college athletes: Results after 1 year. *Am. J. public Health*, **89**, 228–234
- Walsh, M.M., Ellison, J., Hilton, J.F., Chesney, M. & Ernster, V.L. (2000) Spit (smokeless) tobacco use by high school baseball athletes in California. *Tob. Control*, **9** (Suppl. 2), ii32–ii39
- Walton, L.R., Burton, H.R. & Swetnam, L.D. (1995) Effect of mechanization on the physical appearance and chemical composition of burley tobacco. *Recent Adv. Tob. Sci.*, **21**, 3–38
- Wang, H., Yang, B., Zhang, L., Xu, D. & Wang, Z. (2000) Direct block of inward rectifier potassium channels by nicotine. *Toxicol. appl. Pharmacol.*, **164**, 97–101
- Wang, Y., Rotem, E., Andriani, F. & Garlick, J.A. (2001) Smokeless tobacco extracts modulate keratinocyte and fibroblast growth in organotypic culture. *J. dent. Res.*, **80**, 1862–1866
- Warnakulasuriya, K.A.A.S. & McDonald, D.G. (1995) Epithelial cell kinetics in oral leukoplakia. *J. oral Pathol. Med.*, **24**, 165–169
- Wasnik, K.S., Ughade, S.N., Zodpey, S.P. & Ingole, D.L. (1998) Tobacco consumption practices and risk of oropharyngeal cancer: A case-control study in central India. *S.E. Asian J. trop. Med. public Health*, **29**, 827–834
- Wayne, G.F. & Connolly, G.N. (2004) Application, function, and effects of menthol in cigarettes: A survey of tobacco industry documents. *Nicotine Tob. Res.*, **6**, 43–54
- Wedenberg, C., Jönsson, A. & Hirsch, J.M. (1996) Assessment of p53 and Ki-67 expression in snuff-induced lesions. *Br. J. oral maxillofac. Surg.*, **34**, 409–413
- Weeks, W.W., Chaplin, J.F. & Campbell, C.R. (1989) Capillary chromatography: Evaluation of volatiles from flue-cured tobacco varieties. *J. agric Food Chem.*, **37**, 1038–1045
- Wenke, G., Brunnemann, K.D., Hoffmann, D. & Bhide, S.V. (1984) A study of betel quid carcinogenesis. IV. Analysis of the saliva of betel chewers: A preliminary report. *J. Cancer Res. Clin. Oncol.*, **108**, 110–113
- Wennmalm, A., Benthin, G., Granström, E.F., Persson, L., Petersson, A.-S. & Winell, S. (1991) Relation between tobacco use and urinary excretion of thromboxane A₂ and prostacyclin metabolites in young men. *Circulation*, **83**, 1698–1704
- Westman, E.C. & Guthrie, G.P. (1990) Licorice, tobacco chewing, and hypertension. *New Engl. J. Med.*, **322**, 849–850
- Wetter, D.W., McClure, J.B., de Moor, C., Cofta-Gunn, L., Cummings, S., Cinciripini, P.M. & Gritz, E.R. (2002) Concomitant use of cigarettes and smokeless tobacco: Prevalence, correlates, and predictors of tobacco cessation. *Prev. Med.*, **34**, 638–648
- WHO (1988) *Smokeless Tobacco Control* (Technical Report Series 773), Geneva
- WHO (1997) *Tobacco or Health: A Global Status Report*, Geneva

- WHO (2003a) *WHO Framework Convention on Tobacco Control* [available at http://www.who.int/tobacco/framework/WHO_FCTC_english.pdf and http://www.who.int/tobacco/areas/framework/signing_ceremony/countrylist/en; accessed 24 August 2004]
- WHO (2003b) *Recommendation on Smokeless Tobacco Products. WHO Scientific Advisory Committee on Tobacco Product Regulation (SACTob)* (LC/NLM Classification: HD 9130.6), Geneva
- WHO (2006) *WHO ICD 10 International Statistical Classification of Diseases and Related Health Problems, 10th Revision Version for 2006: F17: Mental and Behavioural Disorders due to Use of Tobacco* [<http://www3.who.int/icd/currentversion/fr-icd.htm>]
- WHO EURO (2004) *Tobacco Control Database*, Copenhagen, WHO Regional Office for Europe [available at <http://data.euro.who.int/tobacco/>; accessed 24 August 2004]
- WHO SEARO (2001) *Sentinel Prevalence Study of Tobacco Use In Myanmar, 2001*
- WHO SEARO (2004) *Multisectoral Mechanism for Comprehensive Tobacco Control. Report of a regional consultation, Bangkok, 9–10 July 2003; WHO Project:ICP TOB 001*, New Delhi, WHO, Regional Office for South-East Asia
- Whong, W.-Z., Ames, R.G. & Ong, T.-M. (1984) Mutagenicity of tobacco snuff: Possible health implications for coal miners. *J. Toxicol. environ. Health*, **14**, 491–496
- Whong, W.-Z., Stewart, J.D. & Ong, T.-M. (1985) Formation of bacterial mutagens from the reaction of chewing tobacco with nitrite. *Mutat. Res.*, **158**, 105–110
- Wicklin, B. (2005) *Nordic Tobacco Statistics 1970–2004, Smoking and ‘Snus’ Prevalence, Consumption of Smoking Tobacco and ‘Snus’, Price Movements and Consumer Spending in Sweden, Norway, Denmark, Finland and Iceland, Statistical Report Commissioned by the European Smokeless Tobacco Council*, Stockholm, Statistical Bureau VECA^{HB}
- Wicklin, B. (2006) *Tobacco Statistics: Sales, Consumption and Lung Cancer*, Hässelby, Statistical Bureau VECA [available at <http://www.statveca.com/english/index.html>; accessed 1 June 2006]
- Wiernik, A., Christakopoulos, A., Johansson, L. & Wahlberg, I. (1995) Effect of air-curing on the chemical composition of tobacco. *Recent Adv. Tob. Sci.*, **21**, 39–80
- Will, J.C., Galuska, D.A., Ford, E.S., Mokdad, A. & Calle, E.E. (2001) Cigarette smoking and diabetes mellitus: Evidence of a positive association from a large prospective cohort study. *Int. J. Epidemiol.*, **30**, 540–546
- Williams, R.R. & Horm, J.W. (1977) Association of cancer sites with tobacco and alcohol consumption and socioeconomic status of patients: Interview study from the Third National Cancer Survey. *J. natl Cancer Inst.*, **58**, 525–547
- Williams, N.J., Arheart, K.L. & Klesges, R. (1995) A smokeless tobacco cessation program for postsecondary students. *Health Values*, **19**, 33–42
- Winn, D.M. (1986) Smokeless tobacco and oral/pharynx cancer: The role of cofactors. In: Hoffmann, D. & Harris, C.C., eds, *Mechanisms in Tobacco Carcinogenesis* (Banbury Report No. 23), Cold Spring Harbor, NY, Cold Spring Harbor Laboratory, pp. 361–375
- Winn, D.M., Blot, W.J., Shy, C.M., Pickle, L.W., Toledo, A. & Fraumeni, J.F., Jr (1981a) Snuff dipping and oral cancer among women in the southern United States. *New Engl. J. Med.*, **304**, 745–749
- Winn, D.M., Blot, W.J. & Fraumeni, J.F., Jr (1981b) Snuff dipping and oral cancer. *New Engl. J. Med.*, **305**, 230–231
- Winn, D.M., Ziegler, R.G., Pickle, L.W., Gridley, G., Blot, W.J. & Hoover, R.N. (1984) Diet in the etiology of oral and pharyngeal cancer among women from the southern United States. *Cancer Res.*, **44**, 1216–1222

- Wolfe, M.D. & Carlos, J.P. (1987) Oral health effects of smokeless tobacco use in Navajo Indian adolescents. *Community dent. Oral Epidemiol.*, **15**, 230–235
- Wood, M.W., Medina, J.E., Thompson, G.C., Houck, J.R. & Min, K.-W. (1994) Accumulation of the p53 tumour-suppressor gene product in oral leukoplakia. *Otolaryngol. Head Neck Surg.*, **111**, 758–763
- World Bank (2000) *Economics of Tobacco in Turkey, December 2000* [<http://www1.worldbank.org/tobacco/pdf/country%20briefs/Turkey.pdf>; accessed 24 August 2004]
- Wray, A. & McGuirt, W.F. (1993) Smokeless tobacco usage associated with oral carcinoma. *Arch. otolaryngol. Head Neck Surg.*, **119**, 929–933
- Wu, X., Zhao, H., Suk, R. & Christiani, D.C. (2004) Genetic susceptibility to tobacco-related cancer. *Oncogene*, **23**, 6500–6523
- Wyckham, R.G. (1999) Smokeless tobacco in Canada: Deterring market development. *Tob. Control*, **8**, 411–420
- Wynder, E.L. & Wright, G. (1957) A study of tobacco carcinogenesis. I. The primary fractions. *Cancer*, **10**, 255–271
- Xu, J., Gimenez-Conti, I.B., Cunningham, J.E., Collet, A.M., Luna, M.A., Lanfranchi, H.E., Spitz, M.R. & Conti, C.J. (1998) Alterations of p53, cyclin D1, Rb and *H-ras* in human oral carcinomas related to tobacco use. *Cancer*, **83**, 204–212
- Yanev, I. & Kostianev, S. (2004) Respiratory findings in tobacco industry workers (Letter to the Editor). *Chest*, **125**, 802
- Ye, W., Ekström, A.M., Hansson, L.-E., Bergström, R. & Nyrén, O. (1999) Tobacco, alcohol and the risk of gastric cancer by sub-site and histologic type. *Int. J. Cancer*, **83**, 223–229
- Yildiz, D. (2004) Nicotine, its metabolism and an overview of its biological effects. *Toxicol.*, **43**, 619–632
- Yildiz, D., Liu, Y.-S., Ercal, N. & Armstrong, D.W. (1999) Comparison of pure nicotine- and smokeless tobacco extract-induced toxicities and oxidative stress. *Arch. environ. Contam. Toxicol.*, **37**, 434–439
- Young, T.B., Ford, C.N. & Brandenburg, J.H. (1986) An epidemiologic study of oral cancer in a statewide network. *Am. J. Otolaryngol.*, **7**, 200–208
- Zahm, S.H., Blair, A., Holmes, F.F., Boysen, C.D., Robel, R.J. & Fraumeni, J.F., Jr (1989) A case-control study of soft-tissue sarcoma. *Am. J. Epidemiol.*, **130**, 665–674
- Zahm, S.H., Heineman, E.F. & Vaught, J.B. (1992) Soft tissue sarcoma and tobacco use: Data from a prospective cohort study of United States veterans. *Cancer Causes Control*, **3**, 371–376
- Zaridze, D.G., Kuvshinov, J.P., Matiakin, E., Polakov, B.I., Boyle, P. & Blettner, M. (1985) Chemoprevention of oral and esophageal cancer in Uzbekistan, Union of Soviet Socialist Republics. *Natl Cancer Inst. Monogr.*, **69**, 259–262
- Zaridze, D.G., Matiakin, E.G., Stich, H.F., Hoffmann, D., Blettner, M., Poljakov, B.P., Rosin, M.P. & Brunnemann, K.D. (1986) The effect of nass use and smoking on the risk of oral leukoplakia. *Cancer Detect. Prev.*, **9**, 435–440
- Zheng, W., McLaughlin, J.K., Gridley, G., Bjelke, E., Schuman, L.M., Silverman, D.T., Wacholder, S., Co-Chien, H.T., Blot, W.J. & Fraumeni, J.F., Jr (1993) A cohort study of smoking, alcohol consumption, and dietary factors for pancreatic cancer (United States). *Cancer Causes Control*, **4**, 477–482

- Zheng, T., Cantor, K.P., Zhang, Y., Chiu, B.C.H. & Lynch, C.F. (2001) Risk of brain glioma not associated with cigarette smoking or use of other tobacco products in Iowa. *Cancer Epidemiol. Biomarkers Prev.*, **10**, 413–414
- Zuskin, E., Mustajbegovic, J., Schachter, E.N., Kanceljak, B., Macan, J., Kern, J., Buneta, L. & Pucaric-Cvetkovic, J. (2004) Immunological and respiratory changes in tobacco workers. *Am. J. ind. Med.*, **45**, 76–83