

6. References

- Adams, J.D., LaVoie, E.J., O'Mara-Adams, K.J., Hoffmann, D., Carey, K.D. & Marshall, M.V. (1985a) Pharmacokinetics of N'-nitrosoornicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in laboratory animals. *Cancer Lett.*, **28**, 195–201
- Adams, J.D., LaVoie, E.J. & Hoffmann, D. (1985b) On the pharmacokinetics of tobacco-specific N-nitrosamines in Fischer rats. *Carcinogenesis*, **6**, 509–511
- Adams, J.D., O'Mara-Adams, K.J. & Hoffmann, D. (1987a) Toxic and carcinogenic agents in undiluted mainstream smoke and sidestream smoke of different types of cigarettes. *Carcinogenesis*, **8**, 729–731

- Adams, J.D., Owens-Tucciarone, P. & Hoffmann, D. (1987b) Tobacco-specific N-nitrosamines in dry snuff. *Food chem. Toxicol.*, **25**, 245–246
- Adlkofer, F., Scherer, G., Conze, C., Angerer, J. & Lehnert, G. (1990) Significance of exposure to benzene and other toxic compounds through environmental tobacco smoke. *J. Cancer Res. clin. Oncol.*, **116**, 591–598
- Alaoui-Jamali, M.A., Castonguay, A. & Schuller, H.M. (1988) *In vitro* genotoxicity of 4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone activated by hamster maternal and fetal tissues. *Carcinogenesis*, **9**, 2319–2324
- Amin, S., Desai, D., Hecht, S.S. & Hoffmann, D. (1996) Synthesis of tobacco-specific N-nitrosamines and their metabolites and results of related bioassays. *Crit. Rev. Toxicol.*, **26**, 139–147
- Ammigan, N., Nair, U.J., Amonkar, A.J. & Bhide, S.V. (1989) Modulations in the biotransformation of tobacco extract and *N'*-nitrosonornicotine under differential dietary protein status. *J. Biochem. Toxicol.*, **4**, 7–13
- Ammigan, N., Nair, U.J., Amonkar, A.J. & Bhide, S.V. (1990a) Effect of tobacco extract and *N'*-nitrosonornicotine on the carcinogen metabolising enzymes under different dietary vitamin B status. *Cancer Lett.*, **52**, 153–159
- Ammigan, N., Nagabhushan, M., Nair, U.J., Amonkar, A.J. & Bhide, S.V. (1990b) Effect of nutritional status on mutagenicity of urine excreted by rats treated with standard/experimental carcinogens. *Indian J. exp. Biol.*, **28**, 711–713
- Andersen, R.A., Burton, H.R., Fleming, P.D. & Hamilton-Kemp, T.R. (1989) Effect of storage conditions on nitrosated, acetylated, and oxidized pyridine alkaloid derivatives in smokeless tobacco products. *Cancer Res.*, **49**, 5895–5900
- Anderson, L.M., Hecht, S.S., Kovatch, R.M., Amin, S., Hoffmann, D. & Rice, J.M. (1991) Tumorigenicity of the tobacco-specific carcinogen 4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone in infant mice. *Cancer Lett.*, **58**, 177–181
- Anderson, K.E., Hammons, G.J., Kadlubar, F.F., Potter, J.D., Kaderlik, K.R., Ilett, K.F., Minchin, R.F., Teitel, C.H., Chou, H.-C., Martin, M.V., Guengerich, F.P., Barone, G.W., Lang, N.P. & Peterson, L.A. (1997) Metabolic activation of aromatic amines by human pancreas. *Carcinogenesis*, **18**, 1085–1092
- Anderson, K.E., Carmella, S.G., Ye, M., Bliss, R.L., Le, C., Murphy, L. & Hecht, S.S. (2001) Metabolites of a tobacco-specific lung carcinogen in nonsmoking women exposed to environmental tobacco smoke. *J. natl Cancer Inst.*, **93**, 378–381
- Anderson, K.E., Kliris, J., Murphy, L., Carmella, S.G., Han, S., Link, C., Bliss, R.L., Puumala, S., Murphy, S.E. & Hecht, S.S. (2003) Metabolites of a tobacco-specific lung carcinogen in non-smoking casino patrons. *Cancer Epidemiol. Biomarkers Prev.*, **12**, 1544–1546
- d'Andres, S., Boudoux, R., Renaud, J. & Zuber, J. (2003) TSNA levels in the mainstream smoke of simplified blend prototypes. *Beitr. Tabakforsch. int.*, **20**, 331–340
- Andrews, A.W., Thibault, L.H. & Lijinsky, W. (1978) The relationship between mutagenicity and carcinogenicity of some nitrosamines. *Mutat. Res.*, **51**, 319–326
- Anttila, S., Hukkanen, J., Hakkola, J., Stjernvall, T., Beaune, P., Edwards, R.J., Boobis, A.R., Pelkonen, O. & Raunio, H. (1997) Expression and localization of CYP3A4 and CYP3A5 in human lung. *Am. J. respir. Cell mol. Biol.*, **16**, 242–249
- Aoyama, T., Yamano, S., Guzelian, P.S., Gelboin, H.V. & Gonzalez, F.J. (1990) Five of 12 forms of vaccinia virus-expressed human hepatic cytochrome P450 metabolically activate aflatoxin B₁. *Proc. natl Acad. Sci. USA*, **87**, 4790–4793

- Ashley, D., Beeson, M., Johnson, D., McCraw, J., Richter, P., Pirkle, J., Pechacek, T., Song, S. & Watson, C. (2003) Tobacco-specific nitrosamines in tobacco from U.S. brand and non-U.S. brand cigarettes. *Nicotine Tob. Res.*, **5**, 323–331
- Atawodi, S.E., Lea, S., Nyberg, F., Mukeria, A., Constantinescu, V., Ahrens, W., Brueske-Hohlfeld, I., Fortes, C., Boffetta, P. & Friesen, M.D. (1998) 4-Hydroxy-1-(3-pyridyl)-1-butanone-hemoglobin adducts as biomarkers of exposure to tobacco smoke: Validation of a method to be used in multicenter studies. *Cancer Epidemiol. Biomarkers Prev.*, **7**, 817–821
- Bartsch, H., Malaveille, C., Camus, A.-M., Martel-Planche, G., Brun, G., Hautefeuille, A., Sabadie, N., Barbin, A., Kuroki, T., Drevon, C., Piccoli, C. & Montesano, R. (1980) Bacterial and mammalian mutagenicity tests: Validation and comparative studies on 180 chemicals. In: Montesano, R., Bartsch, H. & Tomatis, L., eds, *Molecular and Cellular Aspects of Carcinogen Screening Tests* (IARC Scientific Publications No. 27), Lyon, pp. 179–241
- Baskaran, K., Laconi, S. & Reddy, M.K. (1994) Transformation of hamster pancreatic duct cells by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), in vitro. *Carcinogenesis*, **15**, 2461–2466
- Becher, R., Lag, M., Schwarze, P.E., Brunborg, G., Soderlund, E.J. & Holme, J.A. (1993) Chemically induced DNA damage in isolated rabbit lung cells. *Mutat. Res.*, **285**, 303–311
- Bedard, L.L., Smith, G.B., Reid, K.R., Petsikas, D. & Massey, T.E. (2002) Investigation of the role of lipoxygenase in bioactivation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in human lung. *Chem. Res. Toxicol.*, **15**, 1267–1273
- Beebe, L.E., Kim, Y.E., Amin, S., Riggs, C.W., Kovatch, R.M. & Anderson, L.M. (1993) Comparison of transplacental and neonatal initiation of mouse lung and liver tumors by N-nitrosodimethylamine (NDMA) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and promotability by a polychlorinated bipheyls mixture (Aroclor 1254). *Carcinogenesis*, **14**, 1545–1548
- Belinsky, S.A., White, C.M., Boucheron, J.A., Richardson, F.C., Swenberg, J.A. & Anderson, M. (1986) Accumulation and persistence of DNA adducts in respiratory tissue of rats following multiple administrations of the tobacco specific carcinogen 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res.*, **46**, 1280–1284
- Belinsky, S.A., White, C.M., Devereux, T.R., Swenberg, J.A. & Anderson, M.W. (1987a) Cell selective alkylation of DNA in rat lung following low dose exposure to the tobacco specific carcinogen 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res.*, **47**, 1143–1148
- Belinsky, S.A., Walker, V.E., Maronpot, R.R., Swenberg, J.A. & Anderson, M.W. (1987b) Molecular dosimetry of DNA adduct formation and cell toxicity in rat nasal mucosa following exposure to the tobacco specific nitrosamine 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone and their relationship to induction of neoplasia. *Cancer Res.*, **47**, 6058–6065
- Belinsky, S.A., Dolan, M.E., White, C.M., Maronpot, R.R., Pegg, A.E. & Anderson, M.W. (1988) Cell specific differences in O⁶-methylguanine–DNA methyltransferase activity and removal of O⁶-methylguanine in rat pulmonary cells. *Carcinogenesis*, **9**, 2053–2058
- Belinsky, S.A., White, C.M., Trushin, N. & Hecht, S.S. (1989a) Cell specificity for the pulmonary metabolism of tobacco-specific nitrosamines in the Fischer rat. *Carcinogenesis*, **10**, 2269–2274
- Belinsky, S.A., Devereux, T.R., Maronpot, R.R., Stoner, G.D. & Anderson, M.W. (1989b) Relationship between the formation of promutagenic adducts and the activation of the K-ras proto-oncogene in lung tumors from A/J mice treated with nitrosamines. *Cancer Res.*, **49**, 5305–5311
- Belinsky, S.A., Foley, J.F., White, C.M., Anderson, M.W. & Maronpot, R.R. (1990) Dose–response relationship between O⁶-methylguanine formation in Clara cells and induction of pulmonary

- neoplasia in the rat by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res.*, **50**, 3772–3780
- Belinsky, S.A., Devereux, T.R., Foley, J.F., Maronpot, R.R. & Anderson, M.W. (1992) Role of the alveolar type II cell in the development and progression of pulmonary tumors induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in the A/J mouse. *Cancer Res.*, **52**, 3164–3173
- Belinsky, S.A., Nikula, K.J., Palmisano, W.A., Michels, R., Saccomanno, G., Gabrielson, E., Baylin, S.B. & Herman, J.G. (1998) Aberrant methylation of *p16^{INK4a}* is an early event in lung cancer and a potential biomarker for early diagnosis. *Proc. natl Acad. Sci. USA*, **95**, 11891–11896
- Bernert, J.T., Jain, R.B., Pirkle, J.L., Wang, L., Miller, B.B. & Sampson, E.J. (2005) Urinary tobacco-specific nitrosamines and 4-aminobiphenyl hemoglobin adducts measured in smokers of either regular or light cigarettes. *Nicotine Tob. Res.*, **7**, 729–738
- Bhide, S.V., Nair, U.J. & Nair, J. (1986) *N*-Nitrosamines in the saliva of tobacco chewers or masheri users. *Food chem. Toxic.*, **24**, 293–297
- Bhide, S.V., Nair, J., Maru, G.B., Nair, U.J., Kameshwar Rao, B.V., Chakraborty, M.K. & Brunnemann, K.D. (1987) Tobacco-specific *N*-nitrosamines [TNSA] in green mature and processed tobacco leaves from India. *Beit. Tabakforsch.*, **14**, 29–32
- Blömeke, B., Greenblatt, M.J., Doan, V.D., Bowman, E.D., Murphy, S.E., Chen, C.C., Kato, S. & Shields, P.G. (1996) Distribution of 7-alkyl-2'-deoxyguanosine adduct levels in human lung. *Carcinogenesis*, **17**, 741–748
- Borgerding, M. & Klus, H. (2005) Analysis of complex mixtures — Cigarette smoke. *Exp. Toxicol. Pathol.*, **57**, 43–73
- Borgerding, M.F., Bodnar, J.A. & Wingate, D.E. (2000) *The 1999 Massachusetts Benchmark Study — Final Report. A Research Study Conducted after Consultation with the Massachusetts Department of Public Health* [http://www.brownandwilliamson.com/APPS/PDF/Final_Report_1999_Mass_Benchmark_Study.pdf]
- Boylard, E., Roe, F.J.C. & Gorrod, J.W. (1964a) Induction of pulmonary tumours in mice by nitrosornicotine, a possible constituent of tobacco smoke. *Nature*, **202**, 1126
- Boylard, E., Roe, F.J.C., Gorrod, J.W. & Mitchley, B.C.V. (1964b) The carcinogenicity of nitrosoanabasine, a possible constituent of tobacco smoke. *Br. J. Cancer*, **18**, 265–270
- Boysen, G., Kenney, P.M.J., Upadhyaya, P., Wang, M. & Hecht, S.S. (2003) Effects of benzyl isothiocyanate and 2-phenethyl isothiocyanate on benzo[a]pyrene and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone metabolism in F-344 rats. *Carcinogenesis*, **24**, 517–525
- Branner, B., Kutzer, C., Zwicknflug, W., Scherer, G., Heller, W.-D. & Richter, E. (1998) Haemoglobin adducts from aromatic amines and tobacco-specific nitrosamines in pregnant smoking and non-smoking women. *Biomarkers*, **3**, 35–47
- Breland, A.B., Acosta, M.C. & Eissenberg, T. (2003) Tobacco specific nitrosamines and potential reduced exposure products for smokers: A preliminary evaluation of Advance™. *Tob. Control*, **12**, 317–321
- Brittebo, E. & Tjälve, H. (1980) Autoradiographic observations on the distribution and metabolism of *N'*-[¹⁴C]nitrosornicotine in mice. *J. Cancer Res. clin. Oncol.*, **98**, 233–242
- Brittebo, E.B. & Tjälve, H. (1981) Formation of tissue-bound *N'*-nitrosornicotine metabolites by the target tissues of Sprague-Dawley and Fisher rats. *Carcinogenesis*, **2**, 959–963
- Brittebo, E.B., Castonguay, A., Furuya, K. & Hecht, S.S. (1983) Metabolism of tobacco-specific nitrosamines by cultured rat nasal mucosa. *Cancer Res.*, **43**, 4343–4348

- Brognaard, J., Clark, A.S., Ni, Y. & Dennis, P.A. (2001) Akt/protein kinase B is constitutively active in non-small cell lung cancer cells and promotes cellular survival and resistance to chemotherapy and radiation. *Cancer Res.*, **61**, 3986–3997
- Brown, B.G., Richter, E., Tricker, A.R., Ayres, P.H. & Doolittle, D.J. (2001a) The effect of a 2-h exposure to cigarette smoke on the metabolic activation of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in A/J mice. *Chem.-biol. Interact.*, **138**, 125–135
- Brown, B., Avalos, J., Lee, C. & Doolittle, D. (2001b) The effect of tobacco smoke, nicotine, and cotinine on the mutagenicity of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL). *Mutat. Res.*, **494**, 21–29
- Brunnemann, K.D. & Hoffmann, D. (1992) Chemical composition of smokeless tobacco products. In: *Smokeless Tobacco or Health. An International Perspective* (Smoking and Tobacco Control Monograph No. 2; NIH Publ. No. 93-3461), Bethesda, MD, National Cancer Institute
- Brunnemann, K.D., Genoble, L. & Hoffmann, D. (1985) N-Nitrosamines in chewing tobacco: An international comparison. *J. agric. Food Chem.*, **33**, 1178–1181
- Brunnemann, K.D., Genoble, L. & Hoffmann, D. (1987a) Identification and analysis of a new tobacco-specific N-nitrosamine, 4-(methylnitrosamino)-4-(3-pyridyl)-1-butanol. *Carcinogenesis*, **8**, 465–469
- Brunnemann, K.D., Rivenson, A., Adams, J.D., Hecht, S.S. & Hoffmann, D. (1987b) A study of snuff carcinogenesis. In: Bartsch, H., O'Neill, I.K. & Schulte-Herman, R., eds, *Relevance of N-Nitroso Compounds to Human Cancer: Exposures and Mechanisms* (IARC Scientific Publications No. 84), Lyon, pp. 456–459
- Brunnemann, K.D., Hornby, A.P. & Stich, H.F. (1987c) Tobacco-specific nitrosamines in the saliva of Inuit snuff dippers in the Northwest territories of Canada. *Cancer Lett.*, **37**, 7–16
- Brunnemann, K.D., Cox, J.E. & Hoffmann, D. (1992) Analysis of tobacco-specific N-nitrosamines in indoor air. *Carcinogenesis*, **13**, 2415–2418
- Brunnemann, K.D., Hoffmann, D., Gairola, C.G. & Lee, B.C. (1994) Low ignition propensity cigarettes: Smoke analysis for carcinogens and testing for mutagenic activity of the smoke particulate matter. *Food chem. Toxicol.*, **32**, 917–922
- Brunnemann, K.D., Mitacek, E.J., Liu, Y., Limsila, T. & Suttajit, M. (1996) Assessment of major carcinogenic tobacco-specific N-nitrosamines in Thai cigarettes. *Cancer Detect. Prev.*, **20**, 114–121
- Brunnemann, K.D., Qi, J. & Hoffmann, D. (2002) Chemical profile of two types of oral snuff tobacco. *Food chem. Toxicol.*, **40**, 1699–1703
- Brunnemann, K.D., Qi, J. & Hoffmann, D. (2004) Levels of TSNA in oral moist snuff in the past 30 years and today (Abstract No. 38). In: *58th Tobacco Science Research Conference Proceedings*, Winston-Salem, NC, Tobacco Science Research
- Burns, D.M., Major, J.M., Shanks, T.G., Thun, M.J. & Samet, J.M. (2001) Smoking lower yield cigarettes and disease risks. In: *Risks Associated with Smoking Cigarettes with Low Tar Machine-measured Yields of Tar and Nicotine* (Smoking and Tobacco Control, Monograph No. 13), Bethesda, MD, National Institutes of Health, National Cancer Institute, pp. 65–158
- Burton, H., Bush, L. & Djordjevic, M. (1989a) Influence of temperature and humidity on the accumulation of tobacco-specific nitrosamines in stored burley tobacco. *J. agric. Food Chem.*, **37**, 1372–1377

- Burton, H.R., Childs, G.H., Jr, Andersen, R.A. & Fleming, P.D. (1989b) Changes in chemical composition of burley tobacco during senescence and curing. 3. Tobacco-specific nitrosamines. *J. agric. Food Chem.*, **37**, 426–430
- Burton, H.R., Dye, N.K. & Bush, L.P. (1992) Distribution of tobacco constituents in tobacco leaf tissue. 1. Tobacco-specific nitrosamines, nitrate, nitrite and alkaloids. *J. agric. Food Chem.*, **40**, 1050–1055
- Bush, L.P., Cui, M., Shi, H., Burton, H.R., Fannin, F.F., Lei, L. & Dye, N. (2001) Formation of tobacco-specific nitrosamines in air-cured tobacco. *Recent Adv. Tob. Sci.*, **27**, 23–46
- Byrd, G.D. & Ogden, M.W. (2003) Liquid chromatographic/tandem mass spectrometric method for the determination of the tobacco-specific nitrosamine metabolite NNAL in smokers' urine. *J. mass Spectrom.*, **38**, 98–107
- Carmella, S.G. & Hecht, S.S. (1987) Formation of hemoglobin adducts upon treatment of F344 rats with the tobacco-specific nitrosamines 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and N'-nitrosonornicotine. *Cancer Res.*, **47**, 2626–2630
- Carmella, S.G., Kagan, S.S., Kagan, M., Foiles, P.G., Palladino, G., Quart, A.M., Quart, E. & Hecht, S.S. (1990a) Mass spectrometric analysis of tobacco-specific nitrosamine hemoglobin adducts in snuff dippers, smokers, and nonsmokers. *Cancer Res.*, **50**, 5438–5445
- Carmella, S.G., Kagan, S.S., Spratt, T.E. & Hecht, S.S. (1990b) Evaluation of cysteine adduct formation in rat hemoglobin by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and related compounds. *Cancer Res.*, **50**, 5453–5459
- Carmella, S.G., Kagan, S.S. & Hecht, S.S. (1992) Evidence that a hemoglobin adduct of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone is a 4-(3-pyridyl)-4-oxobutyl carboxylic acid ester. *Chem. Res. Toxicol.*, **5**, 76–80
- Carmella, S.G., Akerkar, S. & Hecht, S.S. (1993) Metabolites of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in smokers' urine. *Cancer Res.*, **53**, 721–724
- Carmella, S.G., Akerkar, S.A., Richie, J.P., Jr & Hecht, S.S. (1995) Intraindividual and inter-individual differences in metabolites of the tobacco-specific lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in smokers' urine. *Cancer Epidemiol. Biomarkers Prev.*, **4**, 635–642
- Carmella, S.G., Borukhova, A., Akerkar, S.A. & Hecht, S.S. (1997) Analysis of human urine for pyridine-N-oxide metabolites of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, a tobacco-specific lung carcinogen. *Cancer Epidemiol. Biomarkers Prev.*, **6**, 113–120
- Carmella, S.G., Ye, M., Upadhyaya, P. & Hecht, S.S. (1999) Stereochemistry of metabolites of a tobacco-specific lung carcinogen in smokers' urine. *Cancer Res.*, **59**, 3602–3605
- Carmella, S.G., Le, K.-A., Upadhyaya, P. & Hecht, S.S. (2002) Analysis of N- and O-glucuronides of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in human urine. *Chem. Res. Toxicol.*, **15**, 545–550
- Carmella, S.G., Han, S., Frisstad, A., Yang, Y. & Hecht, S.S. (2003) Analysis of total 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in human urine. *Cancer Epidemiol. Biomarkers Prev.*, **12**, 1257–1261
- Carmella, S.G., Han, S.H., Villalta, P.W. & Hecht, S.S. (2005) Analysis of total 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in smokers' blood. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 2669–2672

- Carmella, S.G., Yoder, A. & Hecht, S.S. (2006) Combined analysis of r-1,t-2,3-c-4-tetrahydroxy-1,2,3,4-tetrahydrophenanthrene and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in smokers' plasma. *Cancer Epidemiol. Biomarkers Prev.*, **15**, 1490–1494
- Castonguay, A., Lin, D., Stoner, G.D., Radok, P., Furuya, K., Hecht, S.S., Schut, H.A.J. & Klaunig, J.E. (1983a) Comparative carcinogenicity in A/J mice and metabolism by cultured mouse peripheral lung of *N'*-nitrosornicotine, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, and their analogues. *Cancer Res.*, **43**, 1223–1229
- Castonguay, A., Tjälve, H. & Hecht, S.S. (1983b) Tissue distribution of the tobacco-specific carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and its metabolites in F344 rats. *Cancer Res.*, **43**, 630–638
- Castonguay, A., Stoner, G.D., Schut, H.A. & Hecht, S.S. (1983c) Metabolism of tobacco-specific *N*-nitrosamines by cultured human tissues. *Proc. natl Acad. Sci. USA*, **80**, 6694–6697
- Castonguay, A., Rivenson, A., Trushin, N., Reinhardt, J., Spathopoulos, S., Weiss, C.J., Reiss, B. & Hecht, S.S. (1984a) Effects of chronic ethanol consumption on the metabolism and carcinogenicity of *N'*-nitrosornicotine in F344 rats. *Cancer Res.*, **44**, 2285–2290
- Castonguay, A., Tharp, R. & Hecht, S.S. (1984b) Kinetics of DNA methylation by the tobacco-specific carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in F344 rats. In: O'Neill, I.K., von Borstel, R.C., Miller, C.T., Long, J. & Bartsch, H., eds, *N-Nitroso Compounds: Occurrence, Biological Effects and Relevance to Human Cancer* (IARC Scientific Publications No. 57), Lyon, pp. 805–810
- Castonguay, A., Tjälve, H., Trushin, N., d'Argy, R. & Sperber, G. (1985a) Metabolism and tissue distribution of tobacco-specific *N*-nitrosamines in the marmoset monkey (*Callithrix jacchus*). *Carcinogenesis*, **6**, 1543–1550
- Castonguay, A., Foiles, P.G., Trushin, N. & Hecht, S.S. (1985b) Study of DNA methylation by tobacco-specific *N*-nitrosamines. *Environ. Health Perspect.*, **62**, 197–202
- Castonguay, A., Pepin, P. & Briere, N. (1991) Modulation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone demethylation and denitrosation by rat liver microsomes. *Cancer Lett.*, **59**, 67–74
- Chakradeo, P.P., Nair, J. & Bhide, S.V. (1995) Metabolism of *N'*-nitrosornicotine by adult and fetal human oesophageal cultures. *Cell Biol. int.*, **19**, 53–58
- Chamberlain, W.J. & Chortyk, O.T. (1992) Effects of curing and fertilization on nitrosamine formation in bright and burley tobacco. *Beitr. Tabakforsch.*, **15**, 87–92
- Chamberlain, W.J., Schlotzhauer, W.S. & Chortyk, O.T. (1988) Chemical composition of non-smoking tobacco products. *J. agric. Food Chem.*, **36**, 48–50
- Chen, C.-H.B., Hecht, S.S. & Hoffmann, D. (1978) Metabolic α -hydroxylation of the tobacco-specific carcinogen, *N'*-nitrosornicotine. *Cancer Res.*, **38**, 3639–3645
- Chen, B., Liu, L., Castonguay, A., Maronpot, R.R., Anderson, M.W. & You, M. (1993) Dose-dependent *ras* mutation spectra in *N*-nitrosodiethylamine induced mouse liver tumors and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone induced mouse lung tumors. *Carcinogenesis*, **14**, 1603–1608
- Chen, Y.-P., Johnson, G.K. & Squier, C.A. (1994) Effects of nicotine and tobacco-specific nitrosamines on hamster cheek pouch and gastric mucosa. *J. oral Pathol. Med.*, **23**, 251–255
- Chung, F.-L. & Xu, Y. (1992) Increased 8-oxodeoxyguanosine levels in lung DNA of A/J mice and F344 rats treated with the tobacco-specific nitrosamine 4-(methylnitrosamine)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*, **13**, 1269–1272

- Connolly, G.N. (2001) *Establishing Tolerance Limits for Tobacco Specific Nitrosamines (TSNAs) in Oral Snuff under the Massachusetts Hazardous Substance Act Mg L c 94 s-*, Boston, Massachusetts Department of Public Health
- Correa, E., Joshi, P.A., Castonguay, A. & Schuller, H.M. (1990) The tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone is an active transplacental carcinogen in Syrian golden hamsters. *Cancer Res.*, **50**, 3435–3438
- Counts, M.E., Morton, M.J., Laffoon, S.W., Cox, R.H. & Lipowicz, P.J. (2005) Smoke composition and predicting relationships for international commercial cigarettes smoked with three machine-smoking conditions. *Regul. Toxicol. Pharmacol.*, **41**, 185–227
- Dahl, J.E., Becher, R., Låg, M., Schuller, H.M. & Dybing, E. (1990) Formation of genotoxic products from N-nitrosoheptamethyleneimine (NHMI), 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and N'-nitrosornicotine (NNN) by isolated rabbit lung cells. *Cell Biol. Toxicol.*, **6**, 399–409
- Demkowicz-Dobrzanski, K. & Castonguay, A. (1991) Comparison of DNA alkali-labile sites induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and 4-oxo-4-(3-pyridyl)butanal in rat hepatocytes. *Carcinogenesis*, **12**, 2135–2140
- DeRoton, C., Wiernik, A., Wahlberg, I. & Vidal, B. (2005) Factors influencing the formation of tobacco-specific nitrosamines in French air-cured tobaccos in trials and at the farm level. *Beitr. Tabakforsch. int.*, **21**, 305–320
- Desai, D., Kagan, S.S., Amin, S., Carmella, S.G. & Hecht, S.S. (1993) Identification of 4-(methylnitrosamino)-1-[3-(6-hydroxypyridyl)]-1-butanone as a urinary metabolite of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in rodents. *Chem. Res. Toxicol.*, **6**, 794–799
- Deutsch-Wenzel, R.P., Brune, H., Grimmer, G. & Misfeld, J. (1985) Local application to mouse skin as a carcinogen specific test system for non-volatile nitroso compounds. *Cancer Lett.*, **29**, 85–92
- Devereux, T.R., Anderson, M.W. & Belinsky, S.A. (1988) Factors regulating activation and DNA alkylation by 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone and nitrosodimethylamine in rat lung and isolated lung cells, and the relationship to carcinogenicity. *Cancer Res.*, **48**, 4215–4221
- Devereux, T.R., Anderson, M.W. & Belinsky, S.A. (1991) Role of *ras* protooncogene activation in the formation of spontaneous and nitrosamine-induced lung tumors in the resistant C3H mouse. *Carcinogenesis*, **12**, 299–303
- Devereux, T.R., Belinsky, S.A., Maronpot, R.R., White, C.M., Hegi, M.E., Patel, A.C., Foley, J.F., Greenwell, A. & Anderson, M.W. (1993) Comparison of pulmonary O⁶-methylguanine DNA adduct levels and Ki-*ras* activation in lung tumors from resistant and susceptible mouse strains. *Mol. Carcinog.*, **8**, 177–185
- Dicke, K.E., Skrlin, S.M. & Murphy, S.E. (2005) Nicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-butanone metabolism by cytochrome P450 2B6. *Drug Metab. Dispos.*, **33**, 1760–1764
- Ding, X. & Kaminsky, L.S. (2003) Human extrahepatic cytochromes P450: Function in xenobiotic metabolism and tissue-selective chemical toxicity in the respiratory and gastrointestinal tracts. *Ann. Rev. Pharmacol. Toxicol.*, **43**, 149–173
- Djordjevic, M.V., Brunnemann, K.D. & Hoffmann, D. (1989a) Identification and analysis of a nicotine-derived N-nitrosamino acid and other nitrosamino acids in tobacco. *Carcinogenesis*, **10**, 1725–1731

- Djordjevic, M.V., Gay, S.L., Bush, L.P. & Chaplin, J.F. (1989b) Tobacco-specific nitrosamine accumulation and distribution in flue-cured tobacco alkaloid isolines. *J. agric. Food Chem.*, **37**, 752–756
- Djordjevic, M.V., Sigountos, C.W., Brunnemann, K.D. & Hoffmann, D. (1990) Tobacco-specific nitrosamine delivery in the mainstream smoke of high- and low-yield cigarettes smoked with varying puff volume. In: *CORESTA Symposium Proceedings, Smoke Study Group, Kallithea, Greece*, pp. 54–62
- Djordjevic, M.V., Sigountos, C.W., Hoffmann, D., Brunnemann, K.D., Kagan, M.R., Bush, L.P., Safaev, R., Belitsky, G. & Zaridze, D. (1991) Assessment of major carcinogens and alkaloids in the tobacco and mainstream smoke of USSR cigarettes. *Int. J. Cancer*, **47**, 348–351
- Djordjevic, M.V., Fan, J., Bush, L.P., Brunnemann, K.D. & Hoffmann, D. (1993a) Effects of storage conditions on levels of tobacco-specific N-nitrosamines and N-nitrosamino acids in U.S. moist snuff. *J. agric. Food Chem.*, **41**, 1790–1794
- Djordjevic, M.V., Brunnemann, K.D. & Hoffmann, D. (1993b) The need for regulation of carcinogenic N-nitrosamines in oral snuff. *Food chem. Toxicol.*, **41**, 497–501
- Djordjevic, M.V., Eixarch, L., Bush, L.P. & Hoffmann, D. (1996) A comparison of the yields of selected components in the mainstream smoke of the leading US and Japanese cigarettes. In: *CORESTA Congress Proceedings, Joint Smoke and Technology Groups, Yokohama, Japan*, pp. 200–217
- Djordjevic, M.V., Eixarch, L. & Hoffmann, D. (1997) Self-administered and effective dose of cigar smoke constituents. In: *Proceedings of the 51st Tobacco Chemists' Research Conference, Winston-Salem, NC, September 14–17, 1997*
- Djordjevic, M.V., Stellman, S.D., Takezaki, T. & Tajima, K. (2000a) Cigarette composition as a possible explanation of US–Japan differences in lung cancer rates (Abstract No. 5120). In: *Proceedings of the 91st Annual Meeting of the American Association for Cancer Research, Vol. 41, San Francisco, CA, April 2000*
- Djordjevic, M.V., Prokopczyk, B. & Zatonski, W. (2000b) Tobacco-specific N-nitrosamines (TSNA) in tobacco and mainstream smoke of the leading Polish cigarettes: A comparison with the US and Japanese brands. In: *Proceedings of the CORESTA Congress, Smoke Study Group, STI, Lisbon, Portugal, October 2000*
- Djordjevic, M.V., Stellman, S.D. & Zang, E. (2000c) Doses of nicotine and lung carcinogens delivered to cigarette smokers. *J. natl Cancer Inst.*, **92**, 106–111
- Domellöf, L., Andersson, M., Tjälve, H., Veals, S., Trushin, N. & Hecht, S.S. (1987) Distribution and metabolism of N'-nitrosornicotine in the miniature pig. *Carcinogenesis*, **8**, 1741–1747
- Du, X., Squier, C.A., Kremer, M.J. & Wertz, P.W. (2000) Penetration of N-nitrosornicotine (NNN) across oral mucosa in the presence of ethanol and nicotine. *J. oral Pathol. Med.*, **29**, 80–85
- Egan, H., Preussmann, R., O'Neill, I.K., Eisenbrand, G., Spiegelhalter, B. & Bartsch, H., eds (1983) *Environmental Carcinogens. Selected Methods of Analysis*, Vol. 6, N-Nitroso Compounds (IARC Scientific Publications No. 45), Lyon
- Ellinger-Ziegelbauer, H., Stuart, B., Wahle, B., Bomann, W. & Ahr, H.J. (2004) Characteristic expression profiles induced by genotoxic carcinogens in rat liver. *Toxicol Sci.*, **77**, 19–34
- Falter, B., Kutzer, C. & Richter, E. (1994) Biomonitoring of hemoglobin adducts: Aromatic amines and tobacco-specific nitrosamines. *Clin. Invest.*, **72**, 364–371

- Felicia, N.D., Rekha, G.K. & Murphy, S.E. (2000) Characterization of cytochrome P450 2A4 and 2A5-catalyzed 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) metabolism. *Arch. Biochem. Biophys.*, **384**, 418–424
- Finch, G.L., Nikula, K.J., Belinsky, S.A., Barr, E.B., Stoner, G.D. & Lechner, J.F. (1996) Failure of cigarette smoke to induce or promote lung cancer in the A/J mouse. *Cancer Lett.*, **99**, 161–167
- Finckh, C., Atalla, A., Nagel, G., Stinner, B. & Maser, E. (2001) Expression and NNK reducing activities of carbonyl reductase and 11beta-hydroxysteroid dehydrogenase type 1 in human lung. *Chem.-biol. Interact.*, **130–132**, 761–773
- Fischer, S. & Spiegelhalder, B. (1989) Improved method for the determination of tobacco-specific nitrosamines (TSNA) in tobacco smoke. *Beitr. Tabakforsch. int.*, **14**, 145–153
- Fischer, S., Spiegelhalder, B. & Preussmann, R. (1989a) Preformed tobacco-specific nitrosamines in tobacco — Role of nitrate and influence of tobacco type. *Carcinogenesis*, **10**, 1511–1517
- Fischer, S., Spiegelhalder, B. & Preussmann, R. (1989b) Tobacco-specific nitrosamines in mainstream smoke of West German cigarettes — Tar alone is not a sufficient index for the carcinogenic potential of cigarette smoke. *Carcinogenesis*, **10**, 169–173
- Fischer, S., Spiegelhalder, B. & Preussmann, R. (1990a) Tobacco-specific nitrosamines in European and USA cigarettes. *Arch. Geschwulstforsch.*, **60**, 169–177
- Fischer, S., Castonguay, A., Kaiserman, M., Spiegelhalder, B. & Preussmann, R. (1990b) Tobacco-specific nitrosamines in Canadian cigarettes. *J. Cancer Res. clin. Oncol.*, **116**, 563–568
- Fischer, S., Spiegelhalder, B., Eisenbarth, J. & Preussmann, R. (1990c) Investigations on the origin of tobacco-specific nitrosamines in mainstream smoke of cigarettes. *Carcinogenesis*, **11**, 723–730
- Foiles, P.G., Akerkar, S.A., Carmella, S.G., Kagan, M., Stoner, G.D., Resau, J.H. & Hecht, S.S. (1991) Mass spectrometric analysis of tobacco-specific nitrosamine–DNA adducts in smokers and nonsmokers. *Chem. Res. Toxicol.*, **4**, 364–368
- Foiles, P.G., Miglietta, L.M., Akerkar, S.A., Everson, R.B. & Hecht, S.S. (1988) Detection of O⁶-methyldeoxyguanosine in human placental DNA. *Cancer Res.*, **48**, 4184–4188
- Foiles, P.G., Peterson, L.A., Miglietta, L.M. & Ronai, Z. (1992) Analysis of mutagenic activity and ability to induce replication of polyoma DNA sequences by different model metabolites of the carcinogenic tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Mutat. Res.*, **279**, 91–101
- Fujita, K.-I. & Kamataki, T. (2001) Predicting the mutagenicity of tobacco-related N-nitrosamines in humans using 11 strains of *Salmonella typhimurium* YG7108, each coexpressing a form of human cytochrome P450 along with NADPH-cytochrome P450 reductase. *Environ. mol. Mutag.*, **38**, 339–346
- Furukawa, J., Nishikawa, A., Enami, T., Mitsui, M., Imazawa, T., Tanakamasu, Z., Kim, H.-C., Lee, I.-S., Kasahara, K. & Takahashi, M. (1997) Promotional effects of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) on N-nitrosobis(2-oxopropyl)amine (BOP)-initiated carcinogenesis in hamsters. *Food chem. Toxicol.*, **35**, 387–392
- Georgiadis, P., Samoli, E., Kaila, S., Katsouyanni, K. & Kyrtopoulos, S.A. (2000) Ubiquitous presence of O⁶-methylguanine in human peripheral and cord blood DNA. *Cancer Epidemiol. Biomarkers Prev.*, **9**, 299–305
- Gerde, P., Muggenburg, B.A., Stephens, T., Lewis, J.L., Pyon, K.H. & Dahl, A.R. (1998) A relevant dose of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone is extensively metabolized and rapidly absorbed in the canine tracheal mucosa. *Cancer Res.*, **58**, 1417–1422

- Gervot, L., Rochat, B., Gautier, J.C., Bohnenstengel, F., Kroemer, H., de Berardinis, V., Martin, H., Beaune, P. & de Waziers, I. (1999) Human CYP2B6: Expression, inducibility and catalytic activities. *Pharmacogenetics*, **9**, 295–306
- Gopalakrishnan, R., Morse, M.A., Lu, J., Weghorst, C.M., Sabourin, C.L.K., Stoner, G.D. & Murphy, S.E. (1999) Expression of cytochrome P450 2A3 in rat esophagus: Relevance to N-nitrosobenzylmethylamine. *Carcinogenesis*, **20**, 885–891
- Gopalakrishnan, R., Gupta, A., Carlton, P.S., Morse, M.A. & Stoner, G.D. (2002) Functional role of cytochrome P-450 2A3 in N-nitrosomethylbenzylamine metabolism in rat esophagus. *J. Toxicol. environ. Health*, **65**, 1077–1091
- Gothia (2004) [Available at <http://www.gothiatek.com/templates>]
- Government of British Columbia (2002) *What is in Cigarettes? Mainstream Smoke and Sidestream Smoke Chemical Constituents by Cigarette Brand*. [<http://www.healthplanning.gov.bc.ca/ttdr/index.html>]
- Gray, N., Boyle, P. & Zatonski, W. (1998) Tar concentrations in cigarettes and carcinogen content. *Lancet*, **352**, 787–788
- Gray, N., Zaridze, D., Robertson, C., Krivosheeva, L., Sigacheva, N., Boyle, P. & the International Cigarette Variation Group (2000) Variation within global cigarette brands in tar, nicotine, and certain nitrosamines: Analytic study. *Tob. Control*, **9**, 351
- Gregg, E., Hill, C., Hollywood, M., Kearney, M., McAdam., McLaughlin, D., Purkis, S. & Williams, M. (2005) The UK smoke constituents testing study. Summary of results and comparison with other studies. *Beitr. Tabakforsch. int./Control Tob. Res.*, **21**, 117–138
- Guengerich, F.P. (1993) Purification and characterization of lung enzymes involved in xenobiotic metabolism. In: Gram, T.E., ed., *Metabolic Activation and Toxicity of Chemical Agents in Lung Tissue and Cells*, Oxford, Pergamon Press, pp. 77–87
- Guo, Z., Smith, T.J., Ishizaki, H. & Yang, C.S. (1991a) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) by cytochrome P450IIB1 in a reconstituted system. *Carcinogenesis*, **12**, 2277–2282
- Guo, Z., Smith, T.J., Thomas, P.E. & Yang, C.S. (1991b) Metabolic activation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone as measured by DNA alkylation *in vitro* and its inhibition by isothiocyanates. *Cancer Res.*, **51**, 4798–4803
- Guo, Z., Smith, T.J., Thomas, P.E. & Yang, C.S. (1992) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by inducible and constitutive cytochrome P450 enzymes in rats. *Arch. Biochem. Biophys.*, **298**, 279–286
- Gupta, P. (2004) *Laboratory Testing of Smokeless Tobacco Products. Final Report to the India Office of the WHO* (Allotment No. SE IND TOB 001.RB.02), New Delhi
- Gupta, P.C. & Ray, C.C. (2003) Smokeless tobacco and health in India and South Asia. *Respirology*, **8**, 419–431
- Haglund, J., Henderson, A.P., Golding, B.T. & Törnqvist, M. (2002) Evidence for phosphate adducts in DNA from mice treated with 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Chem. Res. Toxicol.*, **15**, 773–779
- Harris, J.E., Thun, M.J., Mondul, A.M. & Calle, E.E. (2004) Cigarette tar yields in relation to mortality from lung cancer in the Cancer Prevention Study II prospective cohort, 1982–8. *Br. med. J.*, **328**, 72–79

- Hashimoto, K., Ohsawa, K. & Kimura, M. (2004) Mutations induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in the *lacZ* and *cII* genes of MutaTMMouse. *Mutat. Res.*, **560**, 119–131
- Hatsukami, D.K., Lemmonds, C., Zhang, Y., Murphy, S.E., Le, C., Carmella, S.G. & Hecht, S.S. (2004) Evaluation of carcinogen exposure in people who used 'reduced exposure' tobacco products. *J. natl Cancer Inst.*, **96**, 844–852
- Hecht, S.S. (1998) Biochemistry, biology, and carcinogenicity of tobacco-specific N-nitrosamines. *Chem. Res. Toxicol.*, **11**, 559–603
- Hecht, S.S. (2002) Human urinary carcinogen metabolites: Biomarkers for investigating tobacco and cancer. *Carcinogenesis*, **23**, 907–922
- Hecht, S.S. (2003) Tobacco carcinogens, their biomarkers and tobacco-induced cancer. *Nat. Rev. Cancer*, **3**, 733–744
- Hecht, S.S. & Lin, D. (1986) Comparative mutagenicity of 4-(carbethoxynitrosamino)-4-(3-pyridyl)butanal and 4-(carbethoxynitrosamino)-1-(3-pyridyl)-1-butanone, model compounds for alpha-hydroxylation of *N'*-nitrosornicotine. *Carcinogenesis*, **7**, 611–614
- Hecht, S.S. & Trushin, N. (1988) DNA and hemoglobin alkylation by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and its major metabolite 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in F344 rats. *Carcinogenesis*, **9**, 1665–1668
- Hecht, S.S. & Young, R. (1982) Regiospecificity in the metabolism of the homologous cyclic nitrosamines, *N'*-nitrosornicotine and *N'*-nitrosoanabasine. *Carcinogenesis*, **3**, 1195–1199
- Hecht, S.S., Chen, C.-H.B., Dong, M., Ornaf, R.M. & Hoffmann, D. (1977) Chemical studies on tobacco smoke. LI. Studies on non-volatile nitrosamines in tobacco. *Beitr. Tabakforsch. int.*, **9**, 1–6
- Hecht, S.S., Chen, C.B., Hirota, N., Ornaf, R.M., Tso, T.C. & Hoffmann, D. (1978) Tobacco-specific nitrosamines: Formation from nicotine *in vitro* and during tobacco curing and carcinogenicity in Strain A mice. *J. natl Cancer Inst.*, **60**, 819–824
- Hecht, S.S., Chen, C.-H.B., Ohmori, T. & Hoffmann, D. (1980a) Comparative carcinogenicity in F344 rats of the tobacco-specific nitrosamines, *N'*-nitrosornicotine and 4-(*N*-methyl-*N*-nitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res.*, **40**, 298–302
- Hecht, S.S., Chen, C.-H.B. & Hoffmann, D. (1980b) Metabolic β -hydroxylation and N-oxidation of *N'*-nitrosornicotine. *J. med. Chem.*, **23**, 1175–1178
- Hecht, S.S., Reiss, B., Lin, D. & Williams, G.M. (1982) Metabolism of *N'*-nitrosornicotine by cultured rat esophagus. *Carcinogenesis*, **3**, 453–456
- Hecht, S.S., Adams, J.D., Numoto, S. & Hoffmann, D. (1983a) Induction of respiratory tract tumors in Syrian golden hamsters by a single dose of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and the effect of smoke inhalation. *Carcinogenesis*, **4**, 1287–1290
- Hecht, S.S., Young, R. & Maura, Y. (1983b) Comparative carcinogenicity in F344 rats and Syrian golden hamsters of *N'*-nitrosornicotine and *N'*-nitrosornicotine-1-*N*-oxide. *Cancer Lett.*, **20**, 333–340
- Hecht, S.S., Lin, D. & Castonguay, A. (1983c) Effects of α -deuterium substitution on the mutagenicity of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Carcinogenesis*, **4**, 305–310
- Hecht, S.S., Trushin, N., Castonguay, A. & Rivenson, A. (1986a) Comparative tumorigenicity and DNA methylation in F344 rats by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and *N*-nitrosodimethylamine. *Cancer Res.*, **46**, 498–502

- Hecht, S.S., Rivenson, A., Braley, J., DiBello, J., Adams, J.D. & Hoffmann, D. (1986b) Induction of oral cavity tumors in F344 rats by tobacco-specific nitrosamines and snuff. *Cancer Res.*, **46**, 4162–4166
- Hecht, S.S., Lin, D., Castonguay, A. & Rivenson, A. (1987) Effects of alpha-deuterium substitution on the tumorigenicity of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in F344 rats. *Carcinogenesis*, **8**, 291–294
- Hecht, S.S., Abbaspour, A. & Hoffmann, D. (1988a) A study of tobacco carcinogenesis. XLII. Bioassay in A/J mice of some structural analogues of tobacco-specific nitrosamines. *Cancer Lett.*, **42**, 141–145
- Hecht, S.S., Spratt, T.E. & Trushin, N. (1988b) Evidence for 4-(3-pyridyl)-4-oxobutylation of DNA in F344 rats treated with the tobacco-specific nitrosamines 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and N'-nitrosornicotine. *Carcinogenesis*, **9**, 161–165
- Hecht, S.S., Morse, M.A., Amin, S., Stoner, G.D., Jordan, K.G., Choi, C.I. & Chung, F.L. (1989) Rapid single-dose model for lung tumor induction in A/J mice by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and the effect of diet. *Carcinogenesis*, **10**, 1901–1904
- Hecht, S.S., Jordan, K.G., Choi, C.-I. & Trushin, N. (1990) Effects of deuterium substitution on the tumorigenicity of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in A/J mice. *Carcinogenesis*, **11**, 1017–1020
- Hecht, S.S., Carmella, S.G., Murphy, S.E., Akerkar, S., Brunnemann, K.D. & Hoffmann, D. (1993a) A tobacco-specific lung carcinogen in the urine of men exposed to cigarette smoke. *New Engl. J. Med.*, **329**, 1543–1546
- Hecht, S.S., Trushin, N., Reid-Quinn, C.A., Burak, E.S., Jones, A.B., Southers, J.L., Gombar, C.T., Carmella, S.G., Anderson, L.M. & Rice, J.M. (1993b) Metabolism of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in the patas monkey: Pharmacokinetics and characterization of glucuronide metabolites. *Carcinogenesis*, **14**, 229–236
- Hecht, S.S., Chung, F.L., Richie, J.P., Jr, Akerkar, S.A., Borukhova, A., Skowronski, L. & Carmella, S.G. (1995) Effects of watercress consumption on metabolism of a tobacco-specific lung carcinogen in smokers. *Cancer Epidemiol. Biomarkers Prev.*, **4**, 877–884
- Hecht, S.S., Trushin, N., Rigotty, J., Carmella, S.G., Borukhova, A., Akerkar, S., Desai, D., Amin, S. & Rivenson, A. (1996a) Inhibitory effects of 6-phenylhexyl isothiocyanate on 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone metabolic activation and lung tumorigenesis in rats. *Carcinogenesis*, **17**, 2061–2067
- Hecht, S.S., Trushin, N., Rigotty, J., Carmella, S.G., Borukhova, A., Akerkar, S. & Rivenson, A. (1996b) Complete inhibition of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced rat lung tumorigenesis and favorable modification of biomarkers by phenethyl isothiocyanate. *Cancer Epidemiol. Biomarkers Prev.*, **5**, 645–652
- Hecht, S.S., Carmella, S.G., Chen, M., Dor Koch, J.F., Miller, A.T., Murphy, S.E., Jensen, J.A., Zimmerman, C.L. & Hatsukami, D.K. (1999a) Quantitation of urinary metabolites of a tobacco-specific lung carcinogen after smoking cessation. *Cancer Res.*, **59**, 590–596
- Hecht, S.S., Hatsukami, D.K., Bonilla, L.E. & Hochalter, J.B. (1999b) Quantitation of 4-oxo-4-(3-pyridyl)butanoic acid and enantiomers of 4-hydroxy-4-(3-pyridyl)butanoic acid in human urine: A substantial pathway of nicotine metabolism. *Chem. Res. Toxicol.*, **12**, 172–179
- Hecht, S.S., Kenney, P.M.J., Wang, M., Trushin, N., Agarwal, S. Rao, A.V. & Upadhyaya, P. (1999c) Evaluation of butylated hydroxyanisole, myo-inositol, curcumin, esculetin, resveratrol, and lyco-

- pene as inhibitors of benzo[*a*]pyrene plus 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung tumorigenesis in A/J mice. *Cancer Lett.*, **137**, 123–130
- Hecht, S.S., Trushin, N., Chhabra, S.K., Anderson, L.M. & Nerurkar, P.V. (2000) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by cultured monkey lung explants. *Drug Metab. Dispos.*, **28**, 5–9
- Hecht, S.S., Ye, M., Carmella, S.G., Fredrickson, A., Adgate, J.L., Greaves, I.A., Church, T.R., Ryan, A.D., Mongin, S.J. & Sexton, K. (2001) Metabolites of a tobacco-specific lung carcinogen in the urine of elementary school-aged children. *Cancer Epidemiol. Biomarkers Prev.*, **10**, 1109–1116
- Hecht, S.S., Carmella, S.G., Ye, M., Le, K.-A., Jensen, J.A., Zimmerman, C.L. & Hatsukami, D.K. (2002) Quantitation of metabolites of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone after cessation of smokeless tobacco use. *Cancer Res.*, **62**, 129–134
- Hecht, S.S., Murphy, S.E., Carmella, S.G., Zimmerman, C.L., Losey, L., Kramarczuk, I., Roe, M.R., Puumala, S.S., Li, Y.S., Le, C., Jensen, J. & Hatsukami, D.K. (2004a) Effects of reduced cigarette smoking on the uptake of a tobacco-specific lung carcinogen. *J. natl Cancer Inst.*, **96**, 107–115
- Hecht, S.S., Carmella, S.G., Kenney, P.M.J., Low, S.H., Arakawa, K. & Yu, M.C. (2004b) Effects of cruciferous vegetable consumption on urinary metabolites of the tobacco-specific lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in Singapore chinese. *Cancer Epidemiol. Biomarkers Prev.*, **13**, 997–1004
- Hecht, S.S., Villalta, P.W., Sturla, S.J., Cheng, G., Yu, N., Upadhyaya, P. & Wang, M. (2004c) Identification of O²-substituted pyrimidine adducts formed in reactions of 4-(acetoxymethylnitrosamino)-1-(3-pyridyl)-1-butanone and 4-(acetoxymethylnitrosamino)-1-(3-pyridyl)-1-butanol with DNA. *Chem. Res. Toxicol.*, **17**, 588–597
- Hecht, S.S., Murphy, S.E., Carmella, S.G., Jensen, J., Le, C., Joseph, A.M. & Hatsukami, D.K. (2005) Similar uptake of lung carcinogens by smokers of regular, light, and ultralight cigarettes. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 693–698
- Hecht, S.S., Carmella, S.G., Le, K.-A., Murphy, S.E., Boettcher, A.J., Le, C., Koopmeiners, J., An, L. & Hennrikus, D.J. (2006) 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanol and its glucuronides in the urine of infants exposed to environmental tobacco smoke. *Cancer Epidemiol. Biomarkers Prev.*, **15**, 988–992
- Hecht, S.S., Carmella, S.G., Murphy, S.E., Riley, W.T., Le, C., Luo, X., Mooney, M. & Hatsukami, D.K. (2007) Similar exposure to a tobacco-specific carcinogen in smokeless tobacco users and cigarette smokers. *Cancer Epidemiol. Biomarkers Prev.*, **16**, 1567–1572
- Heuman, D.M., Gallagher, E.J., Barwick, J.L., Elshourbagy, N.A. & Guzelian, P.S. (1982) Immunological evidence for induction of a common form of hepatic cytochrome P-450 in rats treated with pregnenolone-16 alpha-carbonitrile or other steroidal or non-steroidal agents. *Mol. Pharmacol.*, **21**, 753–760
- Hilfrich, J., Hecht, S.S. & Hoffmann, D. (1977) A study of tobacco carcinogenesis. XV. Effects of *N'*-nitrosornicotine and *N'*-nitrosoanabasine in Syrian golden hamsters. *Cancer Lett.*, **2**, 169–176
- Hoffmann, D. & Adams, J.D. (1981) Carcinogenic tobacco-specific *N*-nitrosamines in snuff and in the saliva of snuff dippers. *Cancer Res.*, **41**, 4305–4308
- Hoffmann, D. & Hoffmann, I. (2001) The changing cigarette: Chemical studies and bioassays. In: *Risks Associated with Smoking Cigarettes with Low Machine-Measured Yields of Tar and*

- Nicotine* (Smoking and Tobacco Control Monograph No. 13; NIH Publ. No. 02-5074), Bethesda, MD, National Cancer Institute, pp. 159–191
- Hoffmann, D., Raineri, R. Hecht, S.S., Maronpot, R. & Wynder, E.L. (1975) A study of tobacco carcinogenesis. XIV. Effects of *N'*-nitrosornicotine and *N'*-nitrosonanabasine in rats. *J. natl Cancer Inst.*, **55**, 977–981
- Hoffmann, D., Adams, J.D., Piade, J.J. & Hecht, S.S. (1980) Chemical studies on tobacco smoke. LXVIII. Analysis of volatile and tobacco-specific nitrosamines in tobacco products. In: Walker, E.A, Castegnaro, M., Grieciute, L., & Börzsönyi, M., eds, *N-Nitrosocompounds: Analysis, Formation and Occurrence* (IARC Scientific Publications No. 31), Lyon, pp. 507–516
- Hoffmann, D., Castonguay, A., Rivenson, A. & Hecht, S.S. (1981) Comparative carcinogenicity and metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and *N'*-nitrosornicotine in Syrian golden hamsters. *Cancer Res.*, **41**, 2386–2393
- Hoffmann, D., Rivenson, A., Amin, S. & Hecht, S.S. (1984) Dose–response study of the carcinogenicity of tobacco-specific N-nitrosamines in F344 rats. *J. Cancer Res. clin. Oncol.*, **108**, 81–86
- Hoffmann, D., Brunneemann, K.D. & Venitt, S. (1988) Carcinogenic nitrosamines in oral snuff. *Lancet*, **i**, 1232
- Hoffmann, D., Djordjevic, M.V. & Brunneemann, K.D. (1991) New brands of oral snuff. *Food chem. Toxicol.*, **29**, 65–68
- Hoffmann, D., Djordjevic, M., Rivenson, A., Zang, E., Desai, D. & Amin, S. (1993a) A study of tobacco carcinogenesis. LI. Relative potencies of tobacco-specific N-nitrosamines as inducers of lung tumours in A/J mice. *Cancer Letters*, **71**, 25–30
- Hoffmann, D., Rivenson, A., Abbi, R. & Wynder, E.L. (1993b) A study of tobacco carcinogenesis: effect of the fat content of the diet on the carcinogenic activity of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in F344 rats. *Cancer Res.*, **53**, 2758–2761
- Hoffmann, D., Brunneemann, K.D., Prokopczyk, B. & Djordjevic, M.V. (1994) Tobacco-specific N-nitrosamines and areca-derived N-nitrosamines: Chemistry, biochemistry, carcinogenicity, and relevance to humans. *J. Toxicol. environ. Health*, **41**, 1–52
- Hoffmann, D.H., Djordjevic, M.V., Fan, J., Zang, E., Glynn, T. & Connolly, G.N. (1995) Five leading U.S. commercial brands of moist snuff in 1994: Assessment of carcinogenic N-nitrosamines. *J. natl Cancer Inst.*, **87**, 1862–1869
- Hoffmann, D., Hoffmann, I. & El-Bayoumy, K. (2001) The less harmful cigarette: A controversial issue. A tribute to Ernst L. Wynder. *Chem. Res. Toxicol.*, **14**, 767–790
- Hong, J.-Y., Ding, X., Smith, T.J., Coon, M.J. & Yang, C.S. (1992) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), a tobacco-specific carcinogen, by rabbit nasal microsomes and cytochrome P450s NMa and NMb. *Carcinogenesis*, **13**, 2141–2144
- Honkakoski, P. & Negishi, M. (1997) The structure, function, and regulation of cytochrome P450 2A enzymes. *Drug Metab. Rev.*, **29**, 977–996
- Hughes, J.R., Hecht, S.S., Carmella, S.G., Murphy, S.E. & Callas, P. (2004) Smoking behaviour and toxin exposure during six weeks use of a potential reduced exposure product: Omni. *Tob. Control*, **13**, 175–179
- Hukkanen, J., Pelkonen, O., Hakkola, J. & Raunio, H. (2002) Expression and regulation of xenobiotic-metabolizing cytochrome P450 (CYP) enzymes in human lung. *Crit. Rev. Toxicol.*, **32**, 391–411
- Hurt, R.D., Croghan, G.A., Wolter, T.D., Croghan, I.T., Offord, K.P., Williams, G.M., Djordjevic, M.V., Richie, J.P., Jr & Jeffrey, A.M. (2000) Does smoking reduction result in reduction of

- biomarkers associated with harm? A pilot study using a nicotine inhaler. *Nicotine Tob. Res.*, **2**, 327–336
- IARC (1985) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 37, *Tobacco Habits other than Smoking; Betel-quid and Areca-nut Chewing; and Some Related Nitrosamines*, Lyon
- IARC (1986) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 38, *Tobacco Smoking*, Lyon
- IARC (2004) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 83, *Tobacco Smoke and Involuntary Smoking*, Lyon
- Idris, A.M., Nair, J., Ohshima, H., Friesen, M., Brouet, I., Faustman, E.M. & Bartsch, H. (1991) Unusually high levels of carcinogenic tobacco-specific nitrosamines in Sudan snuff (toombak). *Carcinogenesis*, **12**, 1115–1118
- Idris, A.M., Nair, J., Friesen, M., Ohshima, H., Brouet, I., Faustman, E.M. & Bartsch, H. (1992) Carcinogenic tobacco-specific nitrosamines are present at unusually high levels in the saliva of oral snuff users in Sudan. *Carcinogenesis*, **13**, 1001–1005
- Idris, A.M., Ibrahim, S.O., Vasstrand, E.N., Johannessen, A.C., Lillehaug, J.R., Magnusson, B., Wallstrom, M., Hirsch, J.M. & Nilsen, R. (1998) The Swedish snus and the Sudanese toombak: Are they different? *Oral Oncol.*, **34**, 558–566
- Imai, T., Yasuhara, K., Tamufa, T., Takizawa, T., Ueda, M., Hirose, M. & Mitsumori, K. (2002) Inhibitory effects of cinnamaldehyde on 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung carcinogenesis in *rash2* mice. *Cancer Lett.*, **175**, 9–18
- Jalas, J.R. & Hecht, S.S. (2003) Synthesis of stereospecifically deuterated 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) diastereomers and metabolism by A/J mouse lung microsomes and cytochrome p450 2A5. *Chem. Res. Toxicol.*, **16**, 782–793
- Jalas, J.R., Ding, X. & Murphy, S.E. (2003a) Comparative metabolism of the tobacco-specific nitrosamines 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol by rat cytochrome P450 2A3 and human cytochrome P450 2A13. *Drug Metab. Dispos.*, **31**, 1199–1202
- Jalas, J.R., McIntee, E.J., Kenney, P.M.J., Upadhyaya, P., Peterson, L.A. & Hecht, S.S. (2003b) Stereospecific deuterium substitution attenuates the tumorigenicity and metabolism of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Chem. Res. Toxicol.*, **16**, 794–806
- Jalas, J.R., Hecht, S.S. & Murphy, S.E. (2005) Cytochrome P450 enzymes as catalysts of metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), a tobacco-specific carcinogen. *Chem. Res. Toxicol.*, **18**, 95–110
- Jansen, J.G., de Groot, A.J.L., van Teijlingen, C.M.M., Bates, A.D., Vrieling, H. & van Zeeland, A.A. (1996) Induction of *hprt* gene mutations in splenic T-lymphocytes from the rat exposed *in vivo* to DNA methylating agents is correlated with formation of *O*⁶-methylguanine in bone marrow and not in the spleen. *Carcinogenesis*, **17**, 2183–2191
- Jansson, C., Paccou, A. & Österdahl, B.-G. (2003) Analysis of tobacco-specific N-nitrosamines in snuff by ethyl acetate extraction and liquid chromatography–tandem mass spectrometry. *J. Chromatogr. A*, **1008**, 135–143
- Jin, Z., Gao, F., Flagg, T. & Deng, X. (2004) Tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone promotes functional cooperation of Bcl2 and c-Myc through phosphorylation in regulating cell survival and proliferation. *J. Biol. Chem.*, **279**, 40209–40219

- Jorquera, R., Castonguay, A. & Schuller, H.M. (1994) DNA single-strand breaks and toxicity induced by 4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone or N-nitrosodimethylamine in hamster and rat liver. *Carcinogenesis*, **15**, 389–394
- Joseph, A.M., Hecht, S.S., Murphy, S.E., Carmella, S.G., Le, C.T., Zhang, Y., Han, S. & Hatsukami, D.K. (2005) Relationships between cigarette consumption and biomarkers of tobacco toxin exposure. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 2963–2968
- Kato, S., Petruzzelli, S., Bowman, E.D., Turteltaub, K.W., Blomeke, B., Weston, A. & Shields, P.G. (1993) 7-Alkyldeoxyguanosine adduct detection by two-step HPLC and the ³²P-postlabeling assay. *Carcinogenesis*, **14**, 545–550
- Kato, S., Bowman, E.D., Harrington, A.M., Blomeke, B. & Shields, P.G. (1995) Human lung carcinogen–DNA adduct levels mediated by genetic polymorphisms in vivo. *J. natl Cancer Inst.*, **87**, 902–907
- Kawano, R., Takeshima, Y. & Inai, K. (1996) Effects of K-ras gene mutations in the development of lung lesions induced by 4-(N-methyl-N-nitrosamino)-1-(3-pyridyl)-1-butanone in A/J mice. *Jpn. J. Cancer Res.*, **87**, 44–50
- Keyler, D., Pentel, P.R., Kuehl, G., Collins, G. & Murphy, S.E. (2003) Effects of nicotine infusion on the metabolism of the tobacco carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in rats. *Cancer Lett.*, **202**, 1–9
- King, B., Borland, R. & Fowles, J. (2007) Mainstream smoke emissions of Australian and Canadian cigarettes. *Nicotine Tob. Res.*, **9**, 835–844
- Klein-Szanto, A.J.P., Iizasa, T., Momiki, S., Garcia-Palazzo, I., Caamano, J., Metcalf, R., Welsh, J. & Harris, C.C. (1992) A tobacco-specific N-nitrosamine or cigarette smoke condensate causes neoplastic transformation of xenotransplanted human bronchial epithelial cells. *Proc. natl Acad. Sci. USA*, **89**, 6693–6697
- Klus, H., Begutter, H., Scherer, G., Tricker, A.R. & Adlkofer, F. (1992) Tobacco-specific and volatile N-nitrosamines in environmental tobacco smoke of offices. *Indoor Environ.*, **1**, 348–350
- Knasmüller, S., Kassie, F., Zohrer, E., Kundi, M., McCoy, D.G. & Schulte-Hermann, R. (1994) Effects of ethanol treatment on DNA damage induced in *Escherichia coli* K-12 in various organs of mice by N-nitrosornicotine, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and N-nitrosopyrrolidine. *Carcinogenesis*, **15**, 263–270
- Kolar, C. & Lawson, T. (1997) Mutagenicity of carcinogenic nitrosamines when activated by hamster and human pancreatic duct epithelial cells. *Cancer Lett.*, **117**, 149–154
- Koppang, N., Rivenson, A., Reith, A., Dahle, H.K., Evensen, O. & Hoffmann, D. (1992) A study of tobacco carcinogenesis. XLVIII. Carcinogenicity of N'-nitrosornicotine in mink (*Mustela vison*). *Carcinogenesis*, **13**, 1957–1960
- Koppang, N., Rivenson, A., Dahle, H.K. & Hoffmann, D. (1997) A study of tobacco carcinogenesis. LIII. Carcinogenicity of N'-nitrosornicotine (NNN) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in mink (*Mustela vison*). *Cancer Lett.*, **111**, 167–171
- Krause, G., Garganta, F., Vrieling, H. & Scherer, G. (1999) Spontaneous and chemically induced point mutations in *HPRT* cDNA of the metabolically competent human lymphoblastoid cell line, MCL-5. *Mutat. Res.*, **431**, 417–428
- Kresty, L.A., Carmella, S.G., Borukhova, A., Akerkar, S.A., Gopalakrishnan, R., Harris, R.E., Stoner, G.D. & Hecht, S.S. (1996) Metabolites of a tobacco-specific nitrosamine, 4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone (NNK), in the urine of smokeless tobacco users:

- Relationship between urinary biomarkers and oral leukoplakia. *Cancer Epidemiol. Biomarkers Prev.*, **5**, 521–525
- Kumar, R. & Hemminki, K. (1996) Separation of 7-methyl- and 7-(2-hydroxyethyl)-guanine adducts in human DNA samples using a combination of TLC and HPLC. *Carcinogenesis*, **17**, 485–492
- Kumar, R., Siddiqi, M., Tricker, R. & Preussmann, R. (1991) Tobacco-specific *N*-nitrosamines in tobacco and mainstream smoke of Indian cigarettes. *Food chem. Toxicol.*, **29**, 405–407
- Kushida, H., Fujita, K., Suzuki, A., Yamada, M., Endo, T., Nohmi, T. & Kamataki, T. (2000a) Metabolic activation of *N*-alkylnitrosamines in genetically engineered *Salmonella typhimurium* expressing CYP2E1 or CYP2A6 together with human NADPH-cytochrome P450 reductase. *Carcinogenesis*, **21**, 1227–1232
- Kushida, H., Fujita, K.-I., Suzuki, A., Yamada, M., Nohmi, T. & Kamataki, T. (2000b) Development of a *Salmonella* tester strain sensitive to promutagenic *N*-nitrosamines: Expression of recombinant CYP2A6 and human NADPH-cytochrome P450 reductase in *S. typhimurium* YG7108. *Mutat. Res.*, **471**, 135–143
- Lackmann, G.M., Salzberger, U., Tollner, U., Chen, M., Carmella, S.G. & Hecht, S.S. (1999) Metabolites of a tobacco-specific carcinogen in urine from newborns. *J. natl Cancer Inst.*, **91**, 459–465
- Lacroix, D., Desrochers, M., Castonguay, A. & Anderson, A. (1993) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in human kidney epithelial cells transfected with rat CYP2B1 cDNA. *Carcinogenesis*, **14**, 1639–1642
- LaVoie, E.J., Prokopczyk, G., Rigotty, J., Czech, A., Rivenson, A. & Adams, J.D. (1987) Tumorigenic activity of the tobacco-specific nitrosamines 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), 4-(methylnitrosamino)-4-(3-pyridyl)-1-butanol (iso-NNAL) and *N'*-nitrosornicotine (NNN) on topical application to Sencar mice. *Cancer Lett.*, **37**, 277–283
- Lee, C.K., Fulp, C., Bombick, B.R. & Doolittle, D.J. (1996) Inhibition of mutagenicity of *N*-nitrosamines by tobacco smoke and its constituents. *Mutat. Res.*, **367**, 83–92
- Lemmonds, C.A., Hecht, S.S., Jensen, J.A., Murphy, S.E., Carmella, S.G., Zhang, Y. & Hatsukami, D.K. (2005) Smokeless tobacco topography and toxin exposure. *Nicotine Tob. Res.*, **7**, 469–474
- Lijinsky, W., Thomas, B.J. & Kovatch, R.M. (1991) Local and systemic carcinogenic effects of alkylating carcinogens in rats treated by intravesicular administration. *Jpn. J. Cancer Res.*, **82**, 980–986
- Lindberg, R.L.P. & Negishi, M. (1989) Alteration of mouse cytochrome P450_{coh} substrate specificity by mutation of a single amino-acid residue. *Nature*, **339**, 632–634
- Liu, L., Alaoui-Jamali, M.A., el Alami, N. & Castonguay, A. (1990) Metabolism and DNA single strand breaks induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and its analogues in primary culture of rat hepatocytes. *Cancer Res.*, **50**, 1810–1816
- Liu, L., Castonguay, A. & Gerson, S.L. (1992) Lack of correlation between DNA methylation and hepatocarcinogenesis in rats and hamsters treated with 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*, **13**, 2137–2140
- Liu, X.-K., Spratt, T.E., Murphy, S.E. & Peterson, L.A. (1996) Pyridyloxobutylation of guanine residues by 4-[(acetoxymethyl)nitrosamino]-1-(3-pyridyl)-1-butanone generates substrates of *O*⁶-alkylguanine-DNA alkyltransferase. *Chem. Res. Toxicol.*, **9**, 949–953

- Liu, L., Qin, X. & Gerson, S.L. (1999) Reduced lung tumorigenesis in human methylguanine DNA-methyltransferase transgenic mice achieved by expression of transgene within the target cell. *Carcinogenesis*, **20**, 279–284
- Loechler, E.L., Green, C.L. & Essigmann, J.M. (1984) *In vivo* mutagenesis by *O*⁶-methylguanine built into a unique site in a viral genome. *Proc. natl Acad. Sci. USA*, **81**, 6271–6275
- Lorenz, J., Schmassmann, H., Ohnhaus, E. & Oesch, F. (1979) Activities of polycyclic hydrocarbon activating and inactivating enzymes in human lungs of smokers, non-smokers, lung-cancer and non-cancer patients. *Arch. Toxicol.*, **Suppl.**, 483–489
- MacKown, C., Douglass, B., Djordjevic, M. & Bush, L. (1988) Tobacco-specific N-nitrosamines: Formation during processing of midrib and lamina fines. *J. agric. Food Chem.*, **36**, 1031–1035
- Majer, B.J., Mersch-Sundermann, V., Darroudi, F., Laky, B., de Wit, K. & Knasmuller, S. (2004) Genotoxic effects of dietary and lifestyle related carcinogens in human derived hepatoma (HepG2, Hep3B) cells. *Mutat. Res.*, **551**, 153–166
- Malson, J., Sims, K., Murty, R. & Pickworth, W. (2001) Comparison of the nicotine content of tobacco used in bidis and conventional cigarettes. *Tob. Control*, **10**, 181–183
- Maser, E. (1998) 11 β -Hydroxysteroid dehydrogenase responsible for carbonyl reduction of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in mouse lung microsomes. *Cancer Res.*, **58**, 2996–3003
- Maser, E. (2004) Significance of reductases in the detoxification of the tobacco-specific carcinogen NNK. *Trends pharmacol. Sci.*, **25**, 235–237
- Maser, E., Richter, E. & Friebertshauer, J. (1996) The identification of 11 beta-hydroxysteroid dehydrogenase as carbonyl reductase of the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Eur. J. Biochem.*, **238**, 484–489
- Maser, E., Stinner, B. & Atalla, A. (2000) Carbonyl reduction of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) by cytosolic enzymes in human liver and lung. *Cancer Lett.*, **148**, 135–144
- Matzinger, S.A., Gunning, W.T., You, M. & Castonguay, A. (1994) Ki-*ras* mutations in 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-initiated and butylated hydroxytoluene-promoted lung tumors in A/J mice. *Mol. Carcinog.*, **11**, 42–48
- Matzinger, S.A., Crist, K.A., Stoner, G.D., Anderson, M.W., Pereira, M.A., Steele, V.E., Kelloff, G.J., Lubet, R.A. & You, M. (1995) K-*ras* mutations in lung tumors from A/J and A/J \times TSG-*p53* F1 mice treated with 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and phenethyl isothiocyanate. *Carcinogenesis*, **16**, 2487–2492
- McCoy, G.D., Hecht, S.S., Katayama, S. & Wynder, E.L. (1981) Differential effect of chronic ethanol consumption on the carcinogenicity of *N*-nitrosopyrrolidine and *N'*-nitrososnicotine in male Syrian golden hamsters. *Cancer Res.*, **41**, 2849–2854
- McCoy, G., Katayama, S., Young, R., Wyatt, M. & Hecht, S. (1982) Influence of chronic ethanol consumption on the metabolism and carcinogenicity of tobacco-related nitrosamines. In: Bartsch, H., O'Neill, I.K., Castegnaro, M. & Okada, M., eds, *N-Nitroso Compounds: Occurrence and Biological Effects* (IARC Scientific Publications No. 41), Lyon, pp. 635–642
- McIntee, E.J. & Hecht, S.S. (2000) Metabolism of *N'*-nitrososnicotine enantiomers by cultured rat esophagus and *in vivo* in rats. *Chem. Res. Toxicol.*, **13**, 192–199
- McNeill, A., Bedi, R., Islam, S., Alkhatib, M.N. & West, R. (2007) Levels of toxins in oral tobacco products in the UK. *Tob. Control*, **15**, 64–67

- Meger, M., Richter, E., Zwickenpflug, W., Oehlmann, C., Hargaden, M.B., A-Rahim, Y.I. & Vesell, E.S. (1999) Metabolism and disposition of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in rhesus monkeys. *Drug Metab. Dispos.*, **27**, 471–478
- Meger, M., Meger-Kossien, I., Riedel, K. & Scherer, G. (2000) Biomonitoring of environmental tobacco smoke (ETS)-related exposure to 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Biomarkers*, **5**, 33–45
- Melikian, A., Djordjevic, M., Hosey, J., Zhang, J., Chen, S., Zang, E., Muscat, J. & Stellman, S. (2007a) Gender differences relative to smoking behavior and emissions of toxins from mainstream cigarette smoke. *Nicotine Tob. Res.*, **9**, 377–387
- Melikian, A.A., Djordjevic, M.D., Chen, S., Richie, J.P., Jr & Stellman, S.D. (2007b) Effect of delivered dosage of cigarette smoke toxins on the levels of urinary biomarkers of exposure. *Cancer Epidemiol. Biomarkers Prev.*, **16**, 1408–1415
- Mijal, R.S., Thomson, N.M., Fleischer, N.L., Pauly, G.T., Moschel, R.C., Kanugula, S., Fang, Q., Pegg, A.E. & Peterson, L.A. (2004) The repair of the tobacco specific nitrosamine derived adduct *O*⁶-[4-oxo-4-(3-pyridyl)butyl]guanine by *O*⁶-alkylguanine–DNA alkyltransferase variants. *Chem. Res. Toxicol.*, **17**, 424–434
- Miller, J.A. (1994) Research in chemical carcinogenesis with Elizabeth Miller — A trail of discovery with our associates. *Drug Metab. Rev.*, **26**, 1–36
- Mills, N.E., Fishman, C.L., Rom, W.N., Dubin, N. & Jacobson, D.R. (1995) Increased prevalence of *K-ras* oncogene mutations in lung adenocarcinoma. *Cancer Res.*, **55**, 1444–1447
- Milunsky, A., Carmella, S.G., Ye, M. & Hecht, S.S. (2000) A tobacco-specific carcinogen in the fetus. *Prenat. Diagn.*, **20**, 307–310
- Morgan, W., Reece, J.B., Risner, C., Bennett, C., Midgett, C., Johnson, K. & Burton, H. (2004) A collaborative study for the determination of tobacco specific nitrosamines in tobacco. *Beitr. Tabakforsch. int./Control Tob. Res.*, **21**, 192–203
- Morin, A., Porter, A., RataVICIUS, A. & Joly, J. (2004) Evolution of tobacco-specific nitrosamines and microbial populations during flue-curing of tobacco under direct and indirect heating. *Beitr. Tabakforsch. int.*, **21**, 40–46
- Moriya, M. (1993) Single-stranded shuttle phagemid for mutagenesis studies in mammalian cells: 8-Oxoguanine in DNA induces targeted G.C→T.A transversions in simian kidney cells. *Proc. natl Acad. Sci. USA*, **90**, 1122–1126
- Moriya, M., Zhang, W., Johnson, F. & Grollman, A.P. (1994) Mutagenic potency of exocyclic DNA adducts: Marked differences between *Escherichia coli* and simian kidney cells. *Proc. natl Acad. Sci. USA*, **91**, 11899–11903
- Morse, M.A., LaGreca, S.D., Amin, S.G. & Chung, F.-L. (1990a) Effects of indole-3-carbinol on lung tumorigenesis and DNA methylation induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and on the metabolism and disposition of NNK in A/J mice. *Cancer Res.*, **50**, 2613–2617
- Morse, M.A., Eklind, K.I., Toussaint, M., Amin, S.G. & Chung, F.-L. (1990b) Characterization of a glucuronide metabolite of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and its dose-dependent excretion in the urine of mice and rats. *Carcinogenesis*, **11**, 1819–1823
- Morse, M.A., Eklind, K.I., Hecht, S.S., Jordan, K.G., Choi, C.I., Desai, D.H., Amin, S.G. & Chung, F.-L. (1991) Structure–activity relationships for inhibition of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone lung tumorigenesis by arylalkyl isothiocyanates in A/J mice. *Cancer Res.*, **51**, 1846–1850

- Murphy, S.E. & Coletta, K.A. (1993) Two types of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone hemoglobin adducts, from metabolites which migrate into or are formed in red blood cells. *Cancer Res.*, **53**, 777–783
- Murphy, S.E. & Heiblum, R. (1990) Effect of nicotine and tobacco-specific nitrosamines on the metabolism of *N'*-nitrosornicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by rat oral tissue. *Carcinogenesis*, **11**, 1663–1666
- Murphy, S.E. & Spina, D.A. (1994) Evidence for a high-affinity enzyme in rat esophageal microsomes which α -hydroxylates *N'*-nitrosornicotine. *Carcinogenesis*, **15**, 2709–2713
- Murphy, S.E., Heiblum, R. & Trushin, N. (1990a) Comparative metabolism of *N'*-nitrosornicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by cultured F344 rat oral tissue and esophagus. *Cancer Res.*, **50**, 4685–4691
- Murphy, S.E., Palomino, A., Hecht, S.S. & Hoffmann, D. (1990b) Dose–response study of DNA and hemoglobin adduct formation by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in F344 rats. *Cancer Res.*, **50**, 5446–5452
- Murphy, S.E., Carmella, S.G., Idris, A.M. & Hoffmann, D. (1994) Uptake and metabolism of carcinogenic levels of tobacco-specific nitrosamines by Sudanese snuff dippers. *Cancer Epidemiol. Biomarkers Prev.*, **3**, 423–428
- Murphy, S.E., Isaac, I.S., Ding, X. & McIntee, E.J. (2000) Specificity of cytochrome P450 2A3-catalyzed α -hydroxylation of *N'*-nitrosornicotine enantiomers. *Drug Metab. Dispos.*, **28**, 1263–1266
- Murrah, V.A., Gilchrist, E.P. & Moyer, M.P. (1993) Morphologic and growth effects of tobacco-associated chemical carcinogens and smokeless tobacco extracts on human oral epithelial cells in culture. *Oral Surg. oral Med. oral Pathol.*, **75**, 323–332
- Muscat, J.E., Djordjevic, M.V., Colosimo, S., Stellman, S.D. & Richie, J.P., Jr (2005) Racial differences in exposure and glucuronidation of the tobacco-specific carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Cancer*, **103**, 1420–1426
- Mustonen, R., Schoket, B. & Hemminki, K. (1993) Smoking-related DNA adducts: ³²P-Postlabeling analysis of 7-methylguanine in human bronchial and lymphocyte DNA. *Carcinogenesis*, **14**, 151–154
- Nair, J., Ohshima, H., Friesen, M., Croisy, A., Bhide, S.V. & Bartsch, H. (1985) Tobacco-specific and betel nut-specific N-nitroso compounds: Occurrence in saliva and urine of betel quid chewers and formation *in vitro* by nitrosation of betel quid. *Carcinogenesis*, **6**, 295–303
- Nair, U.J., Pakhale, S.S., Spiegelhalter, B., Preussmann, R. & Bhide, S.V. (1987) Carcinogenic and cocarcinogenic constituents of Masheri, a pyrolysed tobacco product. *Indian J. Biochem. Biophys.*, **24**, 257–259
- Nair, J., Pakhale, S.S. & Bhide, S.V. (1989) Carcinogenic tobacco-specific nitrosamines in Indian tobacco products. *Food chem. Toxicol.*, **27**, 751–753
- Nair, U.J., Ammigan, N., Nagabhushan, M., Amonkar, A.J. & Bhide, S.V. (1991) Effect of vitamin A status of rats on metabolizing enzymes after exposure to tobacco extract or *N'*-nitrosornicotine. In: O'Neill, I.K., Chen, J. & Bartsch, H., eds, *Relevance to Human Cancer of N-Nitroso Compounds, Tobacco Smoke and Mycotoxins* (IARC Scientific Publications No. 105), Lyon, pp. 525–528
- Nesnow, S., Ross, J.A., Stoner, G.D. & Mass, M.J. (1995) Mechanistic linkage between DNA adducts, mutations in oncogenes and tumorigenesis of carcinogenic environmental polycyclic aromatic hydrocarbons in strain A/J mice. *Toxicology*, **105**, 403–413

- Nunes, M.G., Desai, D., Koehl, W., Spratt, T.E., Guengerich, F.P. & Amin, S. (1998) Inhibition of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) metabolism in human hepatic microsomes by ipomeanol analogs — An exploratory study. *Cancer Lett.*, **129**, 131–138
- Ohshima, H., Nair, J., Bourgade, M.-C., Friesen, M., Garren, L. & Bartsch, H. (1985) Identification and occurrence of two new *N*-nitrosamino acids in tobacco products: 3-(*N*-Nitroso-*N*-methylamino) propionic acid and 4-(*N*-nitroso-*N*-methylamino) butyric acid. *Cancer Lett.*, **26**, 153–162
- Oreffo, V.I.C., Lin, H.-W., Padmanabhan, R. & Witschi, H. (1993) *K-ras* and *p53* point mutations in 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced hamster lung tumors. *Carcinogenesis*, **14**, 451–455
- Osterdahl, B.-G. & Slorach, S. (1988) Tobacco-specific *N*-nitrosamines in the saliva of habitual male snuff dippers. *Food Addit. Contam.*, **5**, 581–586
- Österdahl, B.-G., Jansson, C. & Paccou, A. (2004) Decreased levels of tobacco-specific *N*-nitrosamines in moist snuff on the Swedish Market. *J. agric. Food Chem.*, **52**, 5085–5088
- Padma, P.R., Lalitha, V.S., Amonkar, A.J. & Bhide, S.V. (1989a) Carcinogenicity studies on the two tobacco-specific *N*-nitrosamines, *N'*-nitrososornicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*, **10**, 1997–2002
- Padma, P.R., Amonkar, A.J. & Bhide, S.V. (1989b) Mutagenic and cytogenetic studies of *N'*-nitrososornicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Lett.*, **46**, 173–180
- Pakhale, S.S. & Maru, G.B. (1998) Distribution of major and minor alkaloids in tobacco, mainstream and sidestream smoke of popular Indian smoking products. *Food chem. Toxicol.*, **36**, 1131–1138
- Papageorge, M.B., Cataldo, E. & Jahngen, E.G.E. (1996) The effect of *N*-nitrososornicotine on the buccal mucosa of Syrian hamsters. *J. oral maxillofac. Surg.*, **54**, 187–190
- Patten, C.J., Smith, T.J., Murphy, S.E., Wang, M.-H., Lee, J., Tynes, R.E., Koch, P. & Yang, C.S. (1996) Kinetic analysis of the activation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by heterologously expressed human P450 enzymes and the effect of P450-specific chemical inhibitors on this activation in human liver microsomes. *Arch. Biochem. Biophys.*, **333**, 127–138
- Patten, C.J., Smith, T.J., Friesen, M.J., Tynes, R.E., Yang, C.S. & Murphy, S.E. (1997) Evidence for cytochrome P450 2A6 and 3A4 as major catalysts for *N'*-nitrososornicotine α -hydroxylation by human liver microsomes. *Carcinogenesis*, **18**, 1623–1630
- Patten, C.J., Peterson, L.A. & Murphy, S.E. (1998) Evidence for metabolic activation of *N'*-nitrososornicotine and *N*-nitrosobenzylmethylamine by a rat nasal coumarin hydroxylase. *Drug Metab. Dispos.*, **26**, 177–180
- Pauly, G.T., Peterson, L.A. & Moschel, R.C. (2002) Mutagenesis by *O*⁶-[4-oxo-4-(3-pyridyl)-butyl]guanine in *Escherichia coli* and human cells. *Chem. Res. Toxicol.*, **15**, 165–169
- Peele, D.M., Riddick, M.G. & Edwards, M.E. (2001) Formation of tobacco-specific nitrosamines in flue-cured tobacco. *Recent Adv. Tob. Sci.*, **27**, 3–12
- Peterson, L.A. & Hecht, S.S. (1991) *O*⁶-Methylguanine is a critical determinant of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone tumorigenesis in A/J mouse lung. *Cancer Res.*, **51**, 5557–5564
- Peterson, L.A., Carmella, S.G. & Hecht, S.S. (1990) Investigations of metabolic precursors to hemoglobin and DNA adducts of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*, **11**, 1329–1333
- Peterson, L.A., Mathew, R. & Hecht, S.S. (1991a) Quantitation of microsomal alpha-hydroxylation of the tobacco-specific nitrosamine, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res.*, **51**, 5495–5500

- Peterson, L.A., Mathew, R., Murphy, S.E., Trushin, N. & Hecht, S.S. (1991b) *In vivo* and *in vitro* persistence of pyridyloxobutyl DNA adducts from 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*, **12**, 2069–2072
- Peterson, L.A., Liu, X.-K. & Hecht, S.S. (1993) Pyridyloxobutyl DNA adducts inhibit the repair of O⁶-methylguanine. *Cancer Res.*, **53**, 2780–2785
- Peterson, L.A., Ng, D.K., Stearns, R.A. & Hecht, S.S. (1994) Formation of NADP(H) analogs of tobacco-specific nitrosamines in rat liver and pancreatic microsomes. *Chem. Res. Toxicol.*, **7**, 599–608
- Peterson, L.A., Thomson, N.M., Crankshaw, D.L., Donaldson, E.E. & Kenney, P.J. (2001) Interactions between methylating and pyridyloxobutylating agents in A/J mouse lungs: Implications for 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung tumorigenesis. *Cancer Res.*, **61**, 5757–5763
- Petruzzelli, S., Tavanti, L.M., Celi, A. & Giuntini, C. (1996) Detection of *N*⁷-methyldeoxyguanosine adducts in human pulmonary alveolar cells. *Am. J. respir. Cell mol. Biol.*, **15**, 216–223
- Pfeifer, G.P., Denissenko, M.F., Olivier, M., Tretyakova, N., Hecht, S.S. & Hainaut, P. (2002) Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers. *Oncogene*, **21**, 7435–7451
- Pohlmann, C., Koops, F., Berg, J., Holz, O., Ehlert, U. & Rüdiger, H.W. (1992) Determinants of a genotoxic effect of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in human diploid fibroblasts. *Clin. Invest.*, **70**, 295–298
- Pool-Zobel, B.L., Klein, R.G., Liegibel, U.M., Kuchenmeister, F., Weber, S. & Schmezer, P. (1992) Systemic genotoxic effects of tobacco-related nitrosamines following oral and inhalational administration to Sprague-Dawley rats. *Clin. Invest.*, **70**, 299–306
- von Pressentin, M.M., Kosinska, W. & Guttenplan, J.B. (1999) Mutagenesis induced by oral carcinogens in *lacZ* mouse (MutaTMMouse) tongue and other oral tissues. *Carcinogenesis*, **20**, 2167–2170
- Prokopczyk, B., Rivenson, A. & Hoffmann, D. (1991) A study of betel quid carcinogenesis. IX. Comparative carcinogenicity of 3-(methylnitrosamino)propionitrile and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone upon local application to mouse skin and rat oral mucosa. *Cancer Lett.*, **60**, 153–157
- Prokopczyk, B., Hoffmann, D., Cox, J.E., Djordevic, M.V. & Brunnemann, K.D. (1992a) Supercritical fluid extraction in the determination of tobacco-specific *N*-nitrosamines in smokeless tobacco. *Chem. Res. Toxicol.*, **5**, 336–340
- Prokopczyk, B., Wu, M., Cox, J.E. & Hoffmann, D. (1992b) Bioavailability of tobacco-specific *N*-nitrosamines to the snuff dipper. *Carcinogenesis*, **13**, 863–866
- Prokopczyk, B., Wu, M., Cox, J.E., Amin, S., Desai, D., Idris, A.M. & Hoffmann, D. (1995) Improved methodology for the quantitative assessment of tobacco-specific *N*-nitrosamines in tobacco by supercritical fluid extraction. *J. agric. Food Chem.*, **43**, 916–922
- Prokopczyk, B., Cox, J.E., Hoffmann, D. & Waggoner, S.E. (1997) Identification of tobacco-specific carcinogen in the cervical mucus of smokers and nonsmokers. *J. natl Cancer Inst.*, **89**, 868–873
- Prokopczyk, B., Trushin, N., Leszczynska, J., Waggoner, S.E. & El Bayoumy, K. (2001) Human cervical tissue metabolizes the tobacco-specific nitrosamine, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, via α -hydroxylation and carbonyl reduction pathways. *Carcinogenesis*, **22**, 107–114

- Prokopczyk, B., Hoffmann, D., Bologna, M., Cunningham, A.J., Trushin, N., Akerkar, S., Boyiri, T., Amin, S., Desai, D., Colosimo, S., Pittman, B., Leder, G., Ramadani, M., Henne-Bruns, D., Beger, H.G. & El Bayoumy, K. (2002) Identification of tobacco-derived compounds in human pancreatic juice. *Chem. Res. Toxicol.*, **15**, 677–685
- Proulx, L.-I., Castonguay, A. & Bissonnette, E.Y. (2004) Cytokine production by alveolar macrophages is down regulated by the α -methylhydroxylation pathway of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Carcinogenesis*, **25**, 997–1003
- Pulling, L.C., Klinge, D.M. & Belinsky, S.A. (2001) *p16^{INK4a}* and *beta-Catenin* alterations in rat liver tumors induced by NNK. *Carcinogenesis*, **22**, 461–466
- Pulling, L.C., Vuilleminot, B.R., Hutt, J.A., Devereux, T.R. & Belinsky, S.A. (2004) Aberrant promoter hypermethylation of the *death-associated protein kinase* gene is early and frequent in murine lung tumors induced by cigarette smoke and tobacco carcinogens. *Cancer Res.*, **64**, 3844–3848
- Ren, Q., Murphy, S.E., Dannenberg, A.J., Park, J.Y., Tephly, T.R. & Lazarus, P. (1999) Glucuronidation of the lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) by rat UDP-glucuronosyltransferase 2B1. *Drug Metab. Dispos.*, **27**, 1010–1016
- Ren, Q., Murphy, S.E., Zheng, Z. & Lazarus, P. (2000) *O*-Glucuronidation of the lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) by human UDP-glucuronosyltransferases 2B7 and 1A9. *Drug Metab. Dispos.*, **28**, 1352–1360
- Richter, E. & Tricker, A.R. (2002) Effect of nicotine, cotinine and phenethyl isothiocyanate on 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) metabolism in the Syrian golden hamster. *Toxicology*, **179**, 95–103
- Richter, E., Friesenegger, S., Engl, J. & Tricker, A.R. (2000) Use of precision-cut tissue slices in organ culture to study metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) by hamster lung, liver and kidney. *Toxicology*, **144**, 83–91
- Rickert, W.S. & Kaiserman, M.J. (1999) Application of proposed Canadian test methods to the analysis of cigarette filler, fine cut tobacco, and tobacco smoke (Abstract No. 16). In: *Proceedings of the 53rd Tobacco Science Research Conference, Montreal, Canada, September 12–15, 1999*
- Risner, C., Reece J.B. & Morgan, W.T. (2001) The determination of tobacco-specific nitrosamines in tobacco: A collaborative investigation of current methodology. *Recent Adv. Tob. Sci.*, **27**, 47–74
- Rivenson, A., Hoffmann, D., Prokopczyk, B., Amin, S. & Hecht, S.S. (1988) Induction of lung and exocrine pancreas tumors in F344 rats by tobacco-specific and areca-derived N-nitrosamines. *Cancer Res.*, **48**, 6912–6917
- Rivenson, A., Djordjevic, M.V., Amin, S. & Hoffmann, R. (1989) A study of tobacco carcinogenesis. XLIV. Bioassay in A/J mice of some N-nitrosamines. *Cancer Lett.*, **47**, 111–114
- Rodenhuis, S. & Slebos, R.J.C. (1992) Clinical significance of *ras* oncogene activation in human lung cancer. *Cancer Res.*, **52**, 2665s–2669s
- Ronai, Z.A., Gradia, S., Peterson, L.A. & Hecht, S.S. (1993) G to A transitions and G to T transversions in codon 12 of the *Ki-ras* oncogene isolated from mouse lung tumors induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and related DNA methylating and pyridyloxobutylating agents. *Carcinogenesis*, **14**, 2419–2422
- Rossignol, G., Alaoui-Jamali, M.A., Castonguay, A. & Schuller, H.M. (1989) Metabolism and DNA damage induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in fetal tissues of the Syrian golden hamster. *Cancer Res.*, **49**, 5671–5676

- Scherer, G., Engl., J., Urban, M., Gilch, G., Janket, D. & Riedel, K. (2007) Relationship between machine-derived smoke yields and biomarkers in cigarette smokers in Germany. *Regul. Toxicol. Pharmacol.*, **47**, 171–183
- Schneider, T., Frei, E. & Wiessler, M. (1999) The profile of urinary metabolites of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in rats is determined by its pulmonary metabolism. *Chem.-biol. Interact.*, **118**, 247–266
- Schrader, E., Hirsch-Ernst, K.I., Richter, E. & Foth, H. (1998) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in isolated rat lung and liver. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **357**, 336–343
- Schrader, E., Hirsch-Ernst, K.I., Scholz, E., Kahl, G.F. & Foth, H. (2000) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in primary cultures of rat alveolar type II cells. *Drug Metab. Dispos.*, **28**, 180–185
- Schuller, H.M. (2002) Mechanisms of smoking-related lung and pancreatic adenocarcinoma development. *Nat. Rev. Cancer*, **2**, 455–463
- Schuller, H.M., Witschi, H.P., Nylén, E., Joshi, P.A., Correa, E. & Becker, K.L. (1990) Pathobiology of lung tumors induced in hamsters by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and the modulating effect of hyperoxia. *Cancer Res.*, **50**, 1960–1965
- Schüller, H.M., Jorquera, R., Riechert, A. & Castonguay, A. (1993) Transplacental induction of pancreas tumors in hamsters by ethanol and the tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res.*, **53**, 2498–2501
- Schüller, H.M., Jorquera, R., Lu, X., Riechert, A. & Castonguay, A. (1994) Transplacental carcinogenicity of low doses of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone administered subcutaneously or intratracheally to hamsters. *J. Cancer Res. clin. Oncol.*, **120**, 200–203
- Schuller, H.M., Zhang, L., Weddle, D.L., Castonguay, A., Walker, K. & Miller, M.S. (2002) The cyclooxygenase inhibitor ibuprofen and the FLAP inhibitor MK886 inhibit pancreatic carcinogenesis induced in hamsters by transplacental exposure to ethanol and the tobacco carcinogen NNK. *J. Cancer Res. clin. Oncol.*, **128**, 525–532
- Schuller, H.M., Plummer, H.K., III & Jull, B.A. (2003) Receptor-mediated effects of nicotine and its nitrosated derivative NNK on pulmonary neuroendocrine cells. *Anat. Rec.*, **270A**, 51–58
- Schulze, J., Richter, E., Binder, U. & Zwickenpflug, W. (1992) Biliary excretion of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in the rat. *Carcinogenesis*, **13**, 1961–1965
- Sellers, E.M., Ramamoorthy, Y., Zeman, M.V., Djordjevic, M.V. & Tyndale, R.F. (2003) The effect of methoxsalen on nicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) metabolism in vivo. *Nicotine Tob. Res.*, **5**, 891–899
- Seo, K.-Y., Jelinsky, S.A. & Loechler, E.L. (2000) Factors that influence the mutagenic patterns of DNA adducts from chemical carcinogens. *Mutat. Res.*, **463**, 215–246
- Sharma, A.K., Prokopczyk, B. & Hoffmann, D. (1991) Supercritical fluid extraction of moist snuff. *J. agric. Food Chem.*, **39**, 508–510
- Shields, P.G., Povey, A.C., Wilson, V.L., Weston, A. & Harris, C.C. (1990) Combined high-performance liquid chromatography/³²P-postlabeling assay of *N*⁷-methyldeoxyguanosine. *Cancer Res.*, **50**, 6580–6584
- Shimada, T., Yun, C.-H., Yamazaki, H., Gautier, J.C., Beaune, P.H. & Guengerich, F.P. (1992) Characterization of human lung microsomal cytochrome P-450 1A1 and its role in the oxidation of chemical carcinogens. *Mol. Pharmacol.*, **41**, 856–864

- Shimada, T., Yamazaki, H., Mimura, M., Inui, Y. & Guengerich, F.P. (1994) Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: Studies with liver microsomes of 30 Japanese and 30 Caucasians. *J. Pharmacol. exp. Ther.*, **270**, 414–423
- Shimada, T., Yamazaki, H., Mimura, M., Wakamiya, N., Ueng, Y.-F., Guengerich, F.P. & Inui, Y. (1996) Characterization of microsomal cytochrome P450 enzymes involved in the oxidation of xenobiotic chemicals in human fetal liver and adult lungs. *Drug Metab. Dispos.*, **24**, 515–522
- Singer, B. & Essigmann, J.M. (1991) Site-specific mutagenesis: Retrospective and prospective. *Carcinogenesis*, **12**, 949–955
- Singer, B. & Grunberger, D. (1983) *Molecular Biology of Mutagens and Carcinogens*, New York, Plenum Press
- Sipowicz, M.A., Amin, S., Desai, D., Kasprzak, K.S. & Anderson, L.M. (1997) Oxidative DNA damage in tissues of pregnant female mice and fetuses caused by the tobacco-specific nitrosamine, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Cancer Lett.*, **117**, 87–91
- Smith, T.J., Guo, Z.Y., Thomas, P.E., Chung, F.-L., Morse, M.A., Elkind, K. & Yang, C.S. (1990) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in mouse lung microsomes and its inhibition by isothiocyanates. *Cancer Res.*, **50**, 6817–6822
- Smith, T.J., Guo, Z., Gonzalez, F.J., Guengerich, F.P., Stoner, G.D. & Yang, C.S. (1992a) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in human lung and liver microsomes and cytochromes P-450 expressed in hepatoma cells. *Cancer Res.*, **52**, 1757–1763
- Smith, T.J., Guo, Z., Hong, J.-Y., Ning, S.M., Thomas, P.E. & Yang, C.S. (1992b) Kinetics and enzyme involvement in the metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in microsomes of rat lung and nasal mucosa. *Carcinogenesis*, **13**, 1409–1414
- Smith, T.J., Guo, Z., Li, C., Ning, S.M., Thomas, P.E. & Yang, C.S. (1993) Mechanisms of inhibition of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone bioactivation in mouse by dietary phenethyl isothiocyanate. *Cancer Res.*, **53**, 3276–3282
- Smith, T.J., Stoner, G.D. & Yang, C.S. (1995) Activation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in human lung microsomes by cytochromes P450, lipoxygenase, and hydroperoxides. *Cancer Res.*, **55**, 5566–5573
- Smith, T.J., Guo, Z., Guengerich, F.P. & Yang, C.S. (1996) Metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) by human cytochrome P450 1A2 and its inhibition by phenethyl isothiocyanate. *Carcinogenesis*, **17**, 809–813
- Smith, T.J., Liao, A.M., Liu, Y., Jones, A.B., Anderson, L.M. & Yang, C.S. (1997) Enzymes involved in the bioactivation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in patas monkey lung and liver microsomes. *Carcinogenesis*, **18**, 1577–1584
- Smith, T.J., Liao, A., Wang, L.-D., Yang, G.-Y., Starcic, S., Philbert, M.A. & Yang, C.S. (1998) Characterization of xenobiotic-metabolizing enzymes and nitrosamine metabolism in the human esophagus. *Carcinogenesis*, **19**, 667–672
- Smith, G.B., Bend, J.R., Bedard, L.L., Reid, K.R., Petsikas, D. & Massey, T.E. (2003) Biotransformation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in peripheral human lung microsomes. *Drug Metab. Dispos.*, **31**, 1134–1141
- Soucek, P. & Gut, I. (1992) Cytochromes P-450 in rats: Structures, functions, properties and relevant human forms. *Xenobiotica*, **22**, 83–103
- Spiegelhalder, B. & Fischer, S. (1991) Formation of tobacco-specific nitrosamines. *Crit. Rev. Toxicol.*, **21**, 241

- Spiegelhalder, B., Kubacki, S. & Fischer, S. (1989) A method for the determination of tobacco-specific nitrosamines (TSNA), nitrate and nitrite in tobacco leaves and processed tobacco. *Beitr. Tabakforsch. int.*, **14**, 135–144
- Spivack, S.D., Hurteau, G.J., Reilly, A.A., Aldous, K.M., Ding, X. & Kaminsky, L.S. (2001) CYP1B1 expression in human lung. *Drug Metab. Dispos.*, **29**, 916–922
- Spratt, T.E., Trushin, N., Lin, D. & Hecht, S.S. (1989) Analysis for N²-(pyridyloxobutyl)deoxyguanosine adducts in DNA of tissues exposed to tritium-labeled 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and N'-nitrosornicotine. *Chem. Res. Toxicol.*, **2**, 169–173
- Squier, C.A. (1986) Penetration of nicotine and nitrosornicotine across porcine oral mucosa. *J. appl. Toxicol.*, **6**, 123–128
- Squier, C.A., Kremer, M.J. & Wertz, P.W. (2003) Effect of ethanol on lipid metabolism and epithelial permeability barrier of skin and oral mucosa in the rat. *J. oral Pathol. Med.*, **32**, 595–599
- StAAF, M., Back, S., Wiernik, A., Wahlberg, I., Long, R. & Young, J. (2005) Formation of tobacco-specific nitrosamines (TSNA) during air-curing: conditions and control. *Beitr. Tabakforsch. int.*, **21**, 321–330
- Staretz, M.E., Murphy, S.E., Patten, C.J., Nunes, M.G., Koehl, W., Amin, S., Koenig, L.A., Guengerich, F.P. & Hecht, S.S. (1997a) Comparative metabolism of the tobacco-related carcinogens benzo[a]pyrene, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol, and N'-nitrosornicotine in human hepatic microsomes. *Drug Metab. Dispos.*, **25**, 154–162
- Staretz, M.E., Koenig, L.A. & Hecht, S.S. (1997b) Effects of long term dietary phenethyl isothiocyanate on the microsomal metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in F344 rats. *Carcinogenesis*, **18**, 1715–1722
- Staretz, M.E., Foiles, P.G., Miglietta, L.M. & Hecht, S.S. (1997c) Evidence for an important role of DNA pyridyloxobutylation in rat lung carcinogenesis by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone: Effects of dose and phenethyl isothiocyanate. *Cancer Res.*, **57**, 259–266
- Stepanov, I. & Hecht, S.S. (2005) Tobacco-specific nitrosamines and their pyridine-N-glucuronides in the urine of smokers and smokeless tobacco users. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 885–891
- Stepanov, I., Carmella, S.G., Hecht, S.S. & Duca, G. (2002) Analysis of tobacco-specific nitrosamines in Moldovan cigarette tobacco. *J. agric. Food Chem.*, **50**, 2793–2797
- Stepanov, I., Hecht, S.S., Ramakrishnan, S. & Gupta, P.C. (2005) Tobacco-specific nitrosamines in smokeless tobacco products marketed in India. *Int. J. Cancer*, **116**, 16–19
- Stepanov, I., Jensen, J., Hatsukami, D. & Hecht, S. (2006a) Tobacco-specific nitrosamines in new tobacco products. *Nicotine Tob. Res.*, **8**, 309–313
- Stepanov, I., Hecht, S.S., Duca, G. & Mardari, I. (2006b) Uptake of the tobacco-specific lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by Moldovan children. *Cancer Epidemiol. Biomarkers Prev.*, **15**, 7–11
- Stepanov, I., Feuer, R., Jensen, J., Hatsukami, D. & Hecht, S.S. (2006c) Mass spectrometric quantitation of nicotine, cotinine, and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in human toenails. *Cancer Epidemiol. Biomarkers Prev.*, **15**, 2378–2383
- Stich, H.F., Parida, B.B. & Brunnemann, K.D. (1992) Localized formation of micronuclei in the oral mucosa and tobacco-specific nitrosamines in the saliva of 'reverse smokers', khaini-tobacco chewers and gudakhu users. *Int. J. Cancer*, **50**, 172–176

- Su, T., Sheng, J.J., Lipinkas, T.W. & Ding, X. (1996) Expression of CYP2A genes in rodent and human nasal mucosa. *Drug Metab. Dispos.*, **24**, 884–890
- Su, T., Bao, Z., Zhang, Q.Y., Smith, T.J., Hong, J.-Y. & Ding, X. (2000) Human cytochrome P450 CYP2A13: Predominant expression in the respiratory tract and its high efficiency metabolic activation of a tobacco-specific carcinogen, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Cancer Res.*, **60**, 5074–5079
- Swedmark, S., Romert, L., Beije, B., Diaz-Pohl, C., Backbro, A. & Jenssen, D. (1994) Comparison of co-cultivation of V79 cells with rat hepatocytes and rat H4IIE hepatoma cells for studying nitrosamine-induced *hprt* gene mutations. *Mutagenesis*, **9**, 281–287
- Szyfter, K., Hemminki, K., Szyfter, W., Szmaja, Z., Banaszewski, J. & Pabiszczak, M. (1996) Tobacco smoke-associated N7-alkylguanine in DNA of larynx tissue and leucocytes. *Carcinogenesis*, **17**, 501–506
- Taioli, E., Garbers, S., Bradlow, H.L., Carmella, S.G., Akerkar, S. & Hecht, S.S. (1997) Effects of indole-3-carbinol on the metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in smokers. *Cancer Epidemiol. Biomarkers Prev.*, **6**, 517–522
- Therriault, M.-J., Proulx, L.-I., Castonguay, A. & Bissonnette, E.Y. (2003) Immunomodulatory effects of the tobacco-specific carcinogen, NNK, on alveolar macrophages. *Clin. exp. Immunol.*, **132**, 232–238
- Thomson, N.M., Kenney, P.M. & Peterson, L.A. (2003) The pyridyloxobutyl DNA adduct, O⁶-[4-oxo-4-(3-pyridyl)butyl]guanine, is detected in tissues from 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-treated A/J mice. *Chem. Res. Toxicol.*, **16**, 1–6
- Tiano, H.F., Wang, R.L., Hosokawa, M., Crespi, C., Tindall, K.R. & Langenbach, R. (1994) Human CYP2A6 activation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK): Mutational specificity in the *gpt* gene of AS52 cells. *Carcinogenesis*, **15**, 2859–2866
- Tjälve, H. (1991) The tissue distribution and the tissue specificity of bioactivation of some tobacco-specific and some other N-nitrosamines. *Crit. Rev. Toxicol.*, **21**, 265–294
- Tjälve, H. & Castonguay, A. (1983) The *in vivo* tissue disposition and *in vitro* target-tissue metabolism of the tobacco-specific carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in Syrian golden hamsters. *Carcinogenesis*, **4**, 1259–1265
- Tjälve, H., Castonguay, A. & Rivenson, A. (1985) Microautoradiographic localization of bound metabolites in the nasal cavities of F344 rats treated with the tobacco-specific carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *J. natl Cancer Inst.*, **74**, 185–189
- Toronto Research Chemicals (2005) *MSDS 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanone [NNK], N'-Nitrosornicotine [NNN], N-Nitrosoanatabine [NAT], N-Nitrosoanabasine [NAB]*, North York, Ontario [available at: <http://www.trc-canada.com>, accessed 02 Feb 2007]
- Toronto Research Chemicals (2006) *Certificates of Analysis*, North York, Ontario [available at: <http://www.trc-canada.com>, accessed 02 Feb 2007]
- Tricker, A.R. & Preussmann, R. (1988) The occurrence of N-nitro compounds in zarda tobacco. *Cancer Lett.*, **42**, 113–118
- Tricker, A.R. & Preussmann, R. (1989) The occurrence of N-nitroso compounds in kiwam tobacco. *Cancer Lett.*, **49**, 221–224
- Tricker, A.R. & Preussmann, R. (1991) Occurrence of and exposure to N-nitroso compounds in tobacco. In: O'Neill, I.K., Chen, J. & Bartsch, H., eds, *Relevance to Human Cancer of N-Nitroso Compounds, Tobacco Smoke and Mycotoxins* (IARC Scientific Publications No. 105), Lyon, 493–495

- Tricker, A.R., Haubner, R., Spiegelhalter, B. & Preussmann, R. (1988) The occurrence of tobacco-specific nitrosamines in oral tobacco products and their potential formation under simulated gastric conditions. *Food chem. Toxicol.*, **26**, 861–865
- Tricker, A.R., Ditrich, C. & Preussmann, R. (1991) N-Nitroso compounds in cigarette tobacco and their occurrence in mainstream tobacco smoke. *Carcinogenesis*, **12**, 257–261
- Trushin, N. & Hecht, S.S. (1999) Stereoselective metabolism of nicotine and tobacco-specific N-nitrosamines to 4-hydroxy-4-(3-pyridyl)butanoic acid in rats. *Chem. Res. Toxicol.*, **12**, 164–171
- Trushin, N., Rivenson, A. & Hecht, S.S. (1994) Evidence supporting the role of DNA pyridyl-oxobutylation in rat nasal carcinogenesis by tobacco-specific nitrosamines. *Cancer Res.*, **54**, 1205–1211
- Tulunay, O.E., Hecht, S.S., Carmella, S.G., Zhang, Y., Lemmonds, C., Murphy, S. & Hatsukami, D.K. (2005) Urinary metabolites of a tobacco-specific lung carcinogen in nonsmoking hospitality workers. *Cancer Epidemiol. Biomarkers Prev.*, **14**, 1283–1286
- Upadhyaya, P., Kenney, P.M.J., Hochalter, J.B., Wang, M. & Hecht, S.S. (1999) Tumorigenicity and metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol enantiomers and metabolites in the A/J mouse. *Carcinogenesis*, **20**, 1577–1582
- Upadhyaya, P., Carmella, S.G., Guengerich, F.P. & Hecht, S.S. (2000) Formation and metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol enantiomers *in vitro* in mouse, rat and human tissues. *Carcinogenesis*, **21**, 1233–1238
- Upadhyaya, P., Zimmerman, C.L. & Hecht, S.S. (2002) Metabolism and pharmacokinetics of N'-nitrosornicotine in the patas monkey. *Drug Metab. Dispos.*, **30**, 1115–1122
- Upadhyaya, P., Sturla, S.J., Tretyakova, N., Ziegel, R., Villalta, P.W., Wang, M. & Hecht, S.S. (2003) Identification of adducts produced by the reaction of 4-(acetoxymethylnitrosamino)-1-(3-pyridyl)-1-butanol with deoxyguanosine and DNA. *Chem. Res. Toxicol.*, **16**, 180–190
- Vondracek, M., Xi, Z., Larsson, P., Baker, V., Mace, K., Pfeifer, A., Tjalve, H., Donato, M.T., Gomez-Lechon, M.J. & Grafstrom, R.C. (2001) Cytochrome P450 expression and related metabolism in human buccal mucosa. *Carcinogenesis*, **22**, 481–488
- Vuillemenot, B.R., Pulling, L.C., Palmisano, W.A., Hutt, J.A. & Belinsky, S.A. (2004) Carcinogen exposure differentially modulates RAR-beta promoter hypermethylation, an early and frequent event in mouse lung carcinogenesis. *Carcinogenesis*, **25**, 623–629
- Waddell, W.J. & Marlowe, C. (1980) Localization of [¹⁴C]nitrosornicotine in tissues of the mouse. *Cancer Res.*, **40**, 3518–3523
- Wang, M., Cheng, G., Sturla, S.J., Shi, Y., McIntee, E.J., Villalta, P.W., Upadhyaya, P. & Hecht, S.S. (2003) Identification of adducts formed by pyridyloxobutylation of deoxyguanosine and DNA by 4-(acetoxymethylnitrosamino)-1-(3-pyridyl)-1-butanol, a chemically activated form of tobacco specific carcinogens. *Chem. Res. Toxicol.*, **16**, 616–626
- Weitberg, A.B. & Corvese, D. (1993) Oxygen radicals potentiate the genetic toxicity of tobacco-specific nitrosamines. *Clin. Genet.*, **43**, 88–91
- Weitberg, A.B. & Corvese, D. (1997) Effect of vitamin E and beta-carotene on DNA strand breakage induced by tobacco-specific nitrosamines and stimulated human phagocytes. *J. exp. clin. Cancer Res.*, **16**, 11–14
- West, K.A., Brognard, J., Clark, A.S., Linnoila, I.R., Yang, X., Swain, S.M., Harris, C., Belinsky, S. & Dennis, P.A. (2003) Rapid Akt activation by nicotine and a tobacco carcinogen modulates the phenotype of normal human airway epithelial cells. *J. clin. Invest.*, **111**, 81–90

- West, K.A., Linnoila, I.R., Belinsky, S.A., Harris, C.C. & Dennis, P.A. (2004) Tobacco carcinogen-induced cellular transformation increases activation of the phosphatidylinositol 3'-kinase/Akt pathway *in vitro* and *in vivo*. *Cancer Res.*, **64**, 446–451
- Westra, W.H., Slebos, R.J.C., Offerhaus, G.J.A., Goodman, S.N., Evers, S.G., Kensler, T.W., Askin, F.B., Rodenhuis, S. & Hruban, R.H. (1993) K-*ras* oncogene activation in lung adenocarcinomas from former smokers. Evidence that K-*ras* mutations are an early and irreversible event in the development of adenocarcinoma of the lung. *Cancer*, **72**, 432–438
- Westra, W.H., Baas, I.O., Hruban, R.H., Askin, F.B., Wilson, K., Offerhaus, G.J.A. & Slebos, R.J.C. (1996) K-*ras* oncogene activation in atypical alveolar hyperplasias of the human lung. *Cancer Res.*, **56**, 2224–2228
- Whitlock, J.P., Jr & Denison, M.S. (1995) Induction of cytochrome P450 enzymes that metabolize xenobiotics. In: Ortiz de Montellano, P.R., ed., *Cytochrome P450: Structure, Mechanism, and Biochemistry*, New York, Plenum Press, pp. 367–390
- Wiener, D., Doerge, D.R., Fang, J.L., Upadhyaya, P. & Lazarus, P. (2004a) Characterization of N-glucuronidation of the lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in human liver: Importance of UDP-glucuronosyltransferase 1A4. *Drug Metab. Dispos.*, **32**, 72–79
- Wiener, D., Fang, J.L., Dossett, N. & Lazarus, P. (2004b) Correlation between UDP-glucuronosyltransferase genotypes and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol glucuronidation phenotype in human liver microsomes. *Cancer Res.*, **64**, 1190–1196
- Williams, G.M. & Laspia, M.F. (1979) The detection of various nitrosamines in the hepatocyte primary culture/DNA repair test. *Cancer Lett.*, **6**, 199–206
- Wilp, J., Zwickelpflug, W. & Richter, E. (2002) Nitrosation of dietary myosmine as risk factor of human cancer. *Food chem. Toxicol.*, **40**, 1223–1228
- Wilson, V.L., Weston, A., Manchester, D.K., Trivers, G.E., Roberts, D.W., Kadlubar, F.F., Wild, C.P., Montesano, R., Willey, J.C., Mann, D.L. & Harris, C.C. (1989) Alkyl and aryl carcinogen adducts detected in human peripheral lung. *Carcinogenesis*, **10**, 2149–2153
- Witschi, H., Espiritu, I., Peake, J.L., Wu, K. Maronpot, R.R. & Pinkerton, K.E. (1997) The carcinogenicity of environmental tobacco smoke. *Carcinogenesis*, **18**, 575–586
- Wong, H.L., Zhang, X., Zhang, Q.-Y., Gu, J., Ding, X., Hecht, S.S. & Murphy, S.E. (2005a) Metabolic activation of the tobacco carcinogen 4-(methylnitrosamino)-(3-pyridyl)-1-butanone by cytochrome P450 2A13 in human fetal nasal microsomes. *Chem. Res. Toxicol.*, **18**, 913–918
- Wong, H.L., Murphy, S.E. & Hecht, S.S. (2005b) Cytochrome P450 2A-catalyzed metabolic activation of structurally similar carcinogenic nitrosamines: N'-Nitrosornicotine enantiomers, N-nitrosopiperidine, and N-nitrosopyrrolidine. *Chem. Res. Toxicol.*, **18**, 61–69
- Wu, Z., Upadhyaya, P., Carmella, S.G., Hecht, S.S. & Zimmerman, C.L. (2002) Disposition of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in bile duct-cannulated rats: Stereoselective metabolism and tissue distribution. *Carcinogenesis*, **23**, 171–179
- Wu, W., Ashley, D. & Watson, C. (2003) Simultaneous determination of five tobacco-specific nitrosamines in mainstream cigarette smoke by isotope dilution liquid chromatography/electrospray ionization tandem mass spectrometry. *Anal. Chem.*, **75**, 4827–4832
- Wu, W., Song, S., Ashley, D. & Watson, C. (2004) Assessment of tobacco-specific nitrosamines in the tobacco and mainstream smoke of bidi cigarettes. *Carcinogenesis*, **25**, 283–287

- Wu, W., Zhang, L., Jain, R., Ashley, D. & Watson, C. (2005) Determination of carcinogenic tobacco-specific nitrosamines in mainstream smoke from U.S.-brand and non-U.S.-brand cigarettes from 14 countries. *Nicotine Tob. Res.*, **7**, 443–451
- Yang, G.Y., Yang, Z.Y., Kim, S., Liao, J., Seril, D.N., Chen, X., Smith, T.J. & Yang, C.S. (1997) Characterization of early pulmonary hyperproliferation and tumor progression and their inhibition by black tea in a 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung tumorigenesis model with A/J mice. *Cancer Res.*, **57**, 1889–1894
- Yim, S.H. & Hee, S.S. (2001) Bacterial mutagenicity of some tobacco aromatic nitrogen bases and their mixtures. *Mutat. Res.*, **492**, 13–27
- Yoo, G.H., Washington, J., Piechocki, M., Ensley, J., Shibuya, T., Oda, D. & Wei, W.-Z. (2000) Progression of head and neck cancer in an in vitro model. *Arch. Otolaryngol. Head Neck Surg.*, **126**, 1313–1318
- You, M., Candrian, U., Maronpot, R.R., Stoner, G.D. & Anderson, M.W. (1989) Activation of the Ki-ras protooncogene in spontaneously occurring and chemically induced lung tumors of the strain A mouse. *Proc. natl Acad. Sci. USA*, **86**, 3070–3074
- Zhang, X., Su, T., Zhang, Q.-Y., Gu, J., Caggana, M., Li, H. & Ding, X. (2002) Genetic polymorphisms of the human CYP2A13 gene: Identification of single-nucleotide polymorphisms and functional characterization of an Arg257Cys variant. *J. Pharmacol. exp. Ther.*, **302**, 416–423
- Zhou, H., Zhu, L.X., Li, K. & Hei, T.K. (1999) Radon, tobacco-specific nitrosamine and mutagenesis in mammalian cells. *Mutat. Res.*, **430**, 145–153
- Zhu, S.Y., Cunningham, M.L., Gray, T.E. & Nettesheim, P. (1991) Cytotoxicity, genotoxicity and transforming activity of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) in rat tracheal epithelial cells. *Mutat. Res.*, **261**, 249–259
- Ziegel, R., Shallop, A., Jones, R. & Tretyakova, N. (2003) K-ras gene sequence effects on the formation of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)-DNA adducts. *Chem. Res. Toxicol.*, **16**, 541–550
- Ziegel, R., Shallop, A., Upadhyaya, P., Jones, R. & Tretyakova, N. (2004) Endogenous 5-methylcytosine protects neighboring guanines from N⁷ and O⁶-methylation and O⁶-pyridyloxobutylation by the tobacco carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Biochemistry*, **43**, 540–549
- Zimmerman, C.L., Wu, Z., Upadhyaya, P. & Hecht, S.S. (2004) Stereoselective metabolism and tissue retention in rats of the individual enantiomers of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL), metabolites of the tobacco-specific nitrosamine, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK). *Carcinogenesis*, **25**, 1237–1242
- Zimonjic, D., Popescu, N.C. & DiPaolo, J.A. (1989) Induction of sister chromatid exchanges by tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in human and hamster cells. *Carcinogenesis*, **10**, 753–755