

7. References

- ACGIH® Worldwide (2005) *Documentation of the TLVs® and BEIs® with Other Worldwide Occupational Exposure Values — 2005 CD ROM*, Cincinnati, OH, American Conference of Governmental Industrial Hygienists
- Adkins, B., Jr, Van Stee, E.W., Simmons, J.E. & Eustis, S.L. (1986) Oncogenic response of strain A/J mice to inhaled chemicals. *J. Toxicol. environ. Health*, **17**, 311–322
- Aguirell, E., Cederberg, H., Ehrenberg, L., Lindahl-Kiessling, K., Rannug, U. & Törnqvist, M. (1991) Genotoxic effects of ethylene oxide and propylene oxide: A comparative study. *Mutat. Res.*, **250**, 229–237
- Albertini, R.A. & Sweeney, L.M. (2007) Propylene oxide: Genotoxicity profile of a rodent nasal carcinogen. *Crit. Rev. Toxicol.*, **37**, 489–520
- Allied Signal Chemicals (1993) *Ethylene Oxide*, Morristown, NJ
- Ambroise, D., Moulin, J.-J., Squinazi, F., Protois, J.-C., Fontana, J.-M. & Wild, P. (2005) Cancer mortality among municipal pest-control workers. *Int. Arch. occup. environ. Health*, **78**, 387–393
- Anon. (2004) Product focus — EO-EG, *Chem. Week*, **April 28**, 38
- Anon. (2005) Chemical profile: Ethylene oxide. *Eur. chem. News*, **25 July** (electronic archive)
- Appelgren, L.-E., Eneroth, G. & Grant, C. (1977) Studies on ethylene oxide: Whole-body autoradiography and dominant lethal test in mice. *Proc. Eur. Soc. Toxicol.*, **18**, 315–317
- Appelgren, L.-E., Eneroth, G., Grant, C., Landström, L.-E. & Tenghagen, K. (1978) Testing of ethylene oxide for mutagenicity using the micronucleus test in mice and rats. *Acta pharmacol. toxicol.*, **43**, 69–71
- Arms, A.D. & Travis, C.C. (1988) *Reference Physiological Parameters in Pharmacokinetic Modeling* (Report No EPA 600 6-88/004), Washington DC, US Environmental Protection Agency.
- Bailey, E., Farmer, P.B. & Shuker, D.E.G. (1987) Estimation of exposure to alkylating carcinogens by the GC–MS determination of adducts to hemoglobin and nucleic acid bases in urine. *Arch. Toxicol.*, **60**, 187–191
- Bastlová, T., Andersson, B., Lambert, B. & Kolman, A. (1993) Molecular analysis of ethylene oxide-induced mutations at the HPRT locus in human diploid fibroblasts. *Mutat. Res.*, **287**, 283–292
- Baumbach, N., Herzog, V. & Schiller, F. (1987) [In vitro studies on the permeation of ethylene oxide through human skin]. *Dermatol. Monatsschr.*, **173**, 328–332 (in German)
- Benson, L.O. & Teta, M.J. (1993) Mortality due to pancreatic and lymphopoietic cancers in chlorohydrin production workers. *Br. J. ind. Med.*, **50**, 710–716
- Bhanot, O.S., Singh, U.S. & Solomon, J.J. (1994) The role of 3-hydroxyethyluridine in mutagenesis by ethylene oxide. *J. biol. Chem.*, **269**, 30056–30064
- Bisanti, L., Maggini, M., Raschetti, R., Alegiani, S.S., Ippolito, F.M., Caffari, B., Segnan, N. & Ponti, A. (1993) Cancer mortality in ethylene oxide workers. *Br. J. ind. Med.*, **50**, 317–324
- Boeniger, M. (1988a) *Health Hazard Evaluation Report, University of Cincinnati Hospital, Cincinnati, OH* (Report No. HETA 86-508), Cincinnati, OH, National Institute for Occupational Safety and Health

- Boeniger, M. (1988b) *Health Hazard Evaluation Report, Humana Audubon Hospital, Louisville, KY* (Report No. HETA 87-378), Cincinnati, OH, National Institute for Occupational Safety and Health
- Boffetta, P., van der Hel, O., Norppa, H., Fabianova, E., Fucic, A., Gundy, S., Lazutka, J., Cebulska-Wasilewska, A., Puskaierova, D., Znaor, A., Kelecseyi, Z., Kurtinaitis, J., Rachtan, J., Forni, A., Vermeulen, R. & Bonassi, S. (2007) Chromosomal aberrations and cancer risk: Results of a cohort study from Central Europe. *Am. J. Epidemiol.*, **165**, 36–43
- Bolt, H.M. (1996) Quantification of endogenous carcinogens: The ethylene oxide paradox. *Biochem. Pharmacol.*, **52**, 1–5
- Bolt, H.M. & Filser, J.G. (1987) Kinetics and disposition in toxicology. Example: Carcinogenic risk estimate for ethylene. *Arch. Toxicol.*, **60**, 73–76
- Bolt, H.M. & Leutbecher, M. (1993) Dose–DNA adduct relationship for ethylene oxide. *Arch. Toxicol.*, **67**, 712–713
- Bolt, H.M. & Thier, R. (2006) Relevance of the deletion polymorphisms of the glutathione S-transferases GSTT1 and GSTM1 in pharmacology and toxicology. *Curr. Drug Metab.*, **7**, 613–628
- Bolt, H.M., Peter, H. & Föst, U. (1988) Analysis of macromolecular ethylene oxide adducts. *Int. Arch. occup. environ. Health*, **60**, 141–144
- Bonassi, S., Znaor, A., Ceppi, M., Lando, C., Chang, W.P., Holland, N., Kirsch-Volders, M., Zeiger, E., Ban, S., Barale, R., Bigatti, M.P., Bolognesi, C., Cebulska-Wasilewska, A., Fabianova, E., Fucic, A., Hagmar, L., Joksic, G., Martelli, A., Migliore, L., Mirkova, E., Scarfi, M.R., Zijno, A., Norppa, H. & Fenech, M. (2007) An increased micronucleus frequency in peripheral blood lymphocytes predicts the risk of cancer in humans. *Carcinogenesis*, **28**, 625–631
- Bono, R., Vincenti, M., Saglia, U., Pignata, C., Russo, R., & Gilli, G. (2002) Tobacco smoke and formation of N-(2-hydroxyethyl)valine in human hemoglobin. *Arch. environ. Health*, **57**, 416–421
- Bowen, D.T., Frew, M.E., Hills, R., Gale, R.E., Wheatley, K., Groves, M.J., Langabeer, S.E., Kottaridis, P.D., Moorman, A.V., Burnett, A.K. & Linch, D.C. (2005) RAS mutation in acute myeloid leukemia is associated with distinct cytogenetic subgroups but does not influence outcome in patients younger than 60 years. *Blood*, **106**, 2113–2119
- Brown, C.D., Wong, B.A. & Fennell, T.R. (1996) In vivo and in vitro kinetics of ethylene oxide metabolism in rats and mice. *Toxicol. appl. Pharmacol.*, **136**, 8–19
- Brown, C.D., Asgharian, B., Turner, M.J. & Fennell, T.R. (1998) Ethylene oxide dosimetry in the mouse. *Toxicol. appl. Pharmacol.*, **148**, 215–221
- Bru gnone, F., Perbellini, L., Faccini, G. & Pasini, F. (1985) Concentration of ethylene oxide in the alveolar air of occupationally exposed workers. *Am. J. ind. Med.*, **8**, 67–72
- Bru gnone, F., Perbellini, L., Faccini, G.B., Pasini, F., Bartolucci, G.B. & DeRosa, E. (1986) Ethylene oxide exposure. Biological monitoring by analysis of alveolar air and blood. *Int. Arch. occup. environ. Health*, **58**, 105–112
- Burgaz, S., Rezanko, R., Kara, S. & Karakaya, A.E. (1992) Thioethers in urine of sterilization personnel exposed to ethylene oxide. *J. clin. pharm. Ther.*, **17**, 169–172
- Byrne, J.L. & Marshall, C.J. (1998) The molecular pathophysiology of myeloid leukaemias: Ras revisited. *Br. J. Haematol.*, **100**, 256–264

- Calleman, C.J., Ehrenberg, L., Jansson, B., Osterman-Golkar, S., Segerbäck, D., Svensson, K. & Wachtmeister, C.A. (1978) Monitoring and risk assessment by means of alkyl groups in hemoglobin in persons occupationally exposed to ethylene oxide. *J. environ. Pathol. Toxicol.*, **2**, 427–442
- Chessor, E., Verhoeven, M., Hon, C.-Y. & Teschke, K. (2005) Evaluation of a modified scavenging system to reduce occupational exposure to nitrous oxide in labor and delivery rooms. *J. occup. environ. Hyg.*, **2**, 314–322
- Chien, Y.C., Liu, H.H., Lin Y.C., Su, P.C., Li, L.H., Chang, C.P. Dang, D.T. & Chen C.Y. (2007) Ethylene oxide sterilization in the medical-supply manufacturing industry: Assessment and control of worker exposure. *J. biomed. Mat. Res. B, appl. Biomat.*, **83**, 527–537
- Christiansen, D.H., Andersen, M.K. & Pedersen-Bjergaard, J. (2001) Mutations with loss of heterozygosity of p53 are common in therapy-related myelodysplasia and acute myeloid leukemia after exposure to alkylating agents and significantly associated with deletion or loss of 5q, a complex karyotype, and a poor prognosis. *J. clin. Oncol.*, **19**, 1405–1413
- Christiansen, D.H., Andersen, M.K., Desta, F. & Pedersen-Bjergaard, J. (2005) Mutations of genes in the receptor tyrosine kinase (RTK)/RAS-BRAF signal transduction pathway in therapy-related myelodysplasia and acute myeloid leukemia. *Leukemia*, **19**, 2232–2240
- Clare, M.G., Dean, B.J., de Jong, G. & van Sittert, N.J. (1985) Chromosome analysis of lymphocytes from workers at an ethylene oxide plant. *Mutat. Res.*, **156**, 109–116
- Coggon, D., Harris, E.C., Poole, J. & Palmer, K.T. (2004) Mortality of workers exposed to ethylene oxide: Extended follow up of a British cohort. *Occup. environ. Med.*, **61**, 358–362
- Csanády, G.A., Denk, B., Pütz, C., Kreuzer, P.E., Kessler, W., Baur, C., Gargas, M.L. & Filser, J.G. (2000) A physiological toxicokinetic model for exogenous and endogenous ethylene and ethylene oxide in rat, mouse, and human: Formation of 2-hydroxyethyl adducts with hemoglobin and DNA. *Toxicol. appl. Pharmacol.*, **165**, 1–26
- Cummins, K.J., Schultz, G.R., Lee, J.S., Nelson, J.H. & Reading, J.C. (1987) The development and evaluation of a hydrobromic acid-coated sampling tube for measuring occupational exposures to ethylene oxide. *Am. ind. Hyg. Assoc. J.*, **48**, 563–573
- Currier, M.F., Carlo, G.L., Poston, P.L. & Ledford, W.E. (1984) A cross sectional study of employees with potential occupational exposure to ethylene oxide. *Br. J. ind. Med.*, **41**, 492–498
- Dahlgran, J.R. & Shingleton, C.R. (1987) Determination of ethylene oxide in ethoxylated surfactants and demulsifiers by headspace gas chromatography. *J. Assoc. off. anal. Chem.*, **70**, 796–798
- Daniel, G., Hoffman, W. & McDonald, B. (2004) OSHA compliance issues: Ethylene oxide in a commercial sterilization operation. *J. occup. environ. Hyg.*, **1**, D121–D125
- Danielson, J.W., Snell, R.P. & Oxborrow, G.S. (1990) Detection and quantitation of ethylene oxide, 2-chloroethanol, and ethylene glycol with capillary gas chromatography. *J. chromatogr. Sci.*, **28**, 97–101
- Dellarco, V.L., Generoso, W.M., Sega, G.A., Fowle, J.R., III & Jacobson-Kram, D. (1990) Review of the mutagenicity of ethylene oxide. *Environ. mol. Mutag.*, **16**, 85–103
- Denk, B., Filser, J.G., Oesterle, D., Deml, E. & Greim, H. (1988) Inhaled ethylene oxide induces preneoplastic foci in rat liver. *J. Cancer Res. clin. Oncol.*, **114**, 35–38
- Deschamps, D., Laurent, A.-M., Festy, B. & Conso, F. (1989) [Study of six ethylene oxide sterilization units in the Poor Law Administration of Paris.] *Arch. Mal. prof.*, **50**, 641–649 (in French)

- Devanney, M.T. (2007) *CEH Marketing Research Report – Ethylene Oxide* (Abstract), Zürich, SRI Consulting [available at <http://www.sriconsulting.com/CEH/Public/Reports/654.5000/> accessed 26.02.2008]
- Dever, J.P., George, K.F., Hoffman, W.C. & Soo, H. (2004) Ethylene oxide. In: Kroschwitz, J.I. & Howe-Grant, M., eds, *Kirk Othmer Encyclopedia of Chemical Technology*, Vol. 10, New York, John Wiley & Sons, pp. 632–673 (on line)
- Dunkelberg, H. (1981) [Carcinogenic activity of ethylene oxide and its reaction products 2-chloroethanol, 2-bromoethanol, ethylene glycol and diethylene glycol. I. Carcinogenicity of ethylene oxide in comparison with 1,2-propylene oxide after subcutaneous administration in mice.] *Zbl. Bakt. Hyg. I. Abt. Orig. B*, **174**, 383–404 (in German)
- Dunkelberg, H. (1982) Carcinogenicity of ethylene oxide and 1,2-propylene oxide upon intragastric administration to rats. *Br. J. Cancer*, **46**, 924–933
- Ehrenberg, L. (1991) Detection and measurement of protein adducts: Aspects of risk assessment. *Prog. clin. biol. Res.*, **372**, 79–87
- Ehrenberg, L. & Törnqvist, M. (1992) Use of biomarkers in epidemiology: Quantitative aspects. *Toxicol. Lett.*, **64–65**, 485–492
- Ehrenberg, L., Hiesche, K.D., Osterman-Golkar, S. & Wennberg, I. (1974) Evaluation of genetic risks of alkylating agents: Tissue doses in the mouse from air contaminated with ethylene oxide. *Mutat. Res.*, **24**, 83–103
- Eide, I., Hagemann, R., Zahlse, K., Tabreke, E., Törnqvist, M., Kumar, R., Vodicka, P. & Hemminki, K. (1995) Uptake, distribution, and formation of hemoglobin and DNA adducts after inhalation of C2–C8 1-alkenes (olefins) in the rat. *Carcinogenesis*, **16**, 1603–1609
- Eide, I., Zhao, C., Kumar, R., Hemminki, K., Wu, K.-Y. & Swenberg, J.A. (1999) Comparison of ³²P-postlabeling and high-resolution GC/MS in quantifying N7-(2-hydroxyethyl)guanine adducts. *Chem. Res. Toxicol.*, **12**, 979–984
- Eisenbrand, G., Muller, N., Denkel, E. & Sterzel, W. (1986) DNA adducts and DNA damage by antineoplastic and carcinogenic N-nitrosocompounds. *J. Cancer Res. clin. Oncol.*, **112**, 196–204
- Elliott, L.J., Ringenburg, V.L., Morelli-Schroth, P., Halperin, W.E. & Herrick, R.F. (1988) Ethylene oxide exposures in hospitals. *Appl. ind. Hyg.*, **3**, 141–145
- Ellison, K.S., Dogliotti, E., Connors, T.D., Basu, A.K. & Essigmann, J.M. (1989) Site-specific mutagenesis by O⁶-alkylguanines located in the chromosomes of mammalian cells: Influence of the mammalian O⁶-alkylguanine–DNA alkyltransferase. *Proc. natl Acad. Sci. USA*, **86**, 8620–8624
- Embree, J.W., Lyon, J.P. & Hine, C.H. (1977) The mutagenic potential of ethylene oxide using the dominant-lethal assay in rats. *Toxicol. appl. Pharmacol.*, **40**, 261–267
- Environmental Protection Agency (1992a) Ethylene oxide; tolerances for residues. *US Code fed. Regul.*, **Title 40**, Part 180.151, p. 311
- Environmental Protection Agency (1992b) Ethylene oxide. *US Code fed. Regul.*, **Title 40**, Part 185.2850, p. 456
- Environmental Protection Agency (2000) *Control of Emissions of Hazardous Air Pollutants from Motor Vehicles and Motor Vehicle Fuels. EPA Technical Support Document* (EPA 420-R-00-023), Washington DC

- Farmer, P.B. & Shuker, D.E.G. (1999) What is the significance of increases in background levels of carcinogen-derived protein and DNA adducts? Some considerations for incremental risk assessment. *Mutat. Res.*, **424**, 275–286
- Farmer, P.B., Bailey, E., Gorf, S.M., Törnqvist, M., Osterman-Golkar, S., Kautiainen, A. & Lewis-Enright, D.P. (1986) Monitoring human exposure to ethylene oxide by the determination of haemoglobin adducts using gas chromatography–mass spectrometry. *Carcinogenesis*, **7**, 637–640
- Farmer, P.B., Neumann, H.G., & Henschler, D. (1987) Estimation of exposure of man to substances reacting covalently with macromolecules. *Arch Toxicol.*, **60**, 251–260
- Farmer, P.B., Cordero, R., & Autrup, H. (1996a) Monitoring human exposure to 2-hydroxyethylating carcinogens. *Environ Health Perspect.*, **104**, 449–452
- Farmer, P.B., Sepai, O., Lawrence, R., Autrup, H., Sabro Nielsen, P., Vestergard, A.B., Waters, R., Leuratti, C., Jones, N.J., Stone, J., Baan, R.A., van Delft, J.H., Steenwinkel, M.J., Kyrtopoulos, S.A., Souliotis, V.L., Theodorakopoulos, N., Bacalis, N.C., Natarajan, A.T., Bates, A.D., Haugen, A., Andreassen, A., Ovrebo, S., Shuker, D.E., Amaning, K.S., Schouft, A., Ellul, A., Garner, R.C., Dingley, K.H., Abbondandolo, A.A., Merlo, F., Cole, J., Aldrich, K., Beare, D., Capulas, E., Rowley, G., Waugh, A.P.W., Povey, A.C., Haque, K., Kirsch-Volders, M., van Hummelen, P. & Castelain, P. (1996b) Biomonitoring human exposure to environmental carcinogenic chemicals. *Mutagenesis*, **11**, 363–381
- Farooqi, Z., Törnqvist, M., Ehrenberg, L. & Natarajan, A.T. (1993) Genotoxic effects of ethylene oxide and propylene oxide in mouse bone marrow cells. *Mutat. Res.*, **288**, 223–228
- Fennell, T.R. & Brown, C.D. (2001) A physiologically based pharmacokinetic model for ethylene oxide in mouse, rat, and human. *Toxicol. appl. Pharmacol.*, **173**, 161–175
- Fennell, T.R., Sumner, S.C. & Walker, V.E. (1992) A model for the formation and removal of hemoglobin adducts. *Cancer Epidemiol. Biomarkers Prev.*, **1**, 213–219
- Fennell, T.R., MacNeela, J.P., Morris, R.W., Watson, M., Thompson, C.L. & Bell, D.A. (2000) Hemoglobin adducts from acrylonitrile and ethylene oxide in cigarette smokers: Effects of glutathione S-transferase T1-null and M1-null genotypes. *Cancer Epidemiol. Biomarkers Prev.*, **9**, 705–712
- Filser, J.G. (1992) The closed chamber technique—uptake, endogenous production, excretion, steady-state kinetics and rates of metabolism of gases and vapors. *Arch. Toxicol.*, **66**, 1–10
- Filser, J.G. & Bolt, H.M. (1984) Inhalation pharmacokinetics based on gas uptake studies. VI. Comparative evaluation of ethylene oxide and butadiene monoxide as exhaled reactive metabolites of ethylene and 1,3-butadiene in rats. *Arch. Toxicol.*, **55**, 219–223
- Filser, J.G., Denk, B., Törnqvist, M., Kessler, W. & Ehrenberg, L. (1992) Pharmacokinetics of ethylene in man; body burden with ethylene oxide and hydroxyethylation of hemoglobin due to endogenous and environmental ethylene. *Arch. Toxicol.*, **66**, 157–163
- Filser, J.G., Schwegler, U., Csanády, G.A., Greim, H., Kreuzer, P.E. & Kessler, W. (1993) Species-specific pharmacokinetics of styrene in rat and mouse. *Arch. Toxicol.*, **67**, 517–530
- Filser, J.G., Kreuzer, P.E., Greim, H. & Bolt H.M. (1994) New scientific arguments for regulation of ethylene oxide residues in skin-care products. *Arch. Toxicol.*, **68**, 401–405
- Flores, G.H. (1983) Controlling exposure to alkylene oxides. *Chem. Eng. News*, **79**, 39–43
- Föst, U., Marczynski, B., Kasemann, R. & Peter, H. (1989) Determination of 7-(2-hydroxyethyl)guanine with gas chromatography/mass spectrometry as a parameter for genotoxicity of ethylene oxide. *Arch. Toxicol.*, **Suppl. 13**, 250–253

- Föst, U., Hallier, E., Ottenwälder, H., Bolt, H.M. & Peter, H. (1991) Distribution of ethylene oxide in human blood and its implications for biomonitoring. *Hum. exp. Toxicol.*, **10**, 25–31
- Föst, U., Törnqvist, M., Leutbecher, M., Granath, F., Hallier, E., & Ehrenberg, L. (1995) Effects of variation in detoxification rate on dose monitoring through adducts. *Hum. exp. Toxicol.*, **14**, 201–203
- Fowles, J., Mitchell, J. & McGrath, H. (2001) Assessment of cancer risk from ethylene oxide residues in spices imported into New Zealand. *Food. chem. Toxicol.*, **39**, 1055–1062
- Fuchs, J., Wullenweber, U., Hengstler, J.G., Bienfait, H.G., Hiltl, G. & Oesch, F. (1994) Genotoxic risk for humans due to work place exposure to ethylene oxide: Remarkable individual differences in susceptibility. *Arch. Toxicol.*, **68**, 343–348
- Galloway, S.M., Berry, P.K., Nichols, W.W., Wolman, S.R., Soper, K.A., Stolley, P.D. & Archer, P. (1986) Chromosome aberrations in individuals occupationally exposed to ethylene oxide and in a large control population. *Mutat. Res.*, **170**, 55–74
- Gardner, M.J., Coggon, D., Pannett, B. & Harris, E.C. (1989) Workers exposed to ethylene oxide: A follow up study. *Br. J. ind. Med.*, **46**, 860–865
- Garman, R.H., Snellings, W.M. & Maronpot, R.R. (1985) Brain tumors in F344 rats associated with chronic inhalation exposure to ethylene oxide. *Neurotoxicology*, **6**, 117–138
- Garman, R.H., Snellings, W.M. & Maronpot, R.R. (1986) Frequency, size and location of brain tumours in F-344 rats chronically exposed to ethylene oxide. *Food chem. Toxicol.*, **24**, 145–153
- Garry, V.F., Hozier, J., Jacobs, D., Wade, R.L. & Gray, D.G. (1979) Ethylene oxide: Evidence of human chromosomal effects. *Environ. Mutag.*, **1**, 375–382
- Garry, V.F., Opp, C.W., Wiencke, J.K. & Lakatua, D. (1982) Ethylene oxide induced sister chromatid exchange in human lymphocytes using a membrane dosimetry system. *Pharmacology*, **25**, 214–221
- Generoso, W.M., Cain, K.T., Krishna, M., Sheu, C.W. & Gryder, R.M. (1980) Heritable translocation and dominant-lethal mutation induction with ethylene oxide in mice. *Mutat. Res.*, **73**, 133–142
- Generoso, W.M., Cumming, R.B., Bandy, J.A. & Cain, K.T. (1983) Increased dominant-lethal effects due to prolonged exposure of mice to inhaled ethylene oxide. *Mutat. Res.*, **119**, 377–379
- Generoso, W.M., Cain, K.T., Hughes, L.A., Sega, G.A., Braden, P.W., Gosslee, D.G. & Shelby, M.D. (1986) Ethylene oxide dose and dose-rate effects in the mouse dominant-lethal test. *Environ. Mutag.*, **8**, 1–7
- Generoso, W.M., Cain, K.T., Cornett, C.V., Cacheiro, N.L.A. & Hughes, L.A. (1990) Concentration–response curves for ethylene-oxide-induced heritable translocations and dominant lethal mutations. *Environ. mol. Mutag.*, **16**, 126–131
- Gérin, M. & Tardif, R. (1986) Urinary N-acetyl-S-2-hydroxyethyl-L-cysteine in rats as biological indicator of ethylene oxide exposure. *Fundam. appl. Toxicol.*, **7**, 419–423
- Glaser, Z.R. (1979) Ethylene oxide: Toxicology review and field study results of hospital use. *J. environ. Pathol. Toxicol.*, **12**, 173–208
- Godderis, L., Aka, P., Mateuca, R., Kirsch-Volders, M., Lison, D. & Veulemans, H. (2006) Dose-dependent influence of genetic polymorphisms on DNA damage induced by styrene oxide, ethylene oxide and gamma-radiation. *Toxicology*, **219**, 220–229
- Golberg, L. (1986) *Hazard Assessment of Ethylene Oxide*, Boca Raton, FL, CRC Press, pp. 3–7

- Greenberg, H.L., Ott, M.G. & Shore, R.E. (1990) Men assigned to ethylene oxide production or other ethylene oxide related chemical manufacturing: A mortality study. *Br. J. ind. Med.*, **47**, 221–230
- Greife, A.L., Hornung, R.W., Stayner, L.G. & Steenland, K.N. (1988) Development of a model for use in estimating exposure to ethylene oxide in a retrospective cohort mortality study. *Scand. J. Work Environ. Health*, **14** (Suppl. 1), 29–30
- Hagmar, L., Welinder, H., Lindén, K., Attewell, R., Osterman-Golkar, S. & Törnqvist, M. (1991) An epidemiological study of cancer risk among workers exposed to ethylene oxide using hemoglobin adducts to validate environmental exposure assessments. *Int. Arch. occup. environ. Health*, **63**, 271–277
- Hagmar, L., Mikoczy, Z. & Welinder, H. (1995) Cancer incidence in Swedish sterilant workers exposed to ethylene oxide. *Occup. environ. Med.*, **52**, 154–156
- Hagmar, L., Bonassi, S., Strömberg, U., Brogger, A., Knudsen, L.E., Norppa, H. & Reuterwall, C. (1998) Chromosomal aberrations in lymphocytes predict human cancer: A report from the European Study Group on Cytogenetic Biomarkers and Health (ESCH). *Cancer Res.*, **58**, 4117–4121
- Hagmar, L., Strömberg, U., Bonassi, S., Hansteen, I.L., Knudsen, L.E., Lindholm, C. & Norppa, H. (2004) Impact of types of lymphocyte chromosomal aberrations on human cancer risk: Results from Nordic and Italian cohorts. *Cancer Res.*, **64**, 2258–2263
- Hallier, E., Langhof, T., Dannappel, D., Leutbecher, M., Schröder, K., Goergens, H.W., Müller, A. & Bolt, H.M. (1993) Polymorphism of glutathione conjugation of methyl bromide, ethylene oxide and dichloromethane in human blood: Influence on the induction of sister chromatid exchanges (SCE) in lymphocytes. *Arch. Toxicol.*, **67**, 173–178
- Hansen, J.P., Allen, J., Brock, K., Falconer, J., Helms, M.J., Shaver, G.C. & Strohm, B. (1984) Normal sister chromatid exchange levels in hospital sterilization employees exposed to ethylene oxide. *J. occup. Med.*, **26**, 29–32
- Haufroid, V., Merz, B., Hofmann, A., Tschopp, A., Lison, D., & Hotz, P. (2007) Exposure to ethylene oxide in hospitals: Biological monitoring and influence of glutathione S-transferase and epoxide hydrolase polymorphisms. *Cancer Epidemiol. Biomarkers Prev.*, **16**, 796–802
- Havlicek, C.S., Hilpert, L.R., Dal, G. & Perotti, D. (1992) *Assessment of Ethylene Oxide Concentrations and Emissions from Sterilization and Fumigation Processes. Final Report*, Sacramento, CA, Prepared for Research Division, California Air Resources Board
- Hayes, J.D., Flanagan, J.U. & Jowsey, I.R. (2005) Glutathione transferases. *Annu. Rev. Pharmacol. Toxicol.*, **45**, 51–88
- Health Canada (1999) *Priority Substances List Assessment Report, Ethylene Oxide*, (Canadian Environmental Protection Act, 1999), Ottawa
- Health Canada (2001) *Priority Substances List; Assessment Report Ethylene Oxide*, Ottawa [available at: <http://www.ec.gc.ca/substances/ese/eng/psap/final/ethyleneoxide.cfm>]
- Heiden Associates (1988a) *An Estimate of Industry Costs for Compliance with Two Ethylene Oxide Workplace STEL Scenarios: Ethylene Oxide Production and Ethoxylation Plants*, Washington DC
- Heiden Associates (1988b) *A Medical Products Industry Profile for Evaluating Compliance with Two Ethylene Oxide Workplace STEL Scenarios: 10 ppm STEL and 5 ppm STEL*, Washington DC

- Hemminki, K., Mutanen, P., Saloniemi, I., Niemi, M.-L. & Vainio, H. (1982) Spontaneous abortions in hospital staff engaged in sterilising instruments with chemical agents. *Br. med. J.*, **285**, 1461–1463
- Hengstler, J.G., Fuchs, J., Gebhard, S. & Oesch, F. (1994) Glycolaldehyde causes DNA–protein crosslinks: A new aspect of ethylene oxide genotoxicity. *Mutat. Res.*, **304**, 229–234
- Hengstler, J.G., Fuchs, J., Tanner, B., Oesch-Bartomowicz, B., Hölz, C. & Oesch, F. (1997) Analysis of DNA single-strand breaks in human venous blood: A technique which does not require isolation of white blood cells. *Environ. mol. Mutag.*, **29**, 58–62
- Herrero, M.E., Arand, M., Hengstler, J.G. & Oesch, F. (1997) Recombinant expression of human microsomal epoxide hydrolase protects V79 chinese hamster cells from styrene oxide- but not from ethylene oxide-induced DNA strand breaks. *Environ. mol. Mutag.*, **30**, 429–439
- Hochberg, V., Shi, X.-C., Moorman, W. & Ong, T. (1990) Induction of micronuclei in rat bone marrow and spleen cells by varied dose-rate of ethylene oxide (Abstract No. 91). *Environ. mol. Mutag.*, **15**, 26
- Hoechst Celanese Corp. (1992) *Material Safety Data Sheet: Ethylene Oxide*, Dallas, TX
- Högstedt, L.C. (1988) Epidemiological studies on ethylene oxide and cancer: An updating. In: Bartsch, H., Hemminki, K. & O'Neill, I.K., eds, *Methods for Detecting DNA Damaging Agents in Humans: Applications in Cancer Epidemiology and Prevention* (IARC Scientific Publications No. 89), Lyon, IARC, pp. 265–270
- Högstedt, C., Rohlén, O., Berndtsson, B.S., Axelsson, O. & Ehrenberg, L. (1979a) A cohort study of mortality and cancer incidence in ethylene oxide production workers. *Br. J. ind. Med.*, **36**, 276–280
- Högstedt, C., Malmqvist, N. & Wadman, B. (1979b) Leukemia in workers exposed to ethylene oxide. *J. Am. med. Assoc.*, **241**, 1132–1133
- Högstedt, B., Gullberg, B., Hedner, K., Kolnig, A.-M., Mitelman, F., Skerfving, S. & Widegren, B. (1983) Chromosome aberrations and micronuclei in bone marrow cells and peripheral blood lymphocytes in humans exposed to ethylene oxide. *Hereditas*, **98**, 105–113
- Högstedt, C., Aringer, L. & Gustavsson, A. (1986) Epidemiologic support for ethylene oxide as a cancer-causing agent. *J. Am. med. Assoc.*, **255**, 1575–1578
- Högstedt, B., Bergmark, E., Törnqvist, M. & Osterman-Golkar, S. (1990) Chromosomal aberrations and micronuclei in lymphocytes in relation to alkylation of hemoglobin in workers exposed to ethylene oxide and propylene oxide. *Hereditas*, **113**, 133–138
- Hong, H.-H.L., Houle, C.D., Ton, T.-V.T. & Sills, R.C. (2007) K-ras Mutations in lung tumors and tumors of other organs are consistent with a common mechanism of ethylene oxide tumorigenesis in the B6C3F1 mouse. *Toxicol. Pathol.*, **35**, 81–85
- Hori, H., Yahata, K., Fujishiro, K., Yoshizumi, K., Li, D., Goto, Y. & Higashi, T. (2002) Personal exposure level and environmental ethylene oxide gas concentration in sterilization facilities of hospitals in Japan. *Appl. occup. environ. Hyg.*, **17**, 634–639
- Hornung, R.W., Greife, A.L., Stayner, L.T., Steenland, N.K., Herrick, R.F., Elliott, L.J., Ringenburg, V.L. & Morawetz, J. (1994) Statistical model for prediction of retrospective exposure to ethylene oxide in an occupational mortality study. *Am. J. ind. Med.*, **25**, 825–836
- Houle, C.D., Ton, T.-V.T., Clayton, N., Huff, J., Hong, H.-H.L. & Sills, R.C. (2006) Frequent *p53* and *H-ras* mutations in benzene- and ethylene oxide-induced mammary gland carcinomas from B6C3F1 mice. *Toxicol. Pathol.*, **34**, 752–762

- IARC (1976) *IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Man*, Vol. 11, *Cadmium, Nickel, Some Epoxides, Miscellaneous Industrial Chemicals and General Considerations on Volatile Anaesthetics*, Lyon, pp. 157–167
- IARC (1985) *IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans*, Vol. 36, *Allyl Compounds, Aldehydes, Epoxides and Peroxides*, Lyon, pp. 189–226
- IARC (1987) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Suppl. 7, *Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs Volumes 1 to 42*, Lyon
- IARC (1991) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 53, *Occupational Exposures in Insecticide Application, and Some Pesticides*, Lyon, pp. 179–249
- IARC (1994) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 60, *Some Industrial Chemicals*, Lyon, pp. 73–159
- IARC (1995) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 63, *Dry Cleaning, Some Chlorinated Solvents and Other Industrial Chemicals*, Lyon
- IARC (1997) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 69, *Polychlorinated Dibenzo-para-Dioxins and Polychlorinated Dibenzofurans*, Lyon
- IARC (1999a) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 71, *Re-evaluation of Some Organic Chemicals, Hydrazine and Hydrogen Peroxide*, Lyon
- IARC (1999b) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 73, *Some Chemicals that Cause Tumours of the Kidney or Urinary Bladder in Rodents and Some Other Substances*, Lyon, pp. 131–182
- IARC (2000) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 75, *Ionizing Radiation, Part 1: X- and γ -Radiation, and Neutrons*, Lyon, pp. 121–359
- IARC (2004) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 83, *Tobacco Smoking*, Lyon, p. 51–1187
- IARC (2006) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 88, *Formaldehyde, 2-Butoxyethanol and 1-tert-Butoxypropan-2-ol*, Lyon
- IARC (2009) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Vol. 94, *Ingested Nitrates and Nitrites, and Cyanobacterial Peptide Toxins*, Lyon (in press)
- IPCS (1985) *Ethylene Oxide* (Environmental Health Criteria 55), Geneva, World Health Organization, International Programme on Chemical Safety
- IPCS-CEC (2001) *International Chemical Safety Card 0155*, Geneva, World Health Organization
- IRSST (2005) [Guide to Sampling Contaminants in Workplace Air], 8th rev. Ed. [www.irsst.qc.ca/fr/_outil_100038.html] (in French)
- Jensen, K.G. (1988) Determination of ethylene oxide residues in processed food products by gas-liquid chromatography after derivatization. *Z. Lebensmitt. Untersuch. Forsch.*, **187**, 535–540
- Jenssen, D. & Ramel, C. (1980) The micronucleus test is part of a short-term mutagenicity test program for the prediction of carcinogenicity evaluated by 143 agents tested. *Mutat. Res.*, **75**, 191–202
- Johanson, G. & Filser, J.G. (1992) Experimental data from closed chamber gas uptake studies in rodents suggest lower uptake rate of chemical than calculated from literature values on alveolar ventilation. *Arch. Toxicol.*, **66**, 291–295
- Jones, A.R. & Wells, G. (1981) The comparative metabolism of 2-bromoethanol and ethylene oxide in the rat. *Xenobiotica*, **11**, 763–770
- Joyner, R.E. (1964) Chronic toxicity of ethylene oxide. *Arch. environ. Health*, **8**, 700–710

- Kao, C.-Y. & Giese, R.W. (2005) Measurement of N7-(2'-hydroxyethyl)guanine in human DNA by gas chromatography electron capture mass spectrometry. *Chem. Res. Toxicol.*, **18**, 70–75
- Kardos, L., Széles, G., Gombkötő, G., Szeremi, M., Tompa, A. & Ádány, R. (2003) Cancer deaths among hospital staff potentially exposed to ethylene oxide: An epidemiological analysis. *Environ. mol. Mutag.*, **42**, 59–60
- Karelová, J., Jablonická, A. & Vargová, M. (1987) Results of cytogenetic testing of workers exposed to ethylene oxide. *J. Hyg. Epidemiol. Microbiol. Immunol.*, **31**, 119–126
- Katoh, T., Higashi, K., Inoue, N. & Tanaka, I. (1989) Lipid peroxidation and the metabolism of glutathione in rat liver and brain following ethylene oxide inhalation. *Toxicology*, **58**, 1–9
- Katoh, T., Higashi, K., Inoue, N. & Tanaka, I. (1990) Different responses of cytosolic and mitochondrial glutathione in rat livers after ethylene oxide exposure. *Toxicol. Lett.*, **54**, 235–239
- Kauppinen, T., Toikkanen, J., Pedersen, D., Young, R., Ahrens, W., Boffetta, P., Hansen, J., Kromhout, H., Maqueda Blasco, J., Mirabelli, D., de la Orden-Rivera, V., Pannett, B., Plato, N., Savela, A., Vincent, R. & Kogevinas, M. (2000) Occupational exposure to carcinogens in the European Union. *Occup. environ. Med.*, **57**, 10–18 [data partially available on the CAREX web site: http://www.ttl.fi/NR/rdonlyres/407B368B-26EF-475D-8F2B-DA0024B853E0/0/5_exposures_by_agent_and_industry.pdf]
- Kautiainen, A. & Törnqvist, M. (1991) Monitoring exposure to simple epoxides and alkenes through gas chromatographic determination of hemoglobin adducts. *Int. Arch. occup. environ. Health*, **63**, 27–31
- Kelsey, K.T., Wiencke, J.K., Eisen, E.A., Lynch, D.W., Lewis, T.R. & Little, J.B. (1988) Persistently elevated sister chromatid exchanges in ethylene oxide-exposed primates: The role of a sub-population of high frequency cells. *Cancer Res.*, **48**, 5045–5050
- Kercher, S.L. & Mortimer, V.D. (1987) Before and after: An evaluation of engineering controls for ethylene oxide sterilization in hospitals. *Appl. ind. Hyg.*, **2**, 7–12
- Kiesselbach, N., Ulm, K., Lange, H.J. & Korallus, U. (1990) A multicentre mortality study of workers exposed to ethylene oxide. *Br. J. ind. Med.*, **47**, 182–188
- Kligerman, A.D., Erexson, G.L., Phelps, M.E. & Wilmer, J.L. (1983) Sister-chromatid exchange induction in peripheral blood lymphocytes of rats exposed to ethylene oxide by inhalation. *Mutat. Res.*, **120**, 37–44
- Koda, S., Kumagai, S. & Ohara, H. (1999) Environmental monitoring and assessment of short-term exposures to hazardous chemicals of a sterilization process in hospital working environments. *Acta med. Okayama*, **53**, 217–223
- Koga, M., Hori, H., Tanaka, I., Akiyama, T. & Inoue, N. (1985) [Quantitative analysis of urinary ethylene glycol in rats exposed to ethylene oxide]. *J. UOEH.*, **7**, 45–49 (in Japanese)
- Koga, M., Hori, H., Tanaka, I., Akiyama, T. & Inoue, N. (1987) [Analysis of urinary metabolites of rats exposed to ethylene oxide]. *J. UOEH.*, **9**, 167–170 (in Japanese)
- Kolman, A., Näslund, M. & Calleman, C.J. (1986) Genotoxic effects of ethylene oxide and their relevance to human cancer. *Carcinogenesis*, **7**, 1245–1250
- Kolman, A., Näslund, M., Osterman-Golkar, S., Scalia-Tomba, G.-P. & Meyer, A. (1989) Comparative studies of in vitro transformation by ethylene oxide and gamma-radiation of C3H/10T1/2 cells. *Mutagenesis*, **4**, 58–61

- Kolman, A., Bohušová, T., Lambert, B. & Simons, J.W.I.M. (1992) Induction of 6-thioguanine-resistant mutants in human diploid fibroblasts *in vitro* with ethylene oxide. *Environ. mol. Mutag.*, **19**, 93–97
- Kolman, A., Chovanec, M., & Osterman-Golkar, S. (2002) Genotoxic effects of ethylene oxide, propylene oxide and epichlorohydrin in humans: Update review (1990–2001). *Mutat. Res.*, **512**, 173–194
- Kreuzer, P.E. (1992) [Permeation Kinetics of Ethylene Oxide in Gaseous Form and Dissolved Other Matrices Through the Skin of Rats, Hamsters and Humans] (GSF-Bericht 19/92), Neuherberg, GSF-Forschungszentrum für Umwelt und Gesundheit (in German)
- Kring, E.V., Damrell, D.J., Basilio, A.N., Jr, McGibney, P.D., Douglas, J.J., Henry, T.J. & Ansul, G.R. (1984) Laboratory validation and field verification of a new passive air monitoring badge for sampling ethylene oxide in air. *Am. ind. Hyg. Assoc. J.*, **45**, 697–707
- Krishnan, K., Gargas, M.L., Fennell, T.R. & Andersen, M.E. (1992) A physiologically based description of ethylene oxide dosimetry in the rat. *Toxicol. ind. Health*, **8**, 121–140
- Lacson, J. (2003) *CEH Marketing Research Report — Ethylene Oxide*, Zurich, SRI Consulting
- La Montagne, A.D., Oakes, J.M. & Lopez Turley, R.N. (2004) Long-term ethylene oxide exposure trends in US hospitals: Relationship with OSHA regulatory and enforcement actions. *Am. J. public Health*, **94**, 1614–1619
- Latif, F., Moschel, R.C., Hemminki, K. & Dipple, A. (1988) Styrene oxide as a stereochemical probe for the mechanism of aralkylation at different sites on guanosine. *Chem. Res. Toxicol.*, **1**, 364–369
- Laurent, C., Frederic, J. & Leonard, A.Y. (1984) Sister chromatid exchange frequency in workers exposed to high levels of ethylene oxide in a hospital sterilization service. *Int. Arch. occup. environ. Health*, **54**, 33–43
- Lerda, D. & Rizzi, R. (1992) Cytogenetic study of persons occupationally exposed to ethylene oxide. *Mutat. Res.*, **281**, 31–37
- Lewis, S.E., Barnett, L.B., Felton, C., Johnson, F.M., Skow, L.C., Cacheiro, N. & Shelby, M.D. (1986) Dominant visible and electrophoretically expressed mutations induced in male mice exposed to ethylene oxide by inhalation. *Environ. Mutag.*, **8**, 867–872
- Li, F., Segal, A. & Solomon, J.J. (1992) In vitro reaction of ethylene oxide with DNA and characterization of DNA adducts. *Chem.-biol. Interact.*, **83**, 35–54
- Liao, P.-C., Li, C.-M., Hung, C.-W. & Chen S.-H. (2001) Quantitative detection of N⁷-(2-hydroxyethyl)guanine adducts in DNA using high-performance liquid chromatography/electrospray ionization tandem mass spectrometry. *J. mass Spectrom.*, **36**, 336–343
- Lide, D.R., ed. (2005) *CRC Handbook of Chemistry and Physics*, 86th Ed., Boca Raton, FL, CRC Press, pp. 3–408
- Lindahl, T. (1979) DNA glycosylases, endonucleases for apurinic/apyrimidinic sites, and base excision-repair. *Prog. Nucleic Acid Res. mol. Biol.*, **22**, 135–192
- Liou, S.H., Lung, J.C., Chen, Y.H., Yang, T., Hsieh, L.L., Chen, C.J. & Wu, T.N. (1999) Increased chromosome-type chromosome aberration frequencies as biomarkers of cancer risk in a black-foot endemic area. *Cancer Res.*, **59**, 1481–1484
- Loeb, L.A. & Preston, B.D. (1986) Mutagenesis by apurinic/apyrimidinic sites. *Ann. Rev. Genet.*, **20**, 201–230

- Lorenti Garcia, C., Darroudi, F., Bates, A.D. & Natarajan, A.T. (2001) Induction and persistence of micronuclei, sister-chromatid exchanges and chromosomal aberrations in splenocytes and bone-marrow cells of rats exposed to ethylene oxide. *Mutat. Res.*, **492**, 59–67
- Lovegren, B.C. & Koketsu, M. (1977a) *BASF-Wyandotte Corporation, Geismar, Louisiana, Task II, Ethylene Oxide Survey Report of the Plant Contact, June 27–28, 1977* (PB81-229775), Springfield, VA, National Technical Information Service
- Lovegren, B.C. & Koketsu, M. (1977b) *Union Carbide Corporation, Institute, West Virginia, Task II, Ethylene Oxide Survey Report of the Plant Contact, July 15–16, 1977* (PB82-106709), Springfield, VA, National Technical Information Service
- Lovegren, B.C. & Koketsu, M. (1977c) *Union Carbide Corporation, Texas City, Texas, Task II, Ethylene Oxide Survey Report of the Plant Contact, June 8–9, 1977* (PB82-108218), Springfield, VA, National Technical Information Service
- Lucas, L.J. & Teta, M.J. (1996) Breast cancer and ethylene oxide exposure. *Int. J. Epidemiol.*, **25**, 685–686
- Ludeke, B.I. & Kleihues, P. (1988) Formation and persistence of *O*⁶-(2-hydroxyethyl)-2'-deoxyguanosine in DNA for various rat tissues following a single dose of *N*-nitroso-*N*-(2-hydroxyethyl)urea. An immuno-slot-blot study. *Carcinogenesis*, **9**, 147–151
- Lynch, D.W., Lewis, T.R., Moorman, W.J., Burg, J.R., Groth, D.H., Khan, A., Ackerman, L.J. & Cockrell, B.Y. (1984a) Carcinogenic and toxicologic effects of inhaled ethylene oxide and propylene oxide in F344 rats. *Toxicol. appl. Pharmacol.*, **76**, 69–84
- Lynch, D.W., Lewis, T.R., Moorman, W.J., Burg, J.R., Gulati, D.K., Kaur, P. & Sabharwal, P.S. (1984b) Sister-chromatid exchanges and chromosome aberrations in lymphocytes from monkeys exposed to ethylene oxide and propylene oxide by inhalation. *Toxicol. appl. Pharmacol.*, **76**, 85–95
- Major, J., Jakab, M.G. & Tompa, A. (1996) Genotoxicological investigation of hospital nurses occupationally exposed to ethylene-oxide: I. Chromosome aberrations, sister-chromatid exchanges, cell cycle kinetics, and UV-induced DNA synthesis in peripheral blood lymphocytes. *Environ. mol. Mutag.*, **27**, 84–92
- Major, J., Jakab, M.G. & Tompa, A. (1999) The frequency of induced premature centromere division in human populations occupationally exposed to genotoxic chemicals. *Mutat. Res.*, **445**, 241–249
- Maples, K.R. & Dahl, A.R. (1993) Levels of epoxides in blood during inhalation of alkenes and alkenes oxides. *Inhal. Toxicol.*, **5**, 43–54
- Margeson, J.H., Steger, J.L. & Homolya, J.B. (1990) Chromatographic methods for analysis of ethylene oxide in emissions from stationary sources. *J. chromatogr. Sci.*, **28**, 204–209
- Marlowe, D.E., Lao, N.T., Eaton, A.R., Page, B.F.J. & Lao, C.S. (1987) Interlaboratory comparison of analytical methods for residual ethylene oxide in medical device materials. *J. pharm. Sci.*, **76**, 333–337
- Marnett, L.J. (2002) Oxy radicals, lipid peroxidation and DNA damage. *Toxicology*, **181–182**, 219–222
- Marsden, D.A., Jones, D.J.L., Lamb, J.H., Tompkins, E.M., Farmer, P.B. & Brown, K. (2007) Determination of endogenous and exogenously derived N7-(2-hydroxyethyl)guanine adducts in ethylene oxide-treated rats. *Chem. Res. Toxicol.*, **20**, 290–299
- Martis, L., Kroes, R., Darby, T.D., & Woods, E.F. (1982) Disposition kinetics of ethylene oxide, ethylene glycol, and 2-chlorethanol in the dog. *J. Toxicol. environ. Health*, **10**, 847–856

- Mayer, J., Warburton, D., Jeffrey, A.M., Pero, R., Walles, S., Andrews, L., Toor, M., Latriano, L., Wazneh, L., Tang, D., Tsai, W.-Y., Kuroda, M. & Perera, F. (1991) Biologic markers in ethylene oxide-exposed workers and controls. *Mutat. Res.*, **248**, 163–176
- McCammon, J., Orgel, D. & Hill, B. (1990) *Health Hazard Evaluation Report, A.E. Staley Manufacturing Co., Decatur, IL* (Report No. HETA-88-348-2081), Cincinnati, OH, National Institute for Occupational Safety and Health
- McKelvey, J.A. & Zemaitis, M.A. (1986) The effects of ethylene oxide (EO) exposure on tissue glutathione levels in rats and mice. *Drug chem. Toxicol.*, **9**, 51–66
- Morgan, R.W., Claxton, K.W., Divine, B.J., Kaplan, S.D. & Harris, V.B. (1981) Mortality among ethylene oxide workers. *J. occup. Med.*, **23**, 767–770
- Mortimer, V.D., Jr & Kercher, S.L. (1989) *Control Technology for Ethylene Oxide Sterilization in Hospitals* (NIOSH Publ. No. 89-120), Cincinnati, OH, National Institute for Occupational Safety and Health
- Moullisseaux, A., Laurent, A.-M., Fabre, M., Jouan, M. & Festy, B. (1983) [Atmospheric levels of ethylene oxide in the occupational environment of sterilization and disinfection facilities.] *Arch. Mal. prof.*, **44**, 1–14 (in French)
- Müller, M., Krämer, A., Angerer, J. & Hallier, E. (1998) Ethylene oxide–protein adduct formation in humans: Influence of glutathione-S-transferase polymorphisms. *Int. Arch. occup. environ. Health*, **71**, 499–502
- Nakashima, K., Furutani, A., Higashi, K., Okuno, F. & Inoue, N. (1987) Glutathione contents in rat livers after acute and chronic exposure to ethylene oxide. *J. UOEH*, **9**, 355–359
- Natarajan, A.T., Preston, R.J., Dellarco, V., Ehrenberg, L., Generoso, W., Lewis, S. & Tate, A.D. (1995) Ethylene oxide: Evaluation of genotoxicity data and an exploratory assessment of genetic risk. *Mutat. Res.*, **330**, 55–70
- National Institute for Occupational Safety and Health (1987) *NIOSH Manual of Analytical Methods*, 3rd Ed., Suppl. 2 (DHHS (NIOSH) Publ. No. 84-100), Washington DC, US Government Printing Office, pp. 1614-1–1614-6
- National Institute for Occupational Safety and Health (1998) *NIOSH Manual of Analytical Methods*, 4th Ed., Suppl. 2 (DHHS (NIOSH) Publ. No. 98-119), Washington DC, US Government Printing Office
- National Library of Medicine (2005) *Hazardous Substances Data Bank Database*, Bethesda, MD [available at <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>]
- National Library of Medicine (2006) *Toxic Chemical Release Inventory (TRI) Data Banks* [TRI87, TRI88, TRI89, TRI90, TRI91, TRI05], Bethesda, MD
- National Toxicology Program (1987) *Toxicology and Carcinogenesis Studies of Ethylene Oxide (CAS No. 75-21-8) in B6C3F₁ Mice (Inhalation Studies)* (NTP Technical Report No. 326; NIH Publication No. 88-2582), Research Triangle Park, NC
- Newman, M.A. & Freund, E. (1989) *Health Hazard Evaluation Report, Washington Hospital, Washington, PA* (Report No. HETA 89-006-2002), Cincinnati, OH, National Institute for Occupational Safety and Health
- NOES (1993) *National Occupational Exposure Survey (1981–1983)*, Cincinnati, OH, National Institute for Occupational Safety and Health
- Norman, S.A., Berlin, J.A., Soper, K.A., Middendorf, B.F. & Stolley, P.D. (1995) Cancer incidence in a group of workers potentially exposed to ethylene oxide. *Int. J. Epidemiol.*, **24**, 276–284

- Norman, S.A., Berlin, J.A., Soper, K.A., Middendorf, B.F. & Stolley, P.D. (1996) Authors' response: Cancer incidence in a group of workers potentially exposed to ethylene oxide. *Int. J. Epidemiol.*, **25**, 686
- Nygren, J. Cedervall, B., Eriksson, S., Dušinská, M. & Kolman, A. (1994) Induction of DNA strand breaks by ethylene oxide in human diploid fibroblasts. *Environ. mol. Mutag.*, **24**, 161–167
- O'Neil, M.J., ed. (2006) *Merck Index*, 14th Ed., Whitehouse Station, NJ, Merck, p. 651
- Occupational Safety and Health Administration (2005) *Regulatory Review of the Occupational Safety and Health Administration's Ethylene Oxide Standard* (29 CFR 1910.1047), Washington DC
- Olsen, G.W., Lacy, S.E., Bodner, K.M., Chau, M., Arceneaux, T.G., Cartmill, J.B., Ramlow, J.M. & Boswell, J.M. (1997) Mortality from pancreatic and lymphopoietic cancer among workers in ethylene and propylene chlorohydrin production. *Occup. environ. Med.*, **54**, 592–598
- Ong, T., Bi, H.-K., Xing, S., Stewart, J. & Moorman, W. (1993) Induction of sister chromatid exchange in spleen and bone marrow cells of rats exposed by inhalation to different dose rates of ethylene oxide. *Environ. mol. Mutag.*, **22**, 147–151
- Oser, J.L., Crandall, M., Phillips, R. & Marlow, D. (1978a) *Indepth Industrial Hygiene Report of Ethylene Oxide Exposure at Union Carbide Corporation, Institute, West Virginia* (PB82-114786), Springfield, VA, National Technical Information Service
- Oser, J.L., Crandall, M. & Rinsky, R. (1978b) *Industrial Hygiene Survey of Dow Chemical Company, Plaquemine, Louisiana* (PB81-229924), Springfield, VA, National Technical Information Service
- Oser, J.L., Young, M., Boyle, T. & Marlow, D. (1979) *Indepth Industrial Hygiene Report of Ethylene Oxide Exposure at Union Carbide Corporation, South Charleston, West Virginia* (PB82-110024), Springfield, VA, National Technical Information Service
- Osterman-Golkar, S. & Bergmark, E. (1988) Occupational exposure to ethylene oxide. Relation between in vivo dose and exposure dose. *Scand. J. Work Environ. Health*, **14**, 372–377
- Osterman-Golkar, S., Farmer, P.B., Segerbäck, D., Bailey, E., Calleman, C.J., Svensson, K. & Ehrenberg, L. (1983) Dosimetry of ethylene oxide in the rat by quantitation of alkylated histidine in hemoglobin. *Teratog. Carcinog. Mutag.*, **3**, 395–405
- Ott, M.G., Teta, M.J. & Greenberg, H.L. (1989) Lymphatic and hematopoietic tissue cancer in a chemical manufacturing environment. *Am. J. ind. Med.*, **16**, 631–643
- Pauwels, W. & Veulemans, H. (1998) Comparison of ethylene, propylene, and styrene 7,8-oxide in vitro adduct formation on N-terminal valine in human haemoglobin and on N-7-guanine in human DNA. *Mutat. Res.*, **418**, 21–33
- Pedersen-Bjergaard, J. & Rowley, J.D. (1994) The balanced and the unbalanced chromosome aberrations of acute myeloid leukemia may develop in different ways and may contribute differently to malignant transformation. *Blood*, **83**, 2780–2786
- Pedersen-Bjergaard, J., Christiansen, D.H., Desta, F. & Andersen, M.K. (2006) Alternative genetic pathways and cooperating genetic abnormalities in the pathogenesis of therapy-related myelodysplasia and acute myeloid leukemia. *Leukemia*, **20**, 1943–1949
- Pero, R.W., Widegren, B., Högstedt, B. & Mitelman, F. (1981) In vivo and in vitro ethylene oxide exposure of human lymphocytes assessed by chemical stimulation of unscheduled DNA synthesis. *Mutat. Res.*, **83**, 271–289

- Pero, R.W., Bryngelsson, T., Widegren, B., Högstedt, B. & Welinder, H. (1982) A reduced capacity for unscheduled DNA synthesis in lymphocytes from individuals exposed to propylene oxide and ethylene oxide. *Mutat. Res.*, **104**, 193–200
- Picut, C.A., Aoyama, H., Holder, J.W., Gold, L.S., Maronpot, R.R. & Dixon, D. (2003) Bromoethane, chloroethane and ethylene oxide induced uterine neoplasms in B6C3F1 mice from 2-year NTP inhalation bioassays: Pathology and incidence data revisited. *Exp. Toxicol. Pathol.*, **55**, 1–9
- Poirier, V. & Papadopoulos, D. (1982) Chromosomal aberrations induced by ethylene oxide in a human amniotic cell line *in vitro*. *Mutat. Res.*, **104**, 255–260
- Popp, W., Wahrenholz, C., Przygoda, H., Brauksiepe, A., Goch, S., Müller, G., Schell, C. & Norporth, K. (1994) DNA–protein cross-links and sister chromatid exchange frequencies in lymphocytes and hydroxyethyl mercapturic acid in urine of ethylene oxide-exposed hospital workers. *Int. Arch. occup. environ. Health*, **66**, 325–332
- Potter, D., Blair, D., Davies, R., Watson, W.P. & Wright, A.S. (1989) The relationships between alkylation of haemoglobin and DNA in Fischer 344 rats exposed to [¹⁴C]ethylene oxide. *Arch. Toxicol.*, **Suppl. 13**, 254–257
- Preston, R.J. (1999) Cytogenetic effects of ethylene oxide, with an emphasis on population monitoring. *Crit. Rev. Toxicol.*, **29**, 263–282
- Preston, R.J. & Abernethy, D.J. (1993) Studies on the induction of chromosomal aberration and sister chromatid exchange in rats exposed to styrene by inhalation. In: Sorsa, M., Peltonen, K., Vainio, N. & Hemminki, K, eds, *Butadiene and Styrene: Assessment of Health Hazards* (IARC Scientific Publications No. 127), Lyon, IARC, pp. 225–233
- Puskar, M.A. & Hecker, L.H. (1989) Field validation of passive dosimeters for the determination of employee exposures to ethylene oxide in hospital product sterilization facilities. *Am. ind. Hyg. Assoc. J.*, **50**, 30–36
- Puskar, M.A., Nowak, J.L. & Hecker, L.H. (1990) Generation of ethylene oxide permissible exposure limit data with on-site sample analysis using the EO Self-Scan™ passive monitor. *Am. ind. Hyg. Assoc. J.*, **51**, 273–279
- Puskar, M.A., Szopinski, F.G. & Hecker, L.H. (1991) Development and validation of a protocol for field validation of passive dosimeters for ethylene oxide excursion limit monitoring. *Am. ind. Hyg. Assoc. J.*, **52**, 145–150
- Rebsdats, S. & Mayer, D. (2005) Ethylene oxide. In: *Ullmann's Encyclopedia of Industrial Chemistry*, 7th Ed., Weinheim, Wiley–VCH Publishers (on line)
- Recio, L., Donner, M., Abernethy, D., Pluta, L., Steen, A.-M., Wong, B.A., James, A. & Preston, R.J. (2004) *In vivo* mutagenicity and mutation spectrum in the bone marrow and testes of B6C3F1 lacI transgenic mice following inhalation exposure to ethylene oxide. *Mutagenesis*, **19**, 215–222
- Ribeiro, L.R., Rabello-Gay, M.N., Salvadori, D.M.F., Pereira, C.A.B. & Beçak, W. (1987a) Cytogenetic effects of inhaled ethylene oxide in somatic and germ cells of mice. *Arch. Toxicol.*, **59**, 332–335
- Ribeiro, L.R., Salvadori, D.M.F., Pereira, C.A.B. & Beçak, W. (1987b) Activity of ethylene oxide in the mouse sperm morphology test. *Arch. Toxicol.*, **60**, 331–333
- Ribeiro, L.R., Salvadori, D.M.F., Rios, A.C.C., Costa, S.L., Tates, A.D., Törnqvist, M. & Natarajan, A.T. (1994) Biological monitoring of workers occupationally exposed to ethylene oxide. *Mutat. Res.*, **313**, 81–87

- Richmond, G.W., Abrahams, R.H., Nemenzo, J.H. & Hine, C.H. (1985) An evaluation of possible effects on health following exposure to ethylene oxide. *Arch. environ. Health*, **40**, 20–25
- Russell, I.B., Cumming, R.B. & Hunsicker, P.R. (1984) Specific-locus mutation rates in the mouse following inhalation of ethylene oxide and application of the results to estimation of human genetic risk. *Mutat. Res.*, **129**, 381–388
- Rusyn, I., Asakura, S., Li, Y., Kosyk, O., Koc, H., Nakamura, J., Upton, P.B. & Swenberg, J.A. (2005) Effects of ethylene oxide and ethylene inhalation on DNA adducts, apurinic/apryrimidinic sites and expression of base excision DNA repair gene in rat brain, spleen, and liver. *DNA Repair*, **4**, 1099–1110
- Saalo, A., Soosaar, A., Vuorela, R. & Kauppinen, T. (2006) [ASA 2004], Helsinki, Finnish Institute of Occupational Health. [available at: http://www.ttl.fi/NR/rdonlyres/5A54A452-7350-4255-8DF3-AF632D9D2775/0/ASA_2004.pdf] (in Finnish)
- Sadtler Research Laboratories (1991) *Sadtler Standard Spectra. 1981–1991 Supplementary Index*, Philadelphia, PA
- Sangster, J. (1989) Octanol–water partition coefficients of simple organic compounds. *J. phys. chem. Ref. Data*, **18**, 1144
- Sarto, F., Cominato, I., Pinton, A.M., Brovedani, P.G., Faccioli, C.M., Bianchi, V. & Levis, A.G. (1984a) Workers exposed to ethylene oxide have increased incidence of sister chromatid exchange. In: Berlin, A., Draper, M., Hemminki, K. & Vainio, H., eds, *Monitoring Human Exposure to Carcinogenic and Mutagenic Agents* (IARC Scientific Publications No. 59), Lyon, IARC, pp. 413–419
- Sarto, F., Cominato, I., Pinton, A.M., Brovedani, P.G., Faccioli, C.M., Bianchi, V. & Levis, A.G. (1984b) Cytogenetic damage in workers exposed to ethylene oxide. *Mutat. Res.*, **138**, 185–195
- Sarto, F., Clonfero, E., Bartolucci, G.B., Franceschi, C., Chiricolo, M. & Levis, A.G. (1987) Sister chromatid exchanges and DNA repair capability in sanitary workers exposed to ethylene oxide: evaluation of the dose–effect relationship. *Am. J. ind. Med.*, **12**, 625–637
- Sarto, F., Tomanin, R., Giacomelli, L., Iannini, G. & Cupiraggi, A.R. (1990) The micronucleus assay in human exfoliated cells of the nose and mouth: Application to occupational exposures to chromic acid and ethylene oxide. *Mutat. Res.*, **244**, 345–351
- Sarto, F., Törnqvist, M.A., Tomanin, R., Bartolucci, G.B., Osterman-Golkar, S.M. & Ehrenberg, L. (1991) Studies of biological and chemical monitoring of low-level exposure to ethylene oxide. *Scand. J. Work Environ. Health*, **17**, 60–64
- Scheick, C., Spittler, G., & Dasenbrock, C. (1997) Thiodiacetic acid—a metabolite of ethylene oxide. *Z. Naturforsch.*, **52C**, 70–76
- Schettgen, T., Broding, H.C., Angerer, J. & Drexler, H. (2002) Hemoglobin adducts of ethylene oxide, propylene oxide, acrylonitrile and acrylamide-biomarkers in occupational and environmental medicine. *Toxicol. Lett.*, **134**, 65–70
- Schulte, P.A., Boeniger, M., Walker, J.T., Schober, S.E., Pereira, M.A., Gulati, D.K., Wojciechowski, J.P., Garza, A., Froelich, R., Strauss, G., Halperin, W.E., Herrick, R. & Griffith, J. (1992) Biologic markers in hospital workers exposed to low levels of ethylene oxide. *Mutat. Res.*, **278**, 237–251
- Schulte, P.A., Walker, J.T., Boeniger, M.F., Tsuchiya, Y. & Halperin, W.E. (1995) Molecular, cytogenetic, and hematologic effects of ethylene oxide on female hospital workers. *J. occup. med. Health*, **37**, 313–320

- Sega, G.A. & Generoso, E.E. (1988) Measurement of DNA breakage in spermiogenic germ-cell stages of mice exposed to ethylene oxide, using an alkaline elution procedure. *Mutat. Res.*, **197**, 93–99
- Sega, G.A., Generoso, E.E. & Brimer, P.A. (1988) Inhalation exposure-rate of ethylene oxide affects the level of DNA breakage and unscheduled DNA synthesis in spermiogenic stages of the mouse. *Mutat. Res.*, **209**, 177–180
- Sega, G.A., Brimer, P.A. & Generoso, E.E. (1991) Ethylene oxide inhalation at different exposure-rates affects binding levels in mouse germ cells and hemoglobin. Possible explanation for the effect. *Mutat. Res.*, **249**, 339–349
- Segerbäck, D. (1990) Reaction products in hemoglobin and DNA after in vitro treatment with ethylene oxide and *N*-(2-hydroxyethyl)-*N*-nitrosourea. *Carcinogenesis*, **11**, 307–312
- Segerbäck, D. (1994) DNA alkylation by ethylene oxide and some mono-substituted epoxides. In: Hemminki, K., Dipple, A., Shuker, D.E.G., Kadlubar, F.F., Segerbäck, D. & Bartsch, H., eds, *DNA Adducts: Identification and Biological Significance* (IARC Scientific Publications No. 125), Lyon, IARC, pp. 37–47
- Shell Chemicals (2005) *MSDS Ethylene Oxide 28.3.2005* [available at <http://shellchemicals.com/msds/1,1098,1136,00.html> accessed 30/10/07]
- Shore, R.E., Gardner, M.J. & Pannett, B. (1993) Ethylene oxide: An assessment of the epidemiological evidence on carcinogenicity. *Br. J. ind. Med.*, **50**, 971–997
- Shults, R.A. & Seitz, T.A. (1992) *Health Hazard Evaluation Report, Valley Hospital, Palmer, Alaska* (Report No. HETA 91-293-2203), Cincinnati, OH, National Institute for Occupational Safety and Health
- Sisk, S.C., Pluta, L.J., Meyer, K.G., Wong, B.C. & Recio, L. (1997) Assessment of the in vivo mutagenicity of ethylene oxide in the tissues of B6C3F1 *lacI* transgenic mice following inhalation exposure. *Mutat. Res.*, **391**, 153–164
- van Sittert, N.J., de Jong, G., Clare, M.G., Davies, R., Dean, B.J., Wren, L.J. & Wright, A.S. (1985) Cytogenetic, immunological, and haematological effects in workers in an ethylene oxide manufacturing plant. *Br. J. ind. Med.*, **42**, 19–26
- van Sittert, N.J., Beulink, G.D.J., van Vliet, E.W.N. & van der Waal, H. (1993) Monitoring occupational exposure to ethylene oxide by the determination of hemoglobin adducts. *Environ. Health Perspect.*, **99**, 217–220
- van Sittert, N.J., Boogaard, P.J., Natarajan, A.T., Tate, A.D., Ehrenberg, L.G. & Törnqvist, M.A. (2000) Formation of DNA adducts and induction of mutagenic effects in rats following 4 weeks inhalation exposure to ethylene oxide as a basis for cancer risk assessment. *Mutat. Res.*, **447**, 27–48
- Smerhovský, Z., Landa, K., Rössner, P., Brabec, M., Zudová, Z., Hola, N., Pokorná, Z., Marecková, J. & Hurychová, D. (2001) Risk of cancer in an occupationally exposed cohort with increased level of chromosomal aberrations. *Environ. Health Perspect.*, **109**, 41–45
- Snellings, W.M., Weil, C.S. & Maronpot, R.R. (1984) A two-year inhalation study of the carcinogenic potential of ethylene oxide in Fischer 344 rats. *Toxicol. appl. Pharmacol.*, **75**, 105–117
- Sobaszek, A., Hache, J.C., Frimat, P., Akakpo, V., Victoire, G. & Furon D. (1999) Working conditions and health effects of ethylene oxide exposure at hospital sterilization sites. *J. occup. environ. Med.*, **41**, 492–499

- Solomon, J. (1999) Cyclic adducts and intermediates induced by simple epoxides. In: Singer, B. & Bartsch, H., eds, *Exocyclic Adducts in Mutagenesis and Carcinogenesis* (IARC Scientific Publications No. 150), Lyon, IARC, pp. 123–125
- Sorsa, M., Ojajärvi, A. & Salomaa, S. (1990) Cytogenetic surveillance of workers exposed to genotoxic chemicals: Preliminary experiences from a prospective cancer study in a cytogenetic cohort. *Teratog. Carcinog. Mutag.*, **10**, 215–221
- Star, E.G. (1980) [Mutagenic and cytotoxic effect of ethylene oxide on human cell cultures.] *Zbl. Bakt. Hyg. I. Abt. Orig. B*, **170**, 548–556 (in German)
- Stayner, L., Steenland, K., Greife, A., Hornung, R., Hayes, R.B., Nowlin, S., Morawetz, J., Ringenburg, V., Elliot, L. & Halperin, W. (1993) Exposure–response analysis of cancer mortality in a cohort of workers exposed to ethylene oxide. *Am. J. Epidemiol.*, **138**, 787–798
- Steenland, K. & Stayner, L. (1993) An epidemiological study of workers potentially exposed to ethylene oxide. *Br. J. ind. Med.*, **50**, 1125–1126
- Steenland, K., Stayner, L., Greife, A., Halperin, W., Hayes, R., Hornung, R. & Nowlin, S. (1991) Mortality among workers exposed to ethylene oxide. *New Engl. J. Med.*, **324**, 1402–1407
- Steenland, K., Whelan, E., Deddens, J., Stayner, L. & Ward, E. (2003) Ethylene oxide and breast cancer incidence in a cohort study of 7576 women (United States). *Cancer Causes Control*, **14**, 531–539
- Steenland, K., Stayner, L. & Deddens, J. (2004) Mortality analyses in a cohort of 18 235 ethylene oxide exposed workers: Follow up extended from 1987 to 1998. *Occup. environ. Med.*, **61**, 2–7
- Steger, J. (1989) *Analytical Method Evaluation for Measuring Ethylene Oxide Emissions from Commercial Dilute-acid Hydrolytic Control Units* (EPA Report No. EPA-600/3-89-016; US NTIS PB89-155253), Research Triangle Park, NC, US Environmental Protection Agency, Office of Research and Development
- Stolley, P.D., Soper, K.A., Galloway, S.M., Nichols, W.W., Norman, S.A. & Wolman, S.R. (1984) Sister-chromatid exchanges in association with occupational exposure to ethylene oxide. *Mutat. Res.*, **129**, 89–102
- Swaen, G.M.H., Slangen, J.M.M., Ott, M.G., Kusters, E., Van Den Langenbergh, G., Arends, J.W. & Zober, A. (1996) Investigation of a cluster of ten cases of Hodgkin's disease in an occupational setting. *Int. Arch. occup. environ. Health*, **68**, 224–228
- Swenberg, J.A., Fedtke, N., Fennell, T.R. & Walker, V.E. (1990) Relationships between carcinogen exposure, DNA adducts and carcinogenesis. In: Clayson, D.B., Munro, I.C., Shubik, P. & Swenberg, J.A. eds, *Progress in Predictive Toxicology*, Amsterdam, Elsevier Scientific Publications, pp. 161–184
- Szopinski, F.G., Puskar, M.A. & Hecker, L.H. (1991) Field validation of three passive dosimeters for excursion limit monitoring of ethylene oxide. *Am. ind. Hyg. Assoc. J.*, **52**, 151–157
- Sociale Zaken & Werkgelegenheid (2006) [Modification of the rules concerning work conditions.] *Off. State J.*, **28 December**, **252**, 23 (in Dutch)
- Takeshita, M., Chang, C.-N., Johnson, F., Will, S. & Grollman, A.P. (1987) Oligodeoxynucleotides containing synthetic abasic sites. *J. biol. Chem.*, **262**, 10171–10179
- Tan, E.-L., Cumming, R.B. & Hsie, A.W. (1981) Mutagenicity and cytotoxicity of ethylene oxide in the CHO/HGPRT system. *Environ. Mutag.*, **3**, 683–686
- Tardif, R., Goyal, R., Brodeur, J., & Gérin, M. (1987) Species differences in the urinary disposition of some metabolites of ethylene oxide. *Fundam. appl. Toxicol.*, **9**, 448–453

- Tates, A.D., Grummt, T., Törnqvist, M., Farmer, P.B., van Dam, F.J., van Mossel, H., Schoemaker, H.M., Osterman-Golkar, S., Uebel, C., Tang, Y.S., Zwinderman, A.H., Natarajan, A.T. & Ehrenberg, L. (1991) Biological and chemical monitoring of occupational exposure to ethylene oxide. *Mutat. Res.*, **250**, 483–497
- Tates, A.D., Boogaard, P.J., Darroudi, F., Natarajan, A.T., Caubo, M.E. & van Sittert, N.J. (1995) Biological effect monitoring in industrial workers following incidental exposure to high concentrations of ethylene oxide. *Mutat. Res.*, **329**, 63–77
- Tates, A.D., van Dam, F.J., Natarajan, A.T., van Teylingen, C.M.M., de Zwart, F.A., Zwinderman, A.H., van Sittert, N.J., Nilsern, A., Nilsern, O.G., Zahlsen, K., Magnusson, A.-L. & Törnqvist, M. (1999) Measurement of HPRT mutations in splenic lymphocytes and haemoglobin adducts in erythrocytes of Lewis rats exposed to ethylene oxide. *Mutat. Res.*, **431**, 397–415
- Tavares, R., Ramos, P., Palminha, J., Bispo, M.A., Paz, I., Bras, A., Rueff, J., Farmer, P.B. & Bailey, E. (1994) Transplacental exposure to genotoxins. Evaluation in haemoglobin of hydroxyethylvaline adduct levels in smoking and non-smoking mothers and their newborns. *Carcinogenesis*, **15**, 1271–1274
- Teta, M.J., Benson, L.O. & Vitale, J.N. (1993) Mortality study of ethylene oxide workers in chemical manufacturing: A 10 year update. *Br. J. ind. Med.*, **50**, 704–709
- Teta, M.J., Sielken, R.L., Jr & Valdez-Flores, C. (1999) Ethylene oxide cancer risk assessment based on epidemiological data: Application of revised regulatory guidelines. *Risk Anal.*, **19**, 1135–1155
- Thier, R. & Bolt, H.M. (2000) Carcinogenicity and genotoxicity of ethylene oxide: New aspects and recent advances. *Crit. Rev. Toxicol.*, **30**, 595–608
- Thier, R., Balkenhol, H., Lewalter, J., Selinski, S., Dommermuth, A. & Bolt, H.M. (2001) Influence of polymorphisms of the human glutathione transferases and cytochrome P450 2E1 enzyme on the metabolism and toxicity of ethylene oxide and acrylonitrile. *Mutat. Res.*, **482**, 41–46
- Thiess, A.M., Schwegler, H., Fleig, I. & Stocker, W.G. (1981) Mutagenicity study of workers exposed to alkylene oxides (ethylene oxide/propylene oxide) and derivatives. *J. occup. Med.*, **23**, 343–347
- Tomkins, D.J., Haines, T., Lawrence, M. & Rosa, N. (1993) A study of sister chromatid exchange and somatic cell mutation in hospital workers exposed to ethylene oxide. *Environ. Health Perspect.*, **101** (Suppl. 3), 159–164
- Tompa, A., Major, J. & Jakab, M.G. (1999) Is breast cancer cluster influenced by environmental and occupational factors among hospital nurses in Hungary? *Pathol. oncol. Res.*, **5**, 117–121
- Tucker, S.P. & Arnold, J.E. (1984) *Evaluation of OSHA Method No. 30 for Ethylene Oxide in Air with 400-mg/200-mg Charcoal Tubes* (NIOSH Report No. 84/02/00; US NTIS PB84-242049), Cincinnati, OH, National Institute for Occupational Safety and Health
- Tucker, J.D., Xu, J., Stewart, J., Baci, P.C., & Ong, T.M. (1986) Detection of sister chromatid exchanges induced by volatile genotoxicants. *Teratog. Carcinog. Mutag.*, **6**, 15–21
- US Census (1980) [http://www2.census.gov/prod2/decennial/documents/1980a_vsC-01.pdf]
- Van Duuren, B.L., Orris, L. & Nelson, N. (1965) Carcinogenicity of epoxides, lactones, and peroxy compounds. Part II. *J. natl Cancer Inst.*, **35**, 707–717
- Vergnes, J.S. & Pritts, I.M. (1994) Effects of ethylene on micronucleus formation in the bone marrow of rats and mice following four week of inhalation exposure. *Mutat. Res.*, **324**, 87–91

- Vogel, E. & Natarajan, A.T. (1982) The relation between reaction kinetics and mutagenic action of mono-functional alkylating agents in higher eukaryotic systems: Interspecies comparisons. In: Hollaender, A. & De Serres, F.J. eds, *Chemical Mutagens VII*, New York, Plenum, pp. 295–336
- Walker, V.E. & Skopek, T.R. (1993) A mouse model for the study of *in vivo* mutational spectra: Sequence specificity of ethylene oxide at the *hprt* locus. *Mutat. Res.*, **288**, 151–162
- Walker, V.E., Fennell, T.R., Boucheron, J.A., Fedtke, N., Ciroussel, F. & Swenberg, J.A. (1990) Macromolecular adducts of ethylene oxide: A literature review and a time-course study on the formation of 7-(2-hydroxyethyl)guanine following exposures of rats by inhalation. *Mutat. Res.*, **233**, 151–164
- Walker, V.E., MacNeela, J.P., Swenberg, J.A., Turner, M.J. & Fennell, T.R. (1992a) Molecular dosimetry of ethylene oxide: Formation and persistence of N-(2-hydroxyethyl)valine in hemoglobin following repeated exposures of rats and mice. *Cancer Res.*, **52**, 4320–4327
- Walker, V.E., Fennell, T.R., Upton, P.B., Skopek, T.R., Prevost, V., Shuker, D.E.G. & Swenberg, J.A. (1992b) Molecular dosimetry of ethylene oxide: Formation and persistence of 7-(2-hydroxyethyl)guanine in DNA following repeated exposures of rats and mice. *Cancer Res.*, **52**, 4328–4334
- Walker, V.E., Fennell, T.R., Upton, P.B., MacNeela, J.P. & Swenberg, J.A. (1993) Molecular dosimetry of DNA and hemoglobin adducts in mice and rats exposed to ethylene oxide. *Environ. Health Perspect.*, **99**, 11–17
- Walker, V.E., Sisk, S.C., Upton, P.B., Wong, B.A. & Recio, L. (1997a) *In vivo* mutagenicity of ethylene oxide at the *hprt* locus in T-lymphocytes of B6C3F1 *lacI* transgenic mice following inhalation exposure. *Mutat. Res.*, **392**, 211–222
- Walker, V.E., Meng, Q. & Clement, N.L. (1997b) Spectra of mutations in *hprt* exon 3 of T-cells from F344 rats and *lacI* transgenic and nontransgenic B6C3F1 mice exposed by inhalation to ethylene oxide. *Environ. mol. Mutag.*, **29** (Suppl. 28), 54
- Walker, V.E., Jones, I.M., Crippen, T.L., Meng, Q., Walker, D.M., Bauer, M.J., Reilly, A.A., Tate, A.D., Nakamura, J., Upton, P.B. & Skopek, T.R. (1999) Relationships between exposure, cell loss and proliferation, and manifestation of *HPRT* mutant T-cells following treatment of preweanling, weanling, and adult male mice with *N*-ethyl-*N*-nitrosourea. *Mutat. Res.*, **431**, 371–388
- Walker, V.E., Wu, K.-Y., Upton, P.B., Ranasinghe, A., Scheller, N., Cho, M.-H., Vergnes, J.S., Skopek, T.R. & Swenberg, J.A. (2000) Biomarkers of exposure and effect as indicators of potential carcinogenic risk arising from *in vivo* metabolism of ethylene to ethylene oxide. *Carcinogenesis*, **21**, 1661–1669
- Weast, R.C. & Astle, M.J. (1985) *CRC Handbook of Data on Organic Compounds*, Vol. I, Boca Raton, FL, CRC Press, p. 627
- WHO (2003) *Ethylene Oxide* (Concise International Chemical Assessment Document 54), Geneva, World Health Organization [available at <http://www.inchem.org/documents/cicads/cicads/cicad54.htm>]
- Wojcik-O'Neill, K.M. & Ello, M. (1991) Equivalency of hydrochloric acid and distilled water extraction media for determining residual ethylene oxide in medical devices. *J. pharm. Sci.*, **80**, 783–784

- Wolfs, P., Dutrieux, M., Scailteur, V., Haxhe, J.-J., Zumofen, M. & Lauwerys, R. (1983) [Monitoring of workers exposed to ethylene oxide in a plant distributing sterilizing gases and in units for sterilizing medical equipment.] *Arch. Mal. prof.*, **44**, 321–328 (in French)
- Wong, O. & Trent, L.S. (1993) An epidemiological study of workers potentially exposed to ethylene oxide. *Br. J. ind. Med.*, **50**, 308–316
- Wu, K.-Y., Ranasinghe, A., Upton, P.B., Walker, V.E. & Swenberg, J.A. (1999a) Molecular dosimetry of endogenous and ethylene oxide-induced N7-(2-hydroxyethyl)guanine formation in tissues of rodents. *Carcinogenesis*, **30**, 1787–1792
- Wu, K.-Y., Scheller, N., Ranasinghe, A., Yen, T.-Y., Sangaiah, R., Giese, R. & Swenberg, J.A. (1999b) A gas chromatography/electron capture/negative chemical ionization high-resolution mass spectrometry method for analysis of endogenous and exogenous N7-(2-hydroxyethyl)guanine in rodents and its potential for human biological monitoring. *Chem. Res. Toxicol.*, **12**, 722–729
- Wu, K.Y., Chiang, S.Y., Huang, T.H., Tseng, Y.S., Chen, Y.L., Kuo, H.W. & Hsieh, C.L. (2004) Formation of N-(2-hydroxyethyl)valine in human hemoglobin-effect of lifestyle factors. *Mutat. Res.*, **559**, 73–82
- Yager, J.W. (1982) Sister chromatid exchanges induced in rabbit lymphocytes by ethylene oxide after inhalation exposure. *Environ. Mutag.*, **4**, 121–134
- Yager, J.W. (1987) Effect of concentration-time parameters on sister-chromatid exchanges induced in rabbit lymphocytes by ethylene oxide inhalation. *Mutat. Res.*, **182**, 343–352
- Yager, J.W. & Benz, R.D. (1982) Sister chromatid exchanges induced in rabbit lymphocytes by ethylene oxide after inhalation exposure. *Environ. Mutag.*, **4**, 121–134
- Yager, J.W., Hines, C.J. & Spear, R.C. (1983) Exposure to ethylene oxide at work increases sister chromatid exchanges in human peripheral lymphocytes. *Science*, **219**, 1221–1223
- Yong, L.C., Schulte, P.A., Wiencke, J.K., Boeniger, M.F., Connally, L.B., Walker, J.T., Whelan, E.A. & Ward, E.M. (2001) Hemoglobin adducts and sister chromatid exchanges in hospital workers exposed to ethylene oxide: Effects of glutathione S-transferase T1 and M1 genotypes. *Cancer Epidemiol. Biomarkers Prev.*, **10**, 539–550
- Yong, L.C., Schulte, P.A., Kao, C.-Y., Giese, R.W., Boeniger, M.F., Strauss, G.H.S., Petersen, M.R. & Wiencke, J.K. (2007) DNA adducts in granulocytes of hospital workers exposed to ethylene oxide. *Am. J. ind. Med.*, **50**, 293–302
- Zamora, P.O., Benson, J.M., Li, A.P. & Brooks, A.L. (1983) Evaluation of an exposure system using cells grown on collagen gels for detecting highly volatile mutagens in the CHO/HGPRT mutation assay. *Environ. Mutag.*, **5**, 795–801
- Zarbl, H. (2006) Cellular oncogenes and carcinogenesis. *In: Warshawsky, D. & Landolph, J.R., Jr, eds, Molecular Carcinogenesis and the Molecular Biology of Cancer*, New York, Taylor and Francis, pp. 103–129
- Zhang, W., Johnson, F., Grollman, A.P. & Shibutani, S. (1995) Miscoding by the exocyclic and related DNA adducts 3,N⁴-etheno-2'-deoxycytidine, 3,N⁴-ethano-2'-deoxycytidine, and 3-(2-hydroxyethyl)-2'-deoxyuridine. *Chem. Res. Toxicol.*, **8**, 157–163
- Zhao, C. & Hemminki, K. (2002) The *in vivo* levels of DNA alkylation products in human lymphocytes are not age dependent: An assay of 7-methyl- and 7-(2-hydroxyethyl)-guanine DNA adducts. *Carcinogenesis*, **23**, 307–310
- Zhao, C., Tyndyk, M., Eide, I. & Hemminki, K. (1999) Endogenous and background DNA adducts by methylating and 2-hydroxyethylating agents. *Mutat. Res.*, **424**, 117–125

Zhong, B.-Z., Gu, Z.-W., Whong, W.-Z., Wallace, W.E. & Ong, T.-M. (1992) Comparative study of micronucleus assay and chromosomal aberration analysis in V79 cells exposed to ethylene oxide. *Teratog. Carcinog. Mutag.*, **11**, 227–233