

## Arsenic and Arsenic Compounds

### References to Supplementary Web Tables, Section 2

- Ades AE, Kazantzis G (1988). Lung cancer in a non-ferrous smelter: the role of cadmium. *Br J Ind Med*, 45:435–442. [PMID:3395580](#)
- Bates MN, Rey OA, Biggs ML *et al.* (2004). Case-control study of bladder cancer and exposure to arsenic in Argentina. *Am J Epidemiol*, 159:381–389 [doi:10.1093/aje/kwh054](#). [PMID:14769642](#)
- Bates MN, Smith AH, Cantor KP (1995). Case-control study of bladder cancer and arsenic in drinking water. *Am J Epidemiol*, 141:523–530. [PMID:7900719](#)
- Beane Freeman LE, Dennis LK, Lynch CF *et al.* (2004). Toenail arsenic content and cutaneous melanoma in Iowa. *Am J Epidemiol*, 160:679–687 [doi:10.1093/aje/kwh267](#). [PMID:15383412](#)
- Berg JW, Burbank F (1972). Correlations between carcinogenic trace metals in water supplies and cancer mortality. *Ann N Y Acad Sci*, 199 1 Geochemical E;249–264 [doi:10.1111/j.1749-6632.1972.tb46460.x](#). [PMID:4506509](#)
- Binks K, Doll R, Gillies M *et al.* (2005). Mortality experience of male workers at a UK tin smelter. *Occup Med (Lond)*, 55:215–226 [doi:10.1093/occmed/kqi026](#). [PMID:15757978](#)
- Buchet JP, Lison D (1998). Mortality by cancer in groups of the Belgian population with a moderately increased intake of arsenic. *Int Arch Occup Environ Health*, 71:125–130 [doi:10.1007/s004200050259](#). [PMID:9553788](#)
- Bulbulyan MA, Jourenkova NJ, Boffetta P *et al.* (1996). Mortality in a cohort of Russian fertilizer workers. *Scand J Work Environ Health*, 22:27–33. [PMID:8685670](#)
- Cebrián ME, Albores A, Aguilar M, Blakely E (1983). Chronic arsenic poisoning in the north of Mexico. *Hum Toxicol*, 2:121–133 [doi:10.1177/096032718300200110](#). [PMID:6840787](#)
- Chen C-J, Chuang Y-C, Lin T-M, Wu H-Y (1985). Malignant neoplasms among residents of a blackfoot disease-endemic area in Taiwan: high-arsenic artesian well water and cancers. *Cancer Res*, 45:5895–5899. [PMID:4053060](#)
- Chen C-J, Chuang Y-C, You S-L *et al.* (1986). A retrospective study on malignant neoplasms of bladder, lung and liver in blackfoot disease endemic area in Taiwan. *Br J Cancer*, 53:399–405. [PMID:3964542](#)
- Chen C-J, Kuo T-L, Wu M-M (1988a). Arsenic and cancers. *Lancet*, 331:414–415 [doi:10.1016/S0140-6736\(88\)91207-X](#). [PMID:2893213](#)
- Chen C-J, Wang C-J (1990). Ecological correlation between arsenic level in well water and age-adjusted mortality from malignant neoplasms. *Cancer Res*, 50:5470–5474. [PMID:2386951](#)
- Chen C-J, Wu M-M, Lee S-S *et al.* (1988b). Atherogenicity and carcinogenicity of high-arsenic artesian well water. Multiple risk factors and related malignant neoplasms of blackfoot disease. *Arteriosclerosis*, 8:452–460. [PMID:3190552](#)
- Chen CL, Hsu LI, Chiou HY *et al.* Blackfoot Disease Study Group (2004). Ingested arsenic, cigarette smoking, and lung cancer risk: a follow-up study in arseniasis-endemic areas in Taiwan. *JAMA*, 292:2984–2990 [doi:10.1001/jama.292.24.2984](#). [PMID:15613666](#)
- Chen SY, Hayes RB, Liang SR *et al.* (1990). Mortality experience of haematite mine workers in China. *Br J Ind Med*, 47:175–181. [PMID:2328225](#)
- Chen W, Bochmann F, Sun Y (2007). Effects of work related confounders on the association between silica exposure and lung cancer: a nested case-control study among Chinese miners and pottery workers. *Int Arch Occup Environ Health*, 80:320–326. [doi:10.1007/s00420-006-0137-0](#) [PMID:16897095](#)

- Chen W, Chen J (2002). Nested case-control study of lung cancer in four Chinese tin mines. *Occup Environ Med*, 59:113–118 [doi:10.1136/oem.59.2.113](https://doi.org/10.1136/oem.59.2.113). PMID:11850554
- Chiang HS, Guo HR, Hong CL *et al.* (1993). The incidence of bladder cancer in the black foot disease endemic area in Taiwan. *Br J Urol*, 71:274–278 [doi:10.1111/j.1464-410X.1993.tb15942.x](https://doi.org/10.1111/j.1464-410X.1993.tb15942.x). PMID:8477313
- Chiou HY, Chiou ST, Hsu YH *et al.* (2001). Incidence of transitional cell carcinoma and arsenic in drinking water: a follow-up study of 8,102 residents in an arseniasis-endemic area in northeastern Taiwan. *Am J Epidemiol*, 153:411–418 [doi:10.1093/aje/153.5.411](https://doi.org/10.1093/aje/153.5.411). PMID:11226969
- Chiou H-Y, Hsueh Y-M, Liaw K-F *et al.* (1995). Incidence of internal cancers and ingested inorganic arsenic: a seven-year follow-up study in Taiwan. *Cancer Res*, 55:1296–1300. PMID:7882325
- Dondon MG, de Vathaire F, Quénel P, Fréry N (2005). Cancer mortality during the 1968–1994 period in a mining area in France. *Eur J Cancer Prev*, 14:297–301. PMID:15902002 [doi:10.1097/00008469-200506000-00017](https://doi.org/10.1097/00008469-200506000-00017)
- Enterline PE, Day R, Marsh GM (1995). Cancers related to exposure to arsenic at a copper smelter. *Occup Environ Med*, 52:28–32 [doi:10.1136/oem.52.1.28](https://doi.org/10.1136/oem.52.1.28). PMID:7697137
- Enterline PE, Marsh GM, Esmen NA *et al.* (1987b). Some effects of cigarette smoking, arsenic, and SO<sub>2</sub> on mortality among US copper smelter workers. *J Occup Med*, 29:831–838. PMID:3681494
- Ferreccio C, González C, Milosavlevic V *et al.* (2000). Lung cancer and arsenic concentrations in drinking water in Chile. *Epidemiology*, 11:673–679. PMID:11055628 [doi:10.1097/00001648-200011000-00010](https://doi.org/10.1097/00001648-200011000-00010)
- Grimsrud TK, Berge SR, Haldorsen T, Andersen A (2005). Can lung cancer risk among nickel refinery workers be explained by occupational exposures other than nickel? *Epidemiology*, 16:146–154 [doi:10.1097/01.ede.0000152902.48916.d7](https://doi.org/10.1097/01.ede.0000152902.48916.d7). PMID:15703528
- Guo H-R, Chiang H-S, Hu H *et al.* (1997). Arsenic in drinking water and incidence of urinary cancers. *Epidemiology*, 8:545–550 [doi:10.1097/00001648-199709000-00012](https://doi.org/10.1097/00001648-199709000-00012). PMID:9270957
- Guo H-R, Lipsitz SR, Hu H, Monson RR (1998). Using ecological data to estimate a regression model for individual data: the association between arsenic in drinking water and incidence of skin cancer. *Environ Res*, 79:82–93 [doi:10.1006/enrs.1998.3863](https://doi.org/10.1006/enrs.1998.3863). PMID:9841806
- Guo H-R, Yu H-S, Hu H, Monson RR (2001). Arsenic in drinking water and skin cancers: cell-type specificity (Taiwan, ROC). *Cancer Causes Control*, 12:909–916 [doi:10.1023/A:1013712203455](https://doi.org/10.1023/A:1013712203455). PMID:11808710
- Hansen ES (1992). A mortality study of Danish stokers. *Br J Ind Med*, 49:48–52. PMID:1733455
- Hinwood AL, Jolley DJ, Sim MR (1999). Cancer incidence and high environmental arsenic concentrations in rural populations: Results of an ecological study. *Int J Environ Health Res*, 9:131–141 [doi:10.1080/09603129973272](https://doi.org/10.1080/09603129973272).
- Hopenhayn-Rich C, Biggs ML, Fuchs A *et al.* (1996). Bladder cancer mortality associated with arsenic in drinking water in Argentina. *Epidemiology*, 7:117–124 [doi:10.1097/00001648-199603000-00003](https://doi.org/10.1097/00001648-199603000-00003). PMID:8834549
- Hopenhayn-Rich C, Biggs ML, Smith AH (1998). Lung and kidney cancer mortality associated with arsenic in drinking water in Córdoba, Argentina. *Int J Epidemiol*, 27:561–569 [doi:10.1093/ije/27.4.561](https://doi.org/10.1093/ije/27.4.561). PMID:9758107
- Hsueh Y-M, Cheng G-S, Wu M-M *et al.* (1995). Multiple risk factors associated with arsenic-induced skin cancer: effects of chronic liver disease and malnutritional status. *Br J Cancer*, 71:109–114. PMID:7819025
- Hsueh Y-M, Chiou H-Y, Huang Y-L *et al.* (1997). Serum β-carotene level, arsenic methylation capability, and incidence of skin cancer. *Cancer Epidemiol Biomarkers Prev*, 6:589–596. PMID:9264271
- Kargas MR, Stukel TA, Morris JS *et al.* (2001). Skin cancer risk in relation to toenail arsenic concentrations in a US population-based case-control study. *Am J Epidemiol*, 153:559–565 [doi:10.1093/aje/153.6.559](https://doi.org/10.1093/aje/153.6.559). PMID:11257063

- Karagas MR, Tosteson TD, Morris JS *et al.* (2004). Incidence of transitional cell carcinoma of the bladder and arsenic exposure in New Hampshire. *Cancer Causes Control*, 15:465–472 [doi:10.1023/B:CACO.0000036452.55199.a3](https://doi.org/10.1023/B:CACO.0000036452.55199.a3). PMID:15286466
- Kennedy C, Bajdik CD, Willemze R, Bouwes Bavinck JN (2005). Chemical exposures other than arsenic are probably not important risk factors for squamous cell carcinoma, basal cell carcinoma and malignant melanoma of the skin. *Br J Dermatol*, 152:194–197 [doi:10.1111/j.1365-2133.2005.06411.x](https://doi.org/10.1111/j.1365-2133.2005.06411.x). PMID:15656837
- Kurttio P, Pukkala E, Kahelin H *et al.* (1999). Arsenic concentrations in well water and risk of bladder and kidney cancer in Finland. *Environ Health Perspect*, 107:705–710 [doi:10.1289/ehp.99107705](https://doi.org/10.1289/ehp.99107705). PMID:10464069
- Kusiak RA, Ritchie AC, Muller J, Springer J (1993). Mortality from lung cancer in Ontario uranium miners. *Br J Ind Med*, 50:920–928. PMID:8217852
- Kusiak RA, Springer J, Ritchie AC, Muller J (1991). Carcinoma of the lung in Ontario gold miners: possible aetiological factors. *Br J Ind Med*, 48:808–817. PMID:1663386
- Lee-Feldstein A (1989). A comparison of several measures of exposure to arsenic. Matched case-control study of copper smelter employees. *Am J Epidemiol*, 129:112–124. PMID:2910055
- Lewis DR, Southwick JW, Ouellet-Hellstrom R *et al.* (1999). Drinking water arsenic in Utah: A cohort mortality study. *Environ Health Perspect*, 107:359–365 [doi:10.1289/ehp.99107359](https://doi.org/10.1289/ehp.99107359). PMID:10210691
- Liaw J, Marshall G, Yuan Y *et al.* (2008). Increased childhood liver cancer mortality and arsenic in drinking water in northern Chile. *Cancer Epidemiol Biomarkers Prev*, 17:1982–1987 [doi:10.1158/1055-9965.EPI-07-2816](https://doi.org/10.1158/1055-9965.EPI-07-2816). PMID:18708388
- Lubin JH, Pottern LM, Stone BJ, Fraumeni JF Jr (2000). Respiratory cancer in a cohort of copper smelter workers: results from more than 50 years of follow-up. *Am J Epidemiol*, 151:554–565. PMID:10733037
- Mabuchi K, Lilienfeld AM, Snell LM (1980). Cancer and occupational exposure to arsenic: a study of pesticide workers. *Prev Med*, 9:51–77 [doi:10.1016/0091-7435\(80\)90059-6](https://doi.org/10.1016/0091-7435(80)90059-6). PMID:7360731
- Marshall G, Ferreccio C, Yuan Y *et al.* (2007). Fifty-year study of lung and bladder cancer mortality in Chile related to arsenic in drinking water. *J Natl Cancer Inst*, 99:920–928 [doi:10.1093/jnci/djm004](https://doi.org/10.1093/jnci/djm004). PMID:17565158
- McLaughlin JK, Chen JQ, Dosemeci M *et al.* (1992). A nested case-control study of lung cancer among silica exposed workers in China. *Br J Ind Med*, 49:167–171. PMID:1313281
- Michaud DS, Hartman TJ, Taylor PR *et al.* (2002). No Association between toenail selenium levels and bladder cancer risk. *Cancer Epidemiol Biomarkers Prev*, 11:1505–1506. PMID:12433737
- Morton W, Starr G, Pohl D *et al.* (1976). Skin cancer and water arsenic in Lane County, Oregon. *Cancer*, 37:2523–2532 [doi:10.1002/1097-0142\(197605\)37:5<2523::AID-CNCR2820370545>3.0.CO;2-B](https://doi.org/10.1002/1097-0142(197605)37:5<2523::AID-CNCR2820370545>3.0.CO;2-B). PMID:1260732
- Nakadaira H, Endoh K, Katagiri M, Yamamoto M (2002). Elevated mortality from lung cancer associated with arsenic exposure for a limited duration. *J Occup Environ Med*, 44:291–299 [doi:10.1097/00043764-200203000-00017](https://doi.org/10.1097/00043764-200203000-00017). PMID:11911031
- Pesch B, Ranft U, Jakubis P *et al.* (2002). Environmental arsenic exposure from a coal-burning power plant as a potential risk factor for nonmelanoma skin carcinoma: results from a case-control study in the district of Prievidza, Slovakia. *Am J Epidemiol*, 155:798–809 [doi:10.1093/aje/155.9.798](https://doi.org/10.1093/aje/155.9.798). PMID:11978583
- Qiao YL, Taylor PR, Yao SX *et al.* (1997). Risk factors and early detection of lung cancer in a cohort of Chinese tin miners. *Ann Epidemiol*, 7:533–541 [doi:10.1016/S1047-2797\(97\)00115-4](https://doi.org/10.1016/S1047-2797(97)00115-4). PMID:9408549
- Rivara MI, Cebrán M, Corey G *et al.* (1997). Cancer risk in an arsenic-contaminated area of Chile. *Toxicol Ind Health*, 13:321–338. PMID:9200798

- Simonato L, Moulin JJ, Javelaud B *et al.* (1994). A retrospective mortality study of workers exposed to arsenic in a gold mine and refinery in France. *Am J Ind Med*, 25:625–633 [doi:10.1002/ajim.4700250503](https://doi.org/10.1002/ajim.4700250503). [PMID:8030634](#)
- Smith AH, Goycolea M, Haque R, Biggs ML (1998). Marked increase in bladder and lung cancer mortality in a region of Northern Chile due to arsenic in drinking water. *Am J Epidemiol*, 147:660–669. [PMID:9554605](#)
- Smith AH, Marshall G, Yuan Y *et al.* (2006). Increased mortality from lung cancer and bronchiectasis in young adults after exposure to arsenic in utero and in early childhood. *Environ Health Perspect*, 114:1293–1296 [doi:10.1289/ehp.8832](https://doi.org/10.1289/ehp.8832). [PMID:16882542](#)
- Sobel W, Bond GG, Baldwin CL, Ducommun DJ (1988). An update of respiratory cancer and occupational exposure to arsenicals. *Am J Ind Med*, 13:263–270 [doi:10.1002/ajim.4700130207](https://doi.org/10.1002/ajim.4700130207). [PMID:3354578](#)
- Steinmaus C, Yuan Y, Bates MN, Smith AH (2003). Case-control study of bladder cancer and drinking water arsenic in the western United States. *Am J Epidemiol*, 158:1193–1201 [doi:10.1093/aje/kwg281](https://doi.org/10.1093/aje/kwg281). [PMID:14652304](#)
- Thun MJ, Schnorr TM, Smith AB *et al.* (1985). Mortality among a cohort of U.S. cadmium production workers—an update. *J Natl Cancer Inst*, 74:325–333. [PMID:3856046](#)
- Tokudome S, Kuratsune M (1976). A cohort study on mortality from cancer and other causes among workers at a metal refinery. *Int J Cancer*, 17:310–317 [doi:10.1002/ijc.2910170306](https://doi.org/10.1002/ijc.2910170306). [PMID:1254355](#)
- Tollestrup K, Daling JR, Allard J (1995). Mortality in a cohort of orchard workers exposed to lead arsenate pesticide spray. *Arch Environ Health*, 50:221–229 [doi:10.1080/00039896.1995.9940391](https://doi.org/10.1080/00039896.1995.9940391). [PMID:7618955](#)
- Tsai S-M, Wang T-N, Ko Y-C (1999). Mortality for certain diseases in areas with high levels of arsenic in drinking water. *Arch Environ Health*, 54:186–193 [doi:10.1080/0003989909602258](https://doi.org/10.1080/0003989909602258). [PMID:10444040](#)
- Tseng WP, Chu HM, How SW *et al.* (1968). Prevalence of skin cancer in an endemic area of chronic arsenicism in Taiwan. *J Natl Cancer Inst*, 40:453–463. [PMID:5644201](#)
- Tsuda T, Babazono A, Yamamoto E *et al.* (1995). Ingested arsenic and internal cancer: a historical cohort study followed for 33 years. *Am J Epidemiol*, 141:198–209. [PMID:7840093](#)
- Wall S (1980). Survival and mortality pattern among Swedish smelter workers. *Int J Epidemiol*, 9:73–87 [doi:10.1093/ije/9.1.73](https://doi.org/10.1093/ije/9.1.73). [PMID:7419333](#)
- Wicklund KG, Daling JR, Allard J, Weiss NS (1988). Respiratory cancer among orchardists in Washington State, 1968 to 1980. *J Occup Med*, 30:561–564. [PMID:3397782](#)
- Wong O, Whorton MD, Foliart DE, Lowengart R (1992). An ecologic study of skin cancer and environmental arsenic exposure. *Int Arch Occup Environ Health*, 64:235–241 [doi:10.1007/BF00378280](https://doi.org/10.1007/BF00378280). [PMID:1468791](#)
- Wu M-M, Kuo T-L, Hwang Y-H, Chen C-J (1989). Dose-response relation between arsenic concentration in well water and mortality from cancers and vascular diseases. *Am J Epidemiol*, 130:1123–1132. [PMID:2589305](#)
- Xuan XZ, Lubin JH, Li JY *et al.* (1993). A cohort study in southern China of tin miners exposed to radon and radon decay products. *Health Phys*, 64:120–131 [doi:10.1097/00004032-199302000-00001](https://doi.org/10.1097/00004032-199302000-00001). [PMID:8449705](#)
- Zaldívar R (1974). Arsenic contamination of drinking water and foodstuffs causing endemic chronic poisoning. *Beitr Pathol*, 151:384–400. [PMID:4838015](#)