

Hepatitis B Virus

References to Supplementary Web Tables, Section 2

- Amin J, Dore GJ, O'Connell DL *et al.* (2006). Cancer incidence in people with hepatitis B or C infection: a large community-based linkage study. *J Hepatol*, 45:197–203.[doi:10.1016/j.jhep.2006.02.014](https://doi.org/10.1016/j.jhep.2006.02.014) [PMID:16684579](https://pubmed.ncbi.nlm.nih.gov/16684579/)
- Anderson LA, Pfeiffer R, Warren JL *et al.* (2008). Hematopoietic malignancies associated with viral and alcoholic hepatitis. *Cancer Epidemiol Biomarkers Prev*, 17:3069–3075.[doi:10.1158/1055-9965.EPI-08-0408](https://doi.org/10.1158/1055-9965.EPI-08-0408) [PMID:18957521](https://pubmed.ncbi.nlm.nih.gov/18957521/)
- Bile K, Aden C, Norder H *et al.* (1993). Important role of hepatitis C virus infection as a cause of chronic liver disease in Somalia. *Scand J Infect Dis*, 25:559–564.[doi:10.3109/00365549309008543](https://doi.org/10.3109/00365549309008543) [PMID:7506842](https://pubmed.ncbi.nlm.nih.gov/7506842/)
- Cenac A, Pedroso ML, Djibo A *et al.* (1995). Hepatitis B, C, and D virus infections in patients with chronic hepatitis, cirrhosis, and hepatocellular carcinoma: a comparative study in Niger. *Am J Trop Med Hyg*, 52:293–296. [PMID:7537942](https://pubmed.ncbi.nlm.nih.gov/7537942/)
- Chang CC, Yu MW, Lu CF *et al.* (1994). A nested case-control study on association between hepatitis C virus antibodies and primary liver cancer in a cohort of 9,775 men in Taiwan. *J Med Virol*, 43:276–280.[doi:10.1002/jmv.1890430315](https://doi.org/10.1002/jmv.1890430315) [PMID:7523582](https://pubmed.ncbi.nlm.nih.gov/7523582/)
- Chen CJ, Yu MW, Wang CJ *et al.* (1993). Multiple risk factors of hepatocellular carcinoma: a cohort study of 13737 male adults in Taiwan. *J Gastroenterol Hepatol*, 8 S1:S83–S87 [doi:10.1111/j.1440-1746.1993.tb01689.x](https://doi.org/10.1111/j.1440-1746.1993.tb01689.x).
- Chen G, Lin W, Shen F *et al.* (2005a). Chronic hepatitis B virus infection and mortality from non-liver causes: results from the Haimen City cohort study. *Int J Epidemiol*, 34:132–137.[doi:10.1093/ije/dyh339](https://doi.org/10.1093/ije/dyh339) [PMID:15659459](https://pubmed.ncbi.nlm.nih.gov/15659459/)
- Chiesa R, Donato F, Tagger A *et al.* (2000). Etiology of hepatocellular carcinoma in Italian patients with and without cirrhosis. *Cancer Epidemiol Biomarkers Prev*, 9:213–216. [PMID:10698484](https://pubmed.ncbi.nlm.nih.gov/10698484/)
- Crook PD, Jones ME, Hall AJ (2003). Mortality of hepatitis B surface antigen-positive blood donors in England and Wales. *Int J Epidemiol*, 32:118–124.[doi:10.1093/ije/dyg039](https://doi.org/10.1093/ije/dyg039) [PMID:12690022](https://pubmed.ncbi.nlm.nih.gov/12690022/)
- Cucuianu A, Patiu M, Duma M *et al.* (1999). Hepatitis B and C virus infection in Romanian non-Hodgkin's lymphoma patients. *Br J Haematol*, 107:353–356.[doi:10.1046/j.1365-2141.1999.01692.x](https://doi.org/10.1046/j.1365-2141.1999.01692.x) [PMID:10583224](https://pubmed.ncbi.nlm.nih.gov/10583224/)
- Dazza MC, Meneses LV, Girard PM *et al.* (1993). Absence of a relationship between antibodies to hepatitis C virus and hepatocellular carcinoma in Mozambique. *Am J Trop Med Hyg*, 48:237–242. [PMID:8383469](https://pubmed.ncbi.nlm.nih.gov/8383469/)
- Donato F, Gelatti U, Tagger A *et al.* (2001). Intrahepatic cholangiocarcinoma and hepatitis C and B virus infection, alcohol intake, and hepatolithiasis: a case-control study in Italy. *Cancer Causes Control*, 12:959–964.[doi:10.1023/A:1013747228572](https://doi.org/10.1023/A:1013747228572) [PMID:11808716](https://pubmed.ncbi.nlm.nih.gov/11808716/)
- Donato F, Tagger A, Chiesa R *et al.* (1997). Hepatitis B and C virus infection, alcohol drinking, and hepatocellular carcinoma: a case-control study in Italy. Brescia HCC Study. *Hepatology*, 26:579–584.[doi:10.1002/hep.510260308](https://doi.org/10.1002/hep.510260308) [PMID:9303486](https://pubmed.ncbi.nlm.nih.gov/9303486/)
- El-Serag HB, Richardson PA, Everhart JE (2001). The role of diabetes in hepatocellular carcinoma: a case-control study among United States Veterans. *Am J Gastroenterol*, 96:2462–2467.[doi:10.1111/j.1572-0241.2001.04054.x](https://doi.org/10.1111/j.1572-0241.2001.04054.x) [PMID:11513191](https://pubmed.ncbi.nlm.nih.gov/11513191/)
- Evans AA, Chen G, Ross EA *et al.* (2002). Eight-year follow-up of the 90,000-person Haimen City cohort: I. Hepatocellular carcinoma mortality, risk factors, and gender differences. *Cancer Epidemiol Biomarkers Prev*, 11:369–376. [PMID:11927497](https://pubmed.ncbi.nlm.nih.gov/11927497/)
- Evans AA, O'Connell AP, Pugh JC *et al.* (1998). Geographic variation in viral load among hepatitis B carriers with differing risks of hepatocellular carcinoma. *Cancer Epidemiol Biomarkers Prev*, 7:559–565. [PMID:9681522](https://pubmed.ncbi.nlm.nih.gov/9681522/)

- Franceschi S, Montella M, Polesel J *et al.* (2006a). Hepatitis viruses, alcohol, and tobacco in the etiology of hepatocellular carcinoma in Italy. *Cancer Epidemiol Biomarkers Prev*, 15:683–689.[doi:10.1158/1055-9965.EPI-05-0702](https://doi.org/10.1158/1055-9965.EPI-05-0702) [PMID:16614109](https://pubmed.ncbi.nlm.nih.gov/16614109/)
- Fukuda K, Shibata A, Hirohata I *et al.* (1993). A hospital-based case-control study on hepatocellular carcinoma in Fukuoka and Saga Prefectures, northern Kyushu, Japan. *Jpn J Cancer Res*, 84:708–714. [PMID:8396564](https://pubmed.ncbi.nlm.nih.gov/8396564/)
- Gelatti U, Covolo L, Franceschini M *et al.*; Brescia HCC Study Group (2005). Coffee consumption reduces the risk of hepatocellular carcinoma independently of its aetiology: a case-control study. *J Hepatol*, 42:528–534.[doi:10.1016/j.jhep.2004.11.039](https://doi.org/10.1016/j.jhep.2004.11.039) [PMID:15868652](https://pubmed.ncbi.nlm.nih.gov/15868652/)
- Gwack J, Hwang SS, Ko KP *et al.* (2007). Fasting serum glucose and subsequent liver cancer risk in a Korean prospective cohort. *J Prev Med Public Health*, 40:23–28.[doi:10.3961/jpmp.2007.40.1.23](https://doi.org/10.3961/jpmp.2007.40.1.23) [PMID:17310595](https://pubmed.ncbi.nlm.nih.gov/17310595/)
- Hadziyannis S, Tabor E, Kaklamani E *et al.* (1995). A case-control study of hepatitis B and C virus infections in the etiology of hepatocellular carcinoma. *Int J Cancer*, 60:627–631.[doi:10.1002/ijc.2910600510](https://doi.org/10.1002/ijc.2910600510) [PMID:7860136](https://pubmed.ncbi.nlm.nih.gov/7860136/)
- Hsing AW, Zhang M, Rashid A *et al.* (2008). Hepatitis B and C virus infection and the risk of biliary tract cancer: a population-based study in China. *Int J Cancer*, 122:1849–1853.[doi:10.1002/ijc.23251](https://doi.org/10.1002/ijc.23251) [PMID:18076042](https://pubmed.ncbi.nlm.nih.gov/18076042/)
- Kaczynski J, Hansson G, Hermodsson S *et al.* (1996). Minor role of hepatitis B and C virus infection in the etiology of hepatocellular carcinoma in a low-endemic area. *Scand J Gastroenterol*, 31:809–813.[doi:10.3109/00365529609010357](https://doi.org/10.3109/00365529609010357) [PMID:8858752](https://pubmed.ncbi.nlm.nih.gov/8858752/)
- Kew MC, Yu MC, Kedda MA *et al.* (1997). The relative roles of hepatitis B and C viruses in the etiology of hepatocellular carcinoma in southern African blacks. *Gastroenterology*, 112:184–187.[doi:10.1016/S0016-5085\(97\)70233-6](https://doi.org/10.1016/S0016-5085(97)70233-6) [PMID:8978357](https://pubmed.ncbi.nlm.nih.gov/8978357/)
- Kim JH, Bang YJ, Park BJ *et al.* (2002). Hepatitis B virus infection and B-cell non-Hodgkin's lymphoma in a hepatitis B endemic area: a case-control study. *Jpn J Cancer Res*, 93:471–477. [PMID:12036441](https://pubmed.ncbi.nlm.nih.gov/12036441/)
- Kirk GD, Lesi OA, Mendy M *et al.* (2004). The Gambia Liver Cancer Study: Infection with hepatitis B and C and the risk of hepatocellular carcinoma in West Africa. *Hepatology*, 39:211–219.[doi:10.1002/hep.20027](https://doi.org/10.1002/hep.20027) [PMID:14752840](https://pubmed.ncbi.nlm.nih.gov/14752840/)
- Kuang SY, Lekawanvijit S, Maneekarn N *et al.* (2005). Hepatitis B 1762T/1764A mutations, hepatitis C infection, and codon 249 p53 mutations in hepatocellular carcinomas from Thailand. *Cancer Epidemiol Biomarkers Prev*, 14:380–384.[doi:10.1158/1055-9965.EPI-04-0380](https://doi.org/10.1158/1055-9965.EPI-04-0380) [PMID:15734961](https://pubmed.ncbi.nlm.nih.gov/15734961/)
- Kumar M, Kumar R, Hissar SS *et al.* (2007). Risk factors analysis for hepatocellular carcinoma in patients with and without cirrhosis: a case-control study of 213 hepatocellular carcinoma patients from India. *J Gastroenterol Hepatol*, 22:1104–1111.[doi:10.1111/j.1440-1746.2007.04908.x](https://doi.org/10.1111/j.1440-1746.2007.04908.x) [PMID:17559381](https://pubmed.ncbi.nlm.nih.gov/17559381/)
- Kuniyoshi M, Nakamuta M, Sakai H *et al.* (2001). Prevalence of hepatitis B or C virus infections in patients with non-Hodgkin's lymphoma. *J Gastroenterol Hepatol*, 16:215–219.[doi:10.1046/j.1440-1746.2001.02406.x](https://doi.org/10.1046/j.1440-1746.2001.02406.x) [PMID:11207904](https://pubmed.ncbi.nlm.nih.gov/11207904/)
- Kuper HE, Tzonou A, Kaklamani E *et al.* (2000). Hepatitis B and C viruses in the etiology of hepatocellular carcinoma; a study in Greece using third-generation assays. *Cancer Causes Control*, 11:171–175.[doi:10.1023/A:1008951901148](https://doi.org/10.1023/A:1008951901148) [PMID:10710202](https://pubmed.ncbi.nlm.nih.gov/10710202/)
- Lee TY, Lee SS, Jung SW *et al.* (2008). Hepatitis B virus infection and intrahepatic cholangiocarcinoma in Korea: a case-control study. *Am J Gastroenterol*, 103:1716–1720.[doi:10.1111/j.1572-0241.2008.01796.x](https://doi.org/10.1111/j.1572-0241.2008.01796.x) [PMID:18557716](https://pubmed.ncbi.nlm.nih.gov/18557716/)
- London WT, Evans AA, Buetow K *et al.* (1995). Molecular and genetic epidemiology of hepatocellular carcinoma: studies in China and Senegal. *Princess Takamatsu Symp*, 25:51–60. [PMID:8875609](https://pubmed.ncbi.nlm.nih.gov/8875609/)
- Lu P, Kuang S, Wang J (1998). [Hepatitis B virus infection and aflatoxin exposure in the development of primary liver cancer]. *Zhonghua Yi Xue Za Zhi (Taipei)*, 78:340–342.
- Lu SN, Lee CM, Changchien CS, Chen CJ (1999). Excess mortality from hepatocellular carcinoma in an HCV-endemic township of an HBV-endemic country (Taiwan). *Trans R Soc Trop Med Hyg*, 93:600–602.[doi:10.1016/S0035-9203\(99\)90063-9](https://doi.org/10.1016/S0035-9203(99)90063-9) [PMID:10717743](https://pubmed.ncbi.nlm.nih.gov/10717743/)
- Marcucci F, Mele A, Spada E *et al.* (2006). High prevalence of hepatitis B virus infection in B-cell non-Hodgkin's lymphoma. *Haematologica*, 91:554–557. [PMID:16585021](https://pubmed.ncbi.nlm.nih.gov/16585021/)

- Nomura A, Stemmermann GN, Chyou PH, Tabor E (1996). Hepatitis B and C virus serologies among Japanese Americans with hepatocellular carcinoma. *J Infect Dis*, 173:1474–1476. [PMID:8648223](#)
- Okuno H, Xie ZC, Lu BY *et al.* (1994). A low prevalence of anti-hepatitis C virus antibody in patients with hepatocellular carcinoma in Guangxi Province, southern China. *Cancer*, 73:58–62. [doi:10.1002/1097-0142\(19940101\)73:1<58::AID-CNCR2820730112>3.0.CO;2-E](#) [PMID:7506119](#)
- Olubuyide IO, Aliyu B, Olaleye OA *et al.* (1997). Hepatitis B and C virus and hepatocellular carcinoma. *Trans R Soc Trop Med Hyg*, 91:38–41. [doi:10.1016/S0035-9203\(97\)90387-4](#) [PMID:9093625](#)
- Omer RE, Kuijsten A, Kadaru AM *et al.* (2004). Population-attributable risk of dietary aflatoxins and hepatitis B virus infection with respect to hepatocellular carcinoma. *Nutr Cancer*, 48:15–21. [doi:10.1207/s15327914nc4801_3](#) [PMID:15203373](#)
- Park BC, Han BH, Ahn SY *et al.* (1995). Prevalence of hepatitis C antibody in patients with chronic liver disease and hepatocellular carcinoma in Korea. *J Viral Hepat*, 2:195–202. [doi:10.1111/j.1365-2893.1995.tb00029.x](#) [PMID:7489347](#)
- Park SC, Jeong SH, Kim J *et al.* (2008). High prevalence of hepatitis B virus infection in patients with B-cell non-Hodgkin's lymphoma in Korea. *J Med Virol*, 80:960–966. [doi:10.1002/jmv.21168](#) [PMID:18428141](#)
- Pyong SJ, Tsukuma H, Hiyama T (1994). Case-control study of hepatocellular carcinoma among Koreans living in Osaka, Japan. *Jpn J Cancer Res*, 85:674–679. [PMID:8071108](#)
- Ribes J, Clèries R, Rubió A *et al.* (2006). Cofactors associated with liver disease mortality in an HBsAg-positive Mediterranean cohort: 20 years of follow-up. *Int J Cancer*, 119:687–694. [doi:10.1002/ijc.21882](#) [PMID:16496403](#)
- Shaib YH, El-Serag HB, Nooka AK *et al.* (2007). Risk factors for intrahepatic and extrahepatic cholangiocarcinoma: a hospital-based case-control study. *Am J Gastroenterol*, 102:1016–1021. [doi:10.1111/j.1572-0241.2007.01104.x](#) [PMID:17324130](#)
- Shin HR, Lee CU, Park HJ *et al.* (1996). Hepatitis B and C virus, *Clonorchis sinensis* for the risk of liver cancer: a case-control study in Pusan, Korea. *Int J Epidemiol*, 25:933–940. [doi:10.1093/ije/25.5.933](#) [PMID:8921477](#)
- Simonetti RG, Cammà C, Fiorello F *et al.* (1992). Hepatitis C virus infection as a risk factor for hepatocellular carcinoma in patients with cirrhosis. A case-control study. *Ann Intern Med*, 116:97–102. [PMID:1309286](#)
- Sonmez M, Bektas O, Yilmaz M *et al.* (2007). The relation of lymphoma and hepatitis B virus/hepatitis C virus infections in the region of East Black Sea, Turkey. *Tumori*, 93:536–539. [PMID:18338485](#)
- Sun CA, Farzadegan H, You SL *et al.* (1996). Mutual confounding and interactive effects between hepatitis C and hepatitis B viral infections in hepatocellular carcinogenesis: a population-based case-control study in Taiwan. *Cancer Epidemiol Biomarkers Prev*, 5:173–178. [PMID:8833617](#)
- Tagger A, Donato F, Ribero ML *et al.* (1999). Case-control study on hepatitis C virus (HCV) as a risk factor for hepatocellular carcinoma: the role of HCV genotypes and the synergism with hepatitis B virus and alcohol. Brescia HCC Study. *Int J Cancer*, 81:695–699. [doi:10.1002/\(SICI\)1097-0215\(19990531\)81:5<695::AID-IJC4>3.0.CO;2-W](#) [PMID:10328218](#)
- Talamini R, Montella M, Crovatto M *et al.* (2004). Non-Hodgkin's lymphoma and hepatitis C virus: a case-control study from northern and southern Italy. *Int J Cancer*, 110:380–385. [doi:10.1002/ijc.20137](#) [PMID:15095303](#)
- Tanaka H, Tsukuma H, Yamano H *et al.* (2004). Prospective study on the risk of hepatocellular carcinoma among hepatitis C virus-positive blood donors focusing on demographic factors, alanine aminotransferase level at donation and interaction with hepatitis B virus. *Int J Cancer*, 112:1075–1080. [doi:10.1002/ijc.20507](#) [PMID:15386355](#)
- Tang B, Kruger WD, Chen G *et al.* (2004). Hepatitis B viremia is associated with increased risk of hepatocellular carcinoma in chronic carriers. *J Med Virol*, 72:35–40. [doi:10.1002/jmv.10559](#) [PMID:14635008](#)
- Tsai JF, Chuang LY, Jeng JE *et al.* (2001). Betel quid chewing as a risk factor for hepatocellular carcinoma: a case-control study. *Br J Cancer*, 84:709–713. [doi:10.1054/bjoc.1999.1597](#) [PMID:11237396](#)

- Tsai JF, Jeng JE, Ho MS *et al.* (1996). Additive effect modification of hepatitis B surface antigen and e antigen on the development of hepatocellular carcinoma. *Br J Cancer*, 73:1498–1502.[doi:10.1038/bjc.1996.283](https://doi.org/10.1038/bjc.1996.283) PMID:8664119
- Wang F, Xu RH, Han B *et al.* (2007). High incidence of hepatitis B virus infection in B-cell subtype non-Hodgkin lymphoma compared with other cancers. *Cancer*, 109:1360–1364.[doi:10.1002/cncr.22549](https://doi.org/10.1002/cncr.22549) PMID:17326056
- Wang LY, Hatch M, Chen CJ *et al.* (1996). Aflatoxin exposure and risk of hepatocellular carcinoma in Taiwan. *Int J Cancer*, 67:620–625.[doi:10.1002/\(SICI\)1097-0215\(19960904\)67:5<620::AID-IJC5>3.0.CO;2-W](https://doi.org/10.1002/(SICI)1097-0215(19960904)67:5<620::AID-IJC5>3.0.CO;2-W) PMID:8782648
- Wang LY, You SL, Lu SN *et al.* (2003). Risk of hepatocellular carcinoma and habits of alcohol drinking, betel quid chewing and cigarette smoking: a cohort of 2416 HBsAg-seropositive and 9421 HBsAg-seronegative male residents in Taiwan. *Cancer Causes Control*, 14:241–250.[doi:10.1023/A:1023636619477](https://doi.org/10.1023/A:1023636619477) PMID:12814203
- Yamamoto S, Kubo S, Hai S *et al.* (2004). Hepatitis C virus infection as a likely etiology of intrahepatic cholangiocarcinoma. *Cancer Sci*, 95:592–595.[doi:10.1111/j.1349-7006.2004.tb02492.x](https://doi.org/10.1111/j.1349-7006.2004.tb02492.x) PMID:15245596
- Yang HI, Lu SN, Liaw YF *et al.*; Taiwan Community-Based Cancer Screening Project Group (2002). Hepatitis B e antigen and the risk of hepatocellular carcinoma. *N Engl J Med*, 347:168–174.[doi:10.1056/NEJMoa013215](https://doi.org/10.1056/NEJMoa013215) PMID:12124405
- Yu MW, You SL, Chang AS *et al.* (1991). Association between hepatitis C virus antibodies and hepatocellular carcinoma in Taiwan. *Cancer Res*, 51:5621–5625. PMID:1655259
- Yu SZ, Zi XL, Chen G (1997b). The relationship between viral hepatitis and primary liver cancer in four areas of China. *Zhonghua Liu Xing Bing Xue Za Zhi*, 18:214–216. PMID:9812521
- Zhang JY, Dai M, Wang X *et al.* (1998). A case-control study of hepatitis B and C virus infection as risk factors for hepatocellular carcinoma in Henan, China. *Int J Epidemiol*, 27:574–578.[doi:10.1093/ije/27.4.574](https://doi.org/10.1093/ije/27.4.574) PMID:9758109
- Zhou YM, Yin ZF, Yang JM *et al.* (2008). Risk factors for intrahepatic cholangiocarcinoma: a case-control study in China. *World J Gastroenterol*, 14:632–635.[doi:10.3748/wjg.14.632](https://doi.org/10.3748/wjg.14.632) PMID:18203300