

Human T-cell Lymphotropic Virus - Type 1

References to Supplementary Web Tables, Section 2

- Adedayo OA, Shehu SM (2004). Human T-cell lymphotropic virus type 1 (HTLV-1) and lymphoid malignancies in Dominica: a seroprevalence study. *Am J Hematol*, 77:336–339.[doi:10.1002/ajh.20208](https://doi.org/10.1002/ajh.20208) [PMID:15551358](https://pubmed.ncbi.nlm.nih.gov/15551358/)
- Arisawa K, Katamine S, Kamihira S *et al.* (2002). A nested case-control study of risk factors for adult T-cell leukemia/lymphoma among human T-cell lymphotropic virus type-I carriers in Japan. *Cancer Causes Control*, 13:657–663.[doi:10.1023/A:1019511224501](https://doi.org/10.1023/A:1019511224501) [PMID:12296513](https://pubmed.ncbi.nlm.nih.gov/12296513/)
- Arisawa K, Sobue T, Yoshimi I *et al.* (2003). Human T-lymphotropic virus type-I infection, survival and cancer risk in southwestern Japan: a prospective cohort study. *Cancer Causes Control*, 14:889–896.[doi:10.1023/B:CACO.0000003853.82298.96](https://doi.org/10.1023/B:CACO.0000003853.82298.96) [PMID:14682446](https://pubmed.ncbi.nlm.nih.gov/14682446/)
- Arisawa K, Soda M, Akahoshi M *et al.* (2006). Human T-cell lymphotropic virus type-1 infection and risk of cancer: 15.4 year longitudinal study among atomic bomb survivors in Nagasaki, Japan. *Cancer Sci*, 97:535–539.[doi:10.1111/j.1349-7006.2006.00212.x](https://doi.org/10.1111/j.1349-7006.2006.00212.x) [PMID:16734733](https://pubmed.ncbi.nlm.nih.gov/16734733/)
- Arisawa K, Soda M, Endo S *et al.* (2000). Evaluation of adult T-cell leukemia/lymphoma incidence and its impact on non-Hodgkin lymphoma incidence in southwestern Japan. *Int J Cancer*, 85:319–324.[doi:10.1002/\(SICI\)1097-0215\(20000201\)85:3<319::AID-IJC4>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1097-0215(20000201)85:3<319::AID-IJC4>3.0.CO;2-B) [PMID:10652420](https://pubmed.ncbi.nlm.nih.gov/10652420/)
- Au WY, Lo JY (2005). HTLV-1-related lymphoma in Hong Kong Chinese. *Am J Hematol*, 78:80–81.[doi:10.1002/ajh.20217](https://doi.org/10.1002/ajh.20217) [PMID:15609285](https://pubmed.ncbi.nlm.nih.gov/15609285/)
- Barbosa HS, Bittencourt AL, Barreto de Araújo I *et al.* (1999). Adult T-cell leukemia/lymphoma in northeastern Brazil: a clinical, histopathologic, and molecular study. *J Acquir Immune Defic Syndr*, 21:65–71. [PMID:10235516](https://pubmed.ncbi.nlm.nih.gov/10235516/)
- Bartholomew C, Jack N, Edwards J *et al.* (1998). HTLV-I serostatus of mothers of patients with adult T-cell leukemia and HTLV-I-associated myelopathy/tropical spastic paraparesis. *J Hum Virol*, 1:302–305. [PMID:10195256](https://pubmed.ncbi.nlm.nih.gov/10195256/)
- Bazarbachi A, Soriano V, Pawson R *et al.* (1997). Mycosis fungoides and Sezary syndrome are not associated with HTLV-I infection: an international study. *Br J Haematol*, 98:927–933.[doi:10.1046/j.1365-2141.1997.3213138.x](https://doi.org/10.1046/j.1365-2141.1997.3213138.x) [PMID:9326191](https://pubmed.ncbi.nlm.nih.gov/9326191/)
- Cabrera ME, Marinov N, Guerra C *et al.* (2003). Chronic lymphoproliferative syndromes in Chile. A prospective study in 132 patients. *Rev Med Chil*, 131:291–298. [PMID:12790078](https://pubmed.ncbi.nlm.nih.gov/12790078/)
- Chan WP, Liang R (1996). Sequence analysis of HTLV-1 provirus associated with adult T-cell leukemia/lymphoma in Hong Kong. *Am J Hematol*, 51:94–95.[doi:10.1002/\(SICI\)1096-8652\(199601\)51:1<94::AID-AJH17>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1096-8652(199601)51:1<94::AID-AJH17>3.0.CO;2-B) [PMID:8571947](https://pubmed.ncbi.nlm.nih.gov/8571947/)
- Farias de Carvalho SM, Pombo de Oliveira MS, Thuler LC *et al.* (1997). HTLV-I and HTLV-II infections in hematologic disorder patients, cancer patients, and healthy individuals from Rio de Janeiro, Brazil. *J Acquir Immune Defic Syndr Hum Retrovirol*, 15:238–242. [PMID:9257659](https://pubmed.ncbi.nlm.nih.gov/9257659/)
- Gabet AS, Mortreux F, Talarmin A *et al.* (2000). High circulating proviral load with oligoclonal expansion of HTLV-1 bearing T cells in HTLV-1 carriers with strongyloidiasis. *Oncogene*, 19:4954–4960.[doi:10.1038/sj.onc.1203870](https://doi.org/10.1038/sj.onc.1203870) [PMID:11042682](https://pubmed.ncbi.nlm.nih.gov/11042682/)
- Gentile TC, Hadlock KG, Uner AH *et al.* (1998). Large granular lymphocyte leukaemia occurring after renal transplantation. *Br J Haematol*, 101:507–512.[doi:10.1046/j.1365-2141.1998.00712.x](https://doi.org/10.1046/j.1365-2141.1998.00712.x) [PMID:9633895](https://pubmed.ncbi.nlm.nih.gov/9633895/)
- Hisada M, Okayama A, Shioiri S *et al.* (1998b). Risk factors for adult T-cell leukemia among carriers of human T-lymphotropic virus type I. *Blood*, 92:3557–3561. [PMID:9808547](https://pubmed.ncbi.nlm.nih.gov/9808547/)
- Hisada M, Okayama A, Spiegelman D *et al.* (2001). Sex-specific mortality from adult T-cell leukemia among carriers of human T-lymphotropic virus type I. *Int J Cancer*, 91:497–499.[doi:10.1002/1097-0215\(20010215\)91:4<497::AID-IJC1044>3.0.CO;2-A](https://doi.org/10.1002/1097-0215(20010215)91:4<497::AID-IJC1044>3.0.CO;2-A) [PMID:11251972](https://pubmed.ncbi.nlm.nih.gov/11251972/)
- Hisada M, Okayama A, Tachibana N *et al.* (1998a). Predictors of level of circulating abnormal lymphocytes among human T-lymphotropic virus type I carriers in Japan. *Int J Cancer*, 77:188–192.[doi:10.1002/\(SICI\)1097-0215\(19980717\)77:2<188::AID-IJC3>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1097-0215(19980717)77:2<188::AID-IJC3>3.0.CO;2-M) [PMID:9650550](https://pubmed.ncbi.nlm.nih.gov/9650550/)

- Hoshida Y, Li T, Dong Z *et al.* (2001). Lymphoproliferative disorders in renal transplant patients in Japan. *Int J Cancer*, 91:869–875.[doi:10.1002/1097-0215\(200002\)9999:9999<::AID-IJC1125>3.0.CO;2-N](https://doi.org/10.1002/1097-0215(200002)9999:9999<::AID-IJC1125>3.0.CO;2-N) [PMID:11275994](https://pubmed.ncbi.nlm.nih.gov/11275994/)
- Kikuchi A, Nishikawa T, Ikeda Y, Yamaguchi K (1997a). Absence of human T-lymphotropic virus type I in Japanese patients with cutaneous T-cell lymphoma. *Blood*, 89:1529–1532. [PMID:9057633](https://pubmed.ncbi.nlm.nih.gov/9057633/)
- Kikuchi A, Ohata Y, Matsumoto H *et al.* (1997b). Anti-HTLV-1 antibody positive cutaneous T-cell lymphoma. *Cancer*, 79:269–274.[doi:10.1002/\(SICI\)1097-0142\(19970115\)79:2<269::AID-CNCR10>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1097-0142(19970115)79:2<269::AID-CNCR10>3.0.CO;2-Z) [PMID:9010100](https://pubmed.ncbi.nlm.nih.gov/9010100/)
- Kim JE, Huh J, Cho K, Kim CW (1998). Pathologic characteristics of primary cutaneous T-cell lymphoma in Korea. *J Korean Med Sci*, 13:31–38. [PMID:9539316](https://pubmed.ncbi.nlm.nih.gov/9539316/)
- Kumagawa M, Suzumiya J, Ohshima K *et al.* (2001). Splenic lymphoproliferative disorders in human T lymphotropic virus type-I endemic area of Japan: clinicopathological, immunohistochemical and genetic analysis of 27 cases. *Leuk Lymphoma*, 41:593–605.[doi:10.3109/10428190109060350](https://doi.org/10.3109/10428190109060350) [PMID:11378577](https://pubmed.ncbi.nlm.nih.gov/11378577/)
- Li H, Loehrer PJ Sr, Hisada M *et al.* (2004). Absence of human T-cell lymphotropic virus type I and human foamy virus in thymoma. *Br J Cancer*, 90:2181–2185. [PMID:15150553](https://pubmed.ncbi.nlm.nih.gov/15150553/)
- Marin O, Hasui K, Remondegui C *et al.* (2002). Adult T-cell leukemia/lymphoma in Jujuy, north-west Argentina. *Pathol Int*, 52:348–357.[doi:10.1046/j.1440-1827.2002.01356.x](https://doi.org/10.1046/j.1440-1827.2002.01356.x) [PMID:12100517](https://pubmed.ncbi.nlm.nih.gov/12100517/)
- Matsumoto S, Yamasaki K, Tsuji K, Shirahama S (2008). Human T lymphotropic virus type 1 infection and gastric cancer development in Japan. *J Infect Dis*, 198:10–15.[doi:10.1086/588733](https://doi.org/10.1086/588733) [PMID:18544011](https://pubmed.ncbi.nlm.nih.gov/18544011/)
- Murphy EL, Hanchard B, Figueroa JP *et al.* (1989). Modelling the risk of adult T-cell leukemia/lymphoma in persons infected with human T-lymphotropic virus type I. *Int J Cancer*, 43:250–253.[doi:10.1002/ijc.2910430214](https://doi.org/10.1002/ijc.2910430214) [PMID:2917802](https://pubmed.ncbi.nlm.nih.gov/2917802/)
- Nakamura S, Tsuneyoshi M (1998). Primary gastric T-cell lymphoma with and without human T-lymphotropic virus type 1. *Cancer*, 82:998–999.[doi:10.1002/\(SICI\)1097-0142\(19980301\)82:5<998::AID-CNCR35>3.0.CO;2-U](https://doi.org/10.1002/(SICI)1097-0142(19980301)82:5<998::AID-CNCR35>3.0.CO;2-U) [PMID:9486597](https://pubmed.ncbi.nlm.nih.gov/9486597/)
- Okayama A, Stuver S, Matsuoka M *et al.* (2004). Role of HTLV-1 proviral DNA load and clonality in the development of adult T-cell leukemia/lymphoma in asymptomatic carriers. *Int J Cancer*, 110:621–625.[doi:10.1002/ijc.20144](https://doi.org/10.1002/ijc.20144) [PMID:15122598](https://pubmed.ncbi.nlm.nih.gov/15122598/)
- Plumelle Y, Gonin C, Edouard A *et al.* (1997). Effect of *Strongyloides stercoralis* infection and eosinophilia on age at onset and prognosis of adult T-cell leukemia. *Am J Clin Pathol*, 107:81–87. [PMID:8980372](https://pubmed.ncbi.nlm.nih.gov/8980372/)
- Pombo De Oliveira MS, Loureiro P, Bittencourt A *et al.*; The Brazilian ATLL Study Group (1999). Geographic diversity of adult t-cell leukemia/lymphoma in Brazil. *Int J Cancer*, 83:291–298.[doi:10.1002/\(SICI\)1097-0215\(19991029\)83:3<291::AID-IJC1>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0215(19991029)83:3<291::AID-IJC1>3.0.CO;2-P) [PMID:10495418](https://pubmed.ncbi.nlm.nih.gov/10495418/)
- Sakata H, Iwakiri R, Koyama T *et al.* (2001). Human T-cell lymphotropic virus-associated primary gastric lymphoma. *Dig Dis Sci*, 46:1381–1386.[doi:10.1023/A:1010667232366](https://doi.org/10.1023/A:1010667232366) [PMID:11478487](https://pubmed.ncbi.nlm.nih.gov/11478487/)
- Satoh M, Toma H, Sugahara K *et al.* (2002). Involvement of IL-2/IL-2R system activation by parasite antigen in polyclonal expansion of CD4(+)25(+) HTLV-1-infected T-cells in human carriers of both HTLV-1 and *S. stercoralis*. *Oncogene*, 21:2466–2475.[doi:10.1038/sj.onc.1205329](https://doi.org/10.1038/sj.onc.1205329) [PMID:11971181](https://pubmed.ncbi.nlm.nih.gov/11971181/)
- Shimada-Hiratsuka M, Fukayama M, Hayashi Y *et al.* (1997). Primary gastric T-cell lymphoma with and without human T-lymphotropic virus type 1. *Cancer*, 80:292–303.[doi:10.1002/\(SICI\)1097-0142\(19970715\)80:2<292::AID-CNCR18>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0142(19970715)80:2<292::AID-CNCR18>3.0.CO;2-P) [PMID:9217043](https://pubmed.ncbi.nlm.nih.gov/9217043/)
- Tokudome S, Maeda Y, Fukada K *et al.* (1991). Follow-up of asymptomatic HTLV-I carriers among blood donors in Kyushu, Japan. *Cancer Causes Control*, 2:75–78.[doi:10.1007/BF00053124](https://doi.org/10.1007/BF00053124) [PMID:1873439](https://pubmed.ncbi.nlm.nih.gov/1873439/)
- Tsukasaki K, Imaizumi Y, Tawara M *et al.* (1999). Diversity of leukaemic cell morphology in ATL correlates with prognostic factors, aberrant immunophenotype and defective HTLV-1 genotype. *Br J Haematol*, 105:369–375.[doi:10.1111/j.1365-2141.1999.01323.x](https://doi.org/10.1111/j.1365-2141.1999.01323.x) [PMID:10233406](https://pubmed.ncbi.nlm.nih.gov/10233406/)
- Wood GS, Salvekar A, Schaffer J *et al.* (1996). Evidence against a role for human T-cell lymphotropic virus type I (HTLV-I) in the pathogenesis of American cutaneous T-cell lymphoma. *J Invest Dermatol*, 107:301–307.[doi:10.1111/1523-1747.ep12363010](https://doi.org/10.1111/1523-1747.ep12363010) [PMID:8751960](https://pubmed.ncbi.nlm.nih.gov/8751960/)
- Wood GS, Schaffer JM, Boni R *et al.* (1997). No evidence of HTLV-I proviral integration in lymphoproliferative disorders associated with cutaneous T-cell lymphoma. *Am J Pathol*, 150:667–673. [PMID:9033279](https://pubmed.ncbi.nlm.nih.gov/9033279/)