

**Table 2.26. Case-control studies of second-hand tobacco smoke and childhood brain and central nervous system cancers**

Reference, study location and period	Organ site (ICD code)	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	(Cases/Controls) OR (95% CI)* * $P < 0.05$	Adjustment for potential confounders	Comments
Schüz <i>et al.</i> (1999) Germany	Central nervous system (CNS)	2358 cases; 1992–1997; < 15 yrs; German Cancer Registry  Residence 15 km within German nuclear installations using cases born after 1975; < 15yrs; diagnosed 1980–1994	2588 population based control matched on age, sex and sampled from district where case lived at date of diagnoses	Structured questionnaire  Telephone Interview	<i>Maternal smoking during pregnancy (cigarettes per day)</i> No (reference) 1–10 11–20 > 20  <i>Paternal smoking before pregnancy (cigarettes per day)</i> No (reference) 1–10 11–20 > 20	CNS ( $n = 399$ )  (320/2062) 1.0 (55/426) 0.8 (0.6–1.1) (17/72) 1.6 (0.9–2.8) (2/11) 0.8 (0.2–3.9)  (195/1336) 1.0 (37/293) 0.8 (0.5–1.2) (112/647) 1.1 (0.8–1.4) (41/264) 1.0 (0.7–1.4)		

**Table 2.26. Case-control studies of second-hand tobacco smoke and childhood brain and central nervous system cancers**

Reference, study location and period	Organ site (ICD code)	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	(Cases/Controls) OR (95% CI)* * $P < 0.05$	Adjustment for potential confounders	Comments
Sorahan <i>et al.</i> (2001) UK	Central nervous system	555 children resident in Yorkshire, West Midlands, and North Western Regional Health Authority areas; < 15 yrs; 1980–1983; Not living with biologic mother excluded;	555 controls matched for sex and date of birth; Practice lists of case GPs	Micro-filmed interview	<i>Fathers before pregnancy (cigarettes per day)</i> Lifelong non-smoker (ref.) < 10 10–19 20–29 30–39 ≥ 40 <i>P</i> -value for trend (2-tailed) <i>Maternal smoking habit during 5<sup>th</sup> mo of pregnancy (cigarettes per day)</i> Lifelong non-smoker (ref.) < 10 10–19 20–29 ≥ 30 <i>P</i> -value for trend (2-tailed)	(cases/controls = 78)  (29/23) 1.0 (2/4) 0.51 (0.09–2.97) (12/9) 1.25 (0.42–3.72) (17/20) 0.52 (0.21–1.33) (1/5) 0.15 (0.01–1.54) (5/4) 0.64 (0.08–4.79) 0.67 (cases/controls = 76)  (32/40) 1.0 (11/2) 6.56 (1.36–31.73)* (16/16) 1.28 (0.55–3.03) (13/14) 1.30 (0.52–3.22) (0/0) 0 0.71	Socioeconomic grouping, ethnic origin, parental age of child at diagnosis	

**Table 2.26. Case-control studies of second-hand tobacco smoke and childhood brain and central nervous system cancers**

Reference, study location and period	Organ site (ICD code)	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	(Cases/Controls) OR (95% CI)* * $P < 0.05$	Adjustment for potential confounders	Comments
Filippini <i>et al.</i> (2002) (SEARCH) International	Brain (ICD-9 191)  Astroglial (ICD-O-2 9 380, 9 382, 9 384, 9 400, 9 401, 9 410, 9 411, 9 420, 9 421, 9 424, 9 440, 9 442,  Primitive neuroectodermal tumour. (ICD-O-2 9 391-2, 9 450, 9 505	1218 cases (< 0-19 yrs) 9 centres; Surveillance of Environmental Aspects Related to Cancer in Humans (SEARCH); 14 yrs in Europe and Australia; 19 yrs in Israel and USA; population based cancer registries; 1980s-1990s	2223 controls (see above)	In-person and phone interviews	<i>Maternal smoking during pregnancy</i>	AT	Age, sex, centre (years vary according to study centre)	
					≤ 2 500 cigarettes	(63/226) 1.0 (0.8-1.4)		
					> 2 500 cigarettes	(48/194) 0.9 (0.6-1.2)		
					≤ 2 500 cigarettes	(24/226) 0.9 (0.6-1.5)		
					> 2 500 cigarettes	(17/194) 0.8 (0.4-1.3)		
					≤ 2 500 cigarettes	(18/226) 1.2 (0.7-2.1)		
					> 2 500 cigarettes	(13/194) 1.1 (0.6-2.0)		
					≤ 2 500 cigarettes	(11/226) 0.6 (0.3-1.1)		
					> 2 500 cigarettes	(12/194) 0.8 (0.4-1.4)		
					≤ 2 500 cigarettes	(117/226) 1.0 (0.8-1.2)		
					> 2 500 cigarettes	(90/194) 0.8 (0.6-1.1)		
					<i>Maternal smoking during pregnancy (years)</i> (No/Yes)			
					0-1	(32/36) 1.3 (0.7-2.3)		
0-4	(80/137) 0.9 (0.6-1.2)							
5-9	(63/117) 1.1 (0.8-1.5)							
10-14	(35/102) 0.7 (0.4-1.0)							
15-19	(32/65) 1.0 (0.6-1.7)							

**Table 2.26. Case-control studies of second-hand tobacco smoke and childhood brain and central nervous system cancers**

Reference, study location and period	Organ site (ICD code)	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	(Cases/Controls) OR (95% CI)* * $P < 0.05$	Adjustment for potential confounders	Comments
Pang <i>et al.</i> (2003) UK	Central nervous system	3838 children with cancer; Hospital based;; < 15 yrs; 1991–1994 in Scotland; 1992–1994 in England and Wales;	7629 control children; matched for sex, date of birth and geographical area of residence at diagnosis; Randomly selected from Family Health Services Authorities (FHSA)	Face-to-face structured interviews; Computerized self-administered questionnaires	<i>Paternal cigarette smoking preconception (cigarettes per day)</i>  Lifelong smoker (ref) 1–19 20+ <i>P</i> for trend <i>Maternal cigarette smoking during pregnancy (cigarettes per day)</i> Nonsmoker (ref) 1–19 20+ <i>P</i> for trend <i>Primitive neuroectodermal tumours</i> Father preconception Mother preconception Mother pregnancy	(cases = 635; controls = 6 987)  (271/3082) 1.00 (101/1003) 1.08 (0.85-1.38) (138/1440) 1.03 (0.82-1.28) 0.706 (cases = 684; controls = 7 577)  (538/5743) 1.00 (111/1306) 0.86 (0.68-1.08) (35/528) 0.62 (0.42–0.93)* 0.011 (Cases) OR (95%CI)  (150) 0.90 (0.63–1.30) (161) 0.96 (0.65–1.42)  (161) 0.55* (0.34–0.88)	Deprivation and parental age at birth of index child	

**Table 2.26. Case-control studies of second-hand tobacco smoke and childhood brain and central nervous system cancers**

Reference, study location and period	Organ site (ICD code)	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	(Cases/Controls) OR (95% CI)* * $P < 0.05$	Adjustment for potential confounders	Comments
Cordier <i>et al.</i> (2004)  7 countries SEARCH International	Brain & Cranial nerves	1218 cases; 1976–1994 (65% between 1985–1989); 9 centres; 14 yrs in Europe and Australia; 19 yrs in Israel and USA;	2223 randomly selected controls from general population were either pair matched (6 centres) or frequency matched (3 centres) to cases by sex and birth year (or by age at one centre)	In-person Interviews at Patients home with biologic mother (some telephone interviews with fathers); occupational exposures to PAHs estimated using job exposure matrix	<u>Maternal exposure during pregnancy from smoking but not occupation</u>		Study centre, child's age, sex and maternal smoking during pregnancy	
					All tumours	(152/305) 0.9 (0.7–1.2)		
					Astroglial	(82/305) 1.0 (0.8–1.3)		
					Primitive neuroectodermal	(31/305) 0.8 (0.5–1.2)		
					Other glial	(22/305) 1.1 (0.7–1.7)		
					<u>from both smoking and occupation</u>			
					All tumours	(13/22) 1.2 (0.6–2.5)		
					Astroglial	(7/22) 1.3 (0.5–3.0)		
					Primitive neuroectodermal	(3/22) 1.2 (0.4–4.2)		
					Other glial	(2/22) 1.5 (0.3–6.6)		
					<u>Paternal preconceptional exposure from smoking but not occupation</u>			
					All tumours	(273/590) 1.1 (0.9–1.3)		
					Astroglial	(155/590) 1.4 (1.1–1.9)		
					Primitive neuroectodermal	(53/590) 0.8 (0.6–1.2)		
Other glial	(32/590) 1.1 (0.6–1.7)							
<u>from both smoking and occupation</u>								
All tumours	(272/464) 1.4 (1.1–1.7)							
Astroglial	(142/464) 1.6 (1.2–2.1)							
Primitive neuroectodermal	(54/464) 1.0 (0.7–1.5)							
Other glial	(34/464) 1.3 (0.8–2.1)							

**Table 2.26. Case-control studies of second-hand tobacco smoke and childhood brain and central nervous system cancers**

Reference, study location and period	Organ site (ICD code)	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	(Cases/Controls) OR (95% CI)* * $P < 0.05$	Adjustment for potential confounders	Comments
Plichart <i>et al.</i> (2008) France	Brain & CNS (ICD-O-3 9 470/3, 9 471/3, 9 473/3, 9 474/3, 9 480/3, 9 508/3, 9 390/3–9393/3, 9 380/3, 9 400/3, 9 420/3, 9 424/3, 9 424/3, 9 440/3, 9 380/3, 9 382/3, 9 430/3, 9 450/3, 9 451/3)	209 incident cases; 2003–2004; Hospital based; < 15 yrs; Residing in mainland France at time of diagnosis of primary malignant CNS tumour;	1681 population based controls matched on age and sex of child	Standardized telephone interview of the biological mothers French national Registry of Childhood Haematopoietic Malignancies and French national Registry of Childhood Solid Tumours (NRCST)	<i>Maternal smoking (pregnancy) (cigarettes per day)</i> No/Yes < 10 ≥ 10 <i>P</i> for trend <i>Paternal smoking (year before birth) (cigarettes per day)</i> No/Yes < 10 ≥ 10 <i>P</i> for trend	CNS  (44/325) 1.1 (0.8–1.6) (29/220) 1.1 (0.7–1.6) (13/99) 1.1 (0.6–2.0) NS  (103/746) 1.2 (0.9–1.6) (52/426) 1.1 (0.8–1.6) (51/310) 1.4 (1.0–2.1) 0.04 (see subtypes below)	Age and sex	
Plichart <i>et al.</i> (2008) Continued	<b><i>Maternal smoking (pregnancy)</i></b>	<b>PNET</b>	<b>Ependymomas</b>	<b>Astrocytomas</b>	<b>Other gliomas</b>			
	No/Yes	(21/325) 1.1 (0.7–1.8)	(6/325) 1.0 (0.4–2.4)	(6/325) 1.3 (0.5–3.2)	(10/325) 1.2 (0.6–2.4)			
	< 10 CPD	(16/220) 1.2 (0.7–2.1)	(4/220) 0.9 (0.3–2.7)	(4/220) 1.3 (0.4–3.8)	(5/220) 0.9 (0.3–2.3)			
	≥ 10 CPD	(5/99) 0.9 (0.3–2.2)	(1/99) 0.6 (0.1–4.3)	(2/99) 1.4 (0.3–5.9)	(4/99) 1.6 (0.5–4.5)			
	<i>P</i> for trend	NS	NS	NS	NS			
	<b><i>Paternal smoking (year before birth)</i></b>							

**Table 2.26. Case-control studies of second-hand tobacco smoke and childhood brain and central nervous system cancers**

Reference, study location and period	Organ site (ICD code)	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	(Cases/Controls) OR (95% CI)* * $P < 0.05$	Adjustment for potential confounders	Comments
	<b>Maternal smoking (pregnancy)</b>	<b>PNET</b>	<b>Ependymomas</b>	<b>Astrocytomas</b>	<b>Other gliomas</b>			
	No/Yes	(49/746) 1.2 (0.8–1.8)	(18/746) 1.6 (0.8–3.3)	(18/746) 3.1 (1.3–7.6)	(17/746) 0.8 (0.4–1.4)			
	< 20 CPD	(27/426) 1.2 (0.7–1.9)	(7/426) 1.0 (0.4–2.6)	(10/426) 3.1 (1.2–8.4)	(8/426) 0.7 (0.3–1.5)			
	≥ 20 CPD	(22/310) 1.2 (0.7–2.1)	(11/310) 2.6 (1.2–5.9)	(8/310) 3.2 (1.2–9.1)	(9/310) 0.9 (0.4–2.0)			
	<i>P</i> for trend	NS	0.02	0.02	NS			