

NICKEL AND NICKEL COMPOUNDS (Group 1*)

A. Evidence for carcinogenicity to humans (*sufficient*)

Early epidemiological studies of populations of workers in nickel refineries in different countries clearly demonstrate excess incidences of cancers of the nasal cavity and lung and, possibly, excesses of cancer of the larynx. Although the carcinogen(s) could not be specified, the cancer hazards seemed to be associated primarily with the early stage of nickel refining. Nickel carbonyl was considered unlikely to be involved, while nickel subsulphide and nickel oxide emerged as the strongest candidates¹.

Later reports from Canada and the USA confirmed the increased risks for lung and sinonasal cancers carried by exposure during nickel refining operations, where the primary exposure was to nickel sulphides (including subsulphide) and nickel oxides²⁻⁴. The early studies of nickel refinery workers in Wales (UK) and Norway were extended and updated. Among workers in South Wales, the elevated risks for cancers of the lung and nasal sinuses persisted until 1930⁵. For both lung cancer (137 cases before 1925) and nasal cancer (56 cases before 1925), the increased risk was significantly associated with employment in calcining and at furnaces and with copper sulphate and nickel sulphate production^{6,7}. In Norway, the highest incidence rates for cancers of the respiratory organs occurred among workers employed in roasting, smelting and electrolysis departments. The increased incidence of nasal cancers (21 cases; relative risk, 26.3) exhibited a very steep decrease with more recent

*This evaluation applies to the group of chemicals as a whole and not necessarily to all individual chemicals within the group (see also Methods, p. 38). After the meeting of the Working Group, the Secretariat became aware of epidemiological and experimental studies in progress on the carcinogenicity of nickel and nickel compounds.

year of first employment; the incidence of lung cancer (82 cases; relative risk, 3.7) gave no indication of a consistent decrease during the period 1916-1959. A slight, statistically nonsignificant excess of laryngeal cancer (5 cases observed, 2.4 expected) was also reported^{8,9}.

Reports of an increased occurrence of lung cancer among nickel smelting workers have also come from New Caledonia, Slovakia and the USSR¹⁰⁻¹⁶.

There have been three case reports of cancers of the respiratory tract in workers who were involved in nickel plating and grinding operations¹.

Three investigations that examined the possible cancer risk associated with exposure to nickel and nickel compounds in nickel alloy plants showed no significant increase in mortality from cancer¹⁷⁻¹⁹. In one of these, excess mortality from lung cancer was noted in maintenance workers; however, it was unclear whether the risk was directly associated with nickel exposures¹⁸. Workers at a gaseous diffusion plant who were exposed to high-purity metallic nickel powder did not exhibit any increase in mortality from respiratory-tract cancers^{20,21}. An incidence study at a hydrometallurgical nickel refining plant in Canada did not indicate an increased risk of cancer. Exposure was to metallic nickel and nickel concentrate dust²².

Other investigations have addressed more complex and mixed exposure conditions and thus provide little evidence to evaluate the specific role of nickel and nickel compounds²³⁻³⁰.

The association of specific types of cancer with nickel exposure has also been examined by means of case-control investigations. One study of cancer of the larynx supported an association with nickel exposure³¹, but another did not³². Studies of sinonasal cancer and lung cancer yielded contradictory results; all suffered from inadequate description of the exposure to nickel³³⁻³⁶. In one of these³⁵, the risk was high in welders with nickel exposure (relative risk, 3.3, 95% confidence interval, 1.2-9.2); however, exposure to nickel compounds was so highly correlated with the presence of chromium that the observed exposure to nickel could have reflected a confounding effect of chromium (see p. 165). A study at an aircraft-engine factory showed no association between lung cancer deaths and exposure to nickel oxides, sulphate, chloride or alloys³⁷.

It is still not possible to state with certainty which specific nickel compounds are human carcinogens, and which are not. A large amount of evidence has accrued that nickel refining carries a carcinogenic risk to workers. The risk is particularly high in those exposed during certain processes, mainly entailing exposure to nickel (sub)sulphides and oxides. The lung and nasal sinuses are the most clearly established target organs.

B. Evidence for carcinogenicity to animals (*sufficient*)

Nickel subsulphide produced malignant tumours in rats after its inhalation¹ or intramuscular^{1,38,39}, intrarenal^{40,41}, intratesticular⁴² or intraocular⁴³ administration and after its insertion into heterotransplanted tracheas⁴⁴; it also produced local sarcomas in mice and rabbits after intramuscular administration^{1,45-47}. Nickel powder, nickel oxide, hydroxide and carbonate, nickelocene and nickel-iron sulphide matte produced local sarcomas in mice, rats, hamsters and rabbits when given intramuscularly^{1,38,48}. Intravenous

administration of nickel carbonyl increased the incidences of various tumours in rats¹, and inhalation of nickel carbonyl produced a low incidence of lung tumours in rats¹. Nickelous acetate administered intraperitoneally to mice produced an excess of lung adenomas and carcinomas⁴⁹. Nickel sulphide produced renal tumours in rats when injected intrarenally⁵⁰.

With few exceptions, the nickel compounds tested produced sarcomas and/or carcinomas at the tissue sites where they were deposited. Bioavailability and persistence in the tissues appear to be important in nickel carcinogenesis.

C. Other relevant data

Studies of the uptake, content and release of nickel in nasal mucosa indicate that workers exposed to water-insoluble nickel salts (e.g., roasting and smelting workers) retain more nickel than those exposed to soluble compounds (e.g., electrolysis workers). Nickel accumulated during active work is retained in the mucous membrane for years after retirement⁵¹⁻⁵³.

Workers exposed to nickel in one refinery had slight excesses of chromosomal aberrations (mainly gaps) in their peripheral lymphocytes, but no increase in the incidence of sister chromatid exchanges was seen⁵⁴.

Nickel compounds did not induce dominant lethal mutations in mice. Soluble nickel compounds caused DNA strand breaks and cross-links in rats treated *in vivo*, and particles of crystalline nickel sulphide bound to DNA in Chinese hamster cells *in vitro*. Nickel compounds were weakly active in inducing chromosomal aberrations and sister chromatid exchanges in human lymphocytes and rodent cells *in vitro*. They induced transformation in several rodent cell systems *in vitro*. Particles of crystalline nickel sulphides induced mutation in a protozoan. In general, negative results were obtained in bacterial mutation assays; nickel compounds induced prophage in bacteria. Insoluble nickel compounds bound to isolated DNA⁵⁴.

References

- ¹IARC Monographs, 11, 75-112, 1976
- ²Chovil, A., Sutherland, R.B. & Halliday, M. (1981) Respiratory cancer in a cohort of nickel sinter plant workers. *Br. J. ind. Med.*, 38, 327-333
- ³Roberts, R.S., Julian, J.A., Muir, D.C.F. & Shannon, H.S. (1984) *Cancer mortality associated with the high-temperature oxidation of nickel subsulfide*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 23-35
- ⁴Enterline, P.E. & Marsh, G.M. (1982) Mortality among workers in a nickel refinery and alloy manufacturing plant in West Virginia. *J. natl Cancer Inst.*, 68, 925-933
- ⁵Doll, R., Mathews, J.D. & Morgan, L.G. (1977) Cancers of the lung and nasal sinuses in nickel workers: a reassessment of the period of risk. *Br. J. ind. Med.*, 34, 102-105

- ⁶Peto, J., Cuckle, H., Doll, R., Hermon, C. & Morgan, L.G. (1984) *Respiratory cancer mortality of Welsh nickel refinery workers*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 37-46
- ⁷Kaldor, J., Peto, J., Easton, D., Doll, R., Hermon, C. & Morgan, L. (1986) Models for respiratory cancer in nickel refinery workers. *J. natl. Cancer Inst.*, 77, 841-848
- ⁸Pedersen, E., Andersen, A. & Høgetveit, A. (1978) Second study of the incidence and mortality of cancer of respiratory organs among workers at a nickel refinery (Abstract). *Ann. clin. Lab. Sci.*, 8, 503-504
- ⁹Magnus, K., Andersen, A. & Høgetveit, A.C. (1982) Cancer of respiratory organs among workers at a nickel refinery in Norway. Second report. *Int. J. Cancer*, 30, 681-685
- ¹⁰Lessard, R., Reed, D., Maheux, B. & Lambert, J. (1978) Lung cancer in New Caledonia, a nickel smelting island. *J. occup. Med.*, 20, 815-817
- ¹¹Olejár, Š., Olejárová, E. & Vrábel, K. (1982) Neoplasia of the lungs in the workers of the nickel smelting plant (Czech.). *Pracov. Léč.*, 34, 280-282
- ¹²Langer, A.M., Rohl, A.N., Selikoff, I.J., Harlow, G.E. & Prinz, M. (1980) Asbestos as a cofactor in carcinogenesis among nickel-processing workers. *Science*, 209, 420-422
- ¹³Meininger, J., Raffinot, P. & Trolly, G. (1982) Cancer in nickel-processing workers in New Caledonia. *Science*, 215, 424-425
- ¹⁴Langer, A.M., Rohl, A.N., Fischbein, A. & Selikoff, I.J. (1982) Cancer in nickel-processing workers in New Caledonia. *Science*, 215, 425-426
- ¹⁵Saknyn, A.V. & Shabynina, N.K. (1970) Some statistical data on the carcinogenous hazards for workers engaged in the production of nickel from oxidized ores (Russ.). *Gig. Tr. prof. Zabol.*, 14, 10-13
- ¹⁶Saknyn, A.V. & Shabynina, N.K. (1973) Epidemiology of malignant new growths at nickel smelters (Russ.). *Gig. Tr. prof. Zabol.*, 17, 25-29
- ¹⁷Cox, J.E., Doll, R., Scott, W.A. & Smith, S. (1981) Mortality of nickel workers: experience of men working with metallic nickel. *Br. J. ind. Med.*, 38, 235-239
- ¹⁸Redmond, C.K. (1984) *Site-specific cancer mortality among workers involved in the production of high nickel alloys*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 73-86
- ¹⁹Cornell, R.G. (1984) *Mortality patterns among stainless-steel workers*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 65-71
- ²⁰Goldbold, J.H., Jr & Tompkins, E.A. (1979) A long-term mortality study of workers occupationally exposed to metallic nickel at the Oak Ridge gaseous diffusion plant. *J. occup. Med.*, 21, 799-806
- ²¹Cragle, D.L., Hollis, D.R., Newport, T.H. & Shy, C.M. (1984) *A retrospective cohort mortality study among workers occupationally exposed to metallic nickel powder at the Oak Ridge gaseous diffusion plant*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 57-63
- ²²Egedahl, R. & Rice, E. (1984) *Cancer incidence at a hydrometallurgical nickel refinery*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 47-55

- ²³Shannon, H.S., Julian, J.A. & Roberts, R.S. (1984) A mortality study of 11,500 nickel workers. *J. natl Cancer Inst.*, *74*, 1251-1258
- ²⁴Cornell, R.G. & Landis, J.R. (1984) *Mortality patterns among nickel/chromium alloy foundry workers*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 87-93
- ²⁵Becker, N., Claude, J. & Frentzel-Beyme, R. (1985) Cancer risk of arc welders exposed to fumes containing chromium and nickel. *Scand. J. Work Environ. Health*, *11*, 75-82
- ²⁶Polednak, A.P. (1981) Mortality among welders, including a group exposed to nickel oxides. *Arch. environ. Health*, *36*, 235-242
- ²⁷Newhouse, M.L., Oakes, D. & Woolley, A.J. (1985) Mortality of welders and other craftsmen at a shipyard in NE England. *Br. J. ind. Med.*, *42*, 406-410
- ²⁸Sorahan, T. & Waterhouse, J.A.H. (1983) Mortality study of nickel-cadmium battery workers by the method of regression models in life tables. *Br. J. ind. Med.*, *40*, 293-300
- ²⁹Sorahan, T. & Waterhouse, J.A.H. (1985) Cancer of prostate among nickel-cadmium battery workers. *Lancet*, *i*, 459
- ³⁰Andersson, K., Elinder, C.G., Hogstedt, C., Kjellström, T. & Spång, G. (1984) Mortality among cadmium and nickel-exposed workers in a Swedish battery factory. *Toxicol. environ. Chem.*, *9*, 53-62
- ³¹Olsen, J. & Sabroe, S. (1984) Occupational causes of laryngeal cancer. *J. Epidemiol. Commun. Health*, *38*, 117-121
- ³²Burch, J.D., Howe, G.R., Miller, A.B. & Semenciw, R. (1981) Tobacco, alcohol, asbestos, and nickel in the etiology of cancer of the larynx: a case-control study. *J. natl Cancer Inst.*, *67*, 1219-1224
- ³³Roush, G.C., Meigs, J.W., Kelly, J., Flannery, J.T. & Burdo, H. (1980) Sinonasal cancer and occupation: a case-control study. *Am. J. Epidemiol.*, *111*, 183-193
- ³⁴Acheson, E.D., Cowdell, R.H. & Rang, E.H. (1981) Nasal cancer in England and Wales: an occupational survey. *Br. J. ind. Med.*, *38*, 218-224
- ³⁵Gérin, M., Siemiatycki, J., Richardson, L., Pellerin, J., Lakhani, R. & Dewar, R. (1984) *Nickel and cancer associations from a multicancer occupation exposure case-referent study: preliminary findings*. In: Sunderman, F.W., Jr, ed., *Nickel in the Human Environment (IARC Scientific Publications No. 53)*, Lyon, International Agency for Research on Cancer, pp. 105-115
- ³⁶Hernberg, S., Collan, Y., Degerth, R., Englund, A., Engzell, U., Kuosma, E., Mutanen, P., Nordlinder, H., Sand Hansen, H., Schultz-Larsen, K., Sjøgaard, H. & Westerholm, P. (1983) Nasal cancer and occupational exposures. Preliminary report of a joint Nordic case-referent study. *Scand. J. Work Environ. Health*, *9*, 208-213
- ³⁷Bernacki, E.J., Parsons, G.E. & Sunderman, F.W., Jr (1978) Investigation of exposure to nickel and lung cancer mortality. Case control study at aircraft engine factory. *Ann. clin. Lab. Sci.*, *8*, 190-194
- ³⁸Sunderman, F.W., Jr & Maenza, R.M. (1976) Comparisons of carcinogenicities of nickel compounds in rats. *Res. Commun. chem. Pathol. Pharmacol.*, *14*, 319-330
- ³⁹Yamashiro, S., Gilman, J.P.W., Hulland, T.J. & Abandowitz, H.M. (1980) Nickel sulphide-induced rhabdomyosarcomata in rats. *Acta pathol. jpn.*, *30*, 9-22
- ⁴⁰Jasmin, G. & Riopelle, J.L. (1976) Renal carcinomas and erythrocytosis in rats following intrarenal injection of nickel subsulfide. *Lab. Invest.*, *35*, 71-78

- ⁴¹Sunderman, F.W., Jr, Maenza, R.M., Hopfer, S.M., Mitchell, J.M., Allpass, P.R. & Damjanov, I. (1979) Induction of renal cancers in rats by intrarenal injection of nickel subsulfide. *J. environ. Pathol. Toxicol.*, 21, 1511-1527
- ⁴²Damjanov, I., Sunderman, F.W., Jr, Mitchell, J.M. & Allpass, P.R. (1978) Induction of testicular sarcomas in Fischer rats by intratesticular injection of nickel subsulfide. *Cancer Res.*, 38, 268-276
- ⁴³Albert, D.M., Gonder, J.R., Papale, J., Craft, J.L., Dohlman, H.G., Reid, M.C. & Sunderman, F.W., Jr (1980) *Induction of ocular neoplasms in Fischer rats by intraocular injection of nickel subsulfide.* In: Brown, S.S. & Sunderman, F.W., Jr, eds, *Nickel Toxicology*, New York, Academic Press, pp. 55-58
- ⁴⁴Yarita, T. & Nettesheim, P. (1978) Carcinogenicity of nickel subsulfide for respiratory tract mucosa. *Cancer Res.*, 38, 3140-3145
- ⁴⁵Hildebrand, H.F. & Biserte, G. (1979) Cylindrical laminated bodies in nickel-subsulphide-induced rhabdomyosarcoma in rabbits. *Eur. J. Cell Biol.*, 19, 276-280
- ⁴⁶Hildebrand, H.F. & Biserte, G. (1979) Nickel sub-sulphide-induced leiomyosarcoma in rabbit white skeletal muscle. A light microscopical and ultrastructural study. *Cancer*, 43, 1358-1374
- ⁴⁷Hildebrand, H.F. & Tetaert, D. (1981) Ni₃S₂-induced leiomyosarcomas in rabbit skeletal muscle: analysis of the tumoral myosin and its significance in the retrodifferentiation concept. *Oncodev. Biol. Med.*, 2, 101-108
- ⁴⁸Kasprzak, K.S., Gabryel, P. & Jarczewska, K. (1983) Carcinogenicity of nickel(II)hydroxides and nickel(II)sulfate in Wistar rats and its relation to the *in vitro* dissolution rates. *Carcinogenesis*, 4, 275-279
- ⁴⁹Stoner, G.D., Shimkin, M.B., Troxell, M.C., Thompson, T.L. & Terry, L.S. (1976) Test for carcinogenicity of metallic compounds by the pulmonary tumor response in strain A mice. *Cancer Res.*, 36, 1744-1747
- ⁵⁰Sunderman, F.W., Jr, McCully, K.S. & Hopfer, S.M. (1984) Association between erythrocytosis and renal cancers in rats following intrarenal injection of nickel compounds. *Carcinogenesis*, 5, 1511-1517
- ⁵¹Torjussen, W. (1979) Nickel as carcinogenic factor in nasal carcinoma. *Acta otolaryngol., Suppl.* 360, 125
- ⁵²Barton, R.T. & Høgetveit, A.C. (1980) Nickel-related cancers of the respiratory tract. *Cancer*, 45, 3061-3064
- ⁵³Torjussen, W. (1985) Occupational nasal cancer caused by nickel and nickel compounds. *Rhinology*, 23, 101-105
- ⁵⁴IARC Monographs, Suppl. 6, 417-420, 1987