

2-NITRONAPHTHALENE

1. Chemical and Physical Data

1.1 Synonyms


Chem. Abstr. Services Reg. No.: 581-89-5

Chem. Abstr. Name: Naphthalene, 2-nitro-

IUPAC Systematic Name: 2-Nitronaphthalene

Synonym: β -Nitronaphthalene

1.2 Structural and molecular formulae and molecular weight



$C_{10}H_7NO_2$

Mol. wt: 173.2

1.3 Chemical and physical properties of the pure substance

(a) *Description:* Yellow needles or plates from ethanol (Weast, 1985)

(b) *Boiling-point:* 165°C at 15°C (Weast, 1985)

(c) *Melting-point:* 79°C (Weast, 1985)

(d) *Spectroscopy data:* Proton and nuclear magnetic resonance and mass spectral data have been reported (Lucchini & Wells, 1976; Kitching *et al.*, 1977; Schuetzle & Jensen, 1985).

(e) *Solubility:* Soluble in ethanol and diethyl ether (Weast, 1985)

1.4 Technical products and impurities

2-Nitronaphthalene is available for research purposes (Aldrich Chemical Co., 1988). It is also available at a purity of 99.7% as a reference material (Belliardo *et al.*, 1988).

2. Production, Use, Occurrence and Analysis

2.1 Production and use

(a) Production

2-Nitronaphthalene is a by-product (2–3%) from the commercial preparation of 1-nitronaphthalene by direct nitration of naphthalene (Verschueren, 1983). It can be produced by the Bucherer reaction, starting with 2-naphthalenol (2-naphthol), and by other indirect methods; however, the Bucherer method, which yields 2-naphthylamine (see IARC, 1987), is seldom used (Gaydos, 1981).

(b) Use

No evidence was found that 2-nitronaphthalene has been used for commercial applications.

2.2 Occurrence

(a) Engine exhaust

Emissions from a heavy-duty diesel engine contained 2-nitronaphthalene at concentrations of 0.94 mg/kg in particulates when the engine was run at 100% load and 1200 revolutions/min and 0.87 mg/kg in particulates when the same engine was run at 75% load and 1800 revolutions/min (Draper, 1986).

(b) Other occurrence

Toners for use in photocopy machines have been produced in quantity since the late 1950s and have seen widespread use. 'Long-flow' furnace black was first used in photocopy toners in 1967; its manufacture involved an oxidation whereby some nitration also occurred. Subsequent changes in the production technique reduced the total extractable nitropyrene content from an uncontrolled level of 5–100 mg/kg to below 0.3 mg/kg (Rosenkranz *et al.*, 1980; Sanders, 1981; Butler *et al.*, 1983), and toners produced from this carbon black since 1980 have not been found to contain detectable levels of mutagenicity or, hence, nitropyrenes (Rosenkranz *et al.*, 1980; Butler *et al.*, 1983). One sample of a formerly available commercial carbon black was reported to contain detectable concentrations of unspecified nitronaphthalene (Ramdahl & Urdal, 1982).

2-Nitronaphthalene was detected in urban ambient airborne particulates in St Louis, MO, USA (Ramdahl & Urdal, 1982; Ramdahl *et al.*, 1982). In a laboratory experiment, 2-nitronaphthalene was formed by a gas-phase naphthalene nitration reaction at room temperature and atmospheric pressure with a yield of about 8% (Pitts *et al.*, 1985). It was postulated that significant amounts of 2-nitronaphthalene result from the night-time gas-phase reaction of naphthalene with nitrogen pentoxide (Pitts, 1987). Approximately equal

amounts of 1- and 2-nitronaphthalene were found in day-time samples of ambient air in Torrance, CA, USA. 2-Nitronaphthalene concentrations were 1.1 ng/m³ of sample during night-time sampling and 2.9 ng/m³ of sample during day-time sampling (Arey *et al.*, 1987).

2.3 Analysis

See the monograph on 1-nitropyrene.

3. Biological Data Relevant to the Evaluation of Carcinogenic Risk to Humans

3.1 Carcinogenicity studies in animals¹

(a) Oral administration

Monkey: One female rhesus monkey (*Macaca mulatta*) that had been infected with a laboratory strain of *Plasmodium cynomolgi* and cured with chloroquine at least 12 weeks before the beginning of the experiment received a daily dose of 242 mg/kg bw 2-nitronaphthalene ['purified' but purity not given] administered orally in hard gelatin capsules on six days per week for 54 months. The monkey was then killed (terminal weight, 5.7 kg) and necropsied, three papillomas were found in the urinary bladder; no tumour was found in other organs (Conzelman *et al.*, 1970). [The Working Group noted that results were available for only one animal.]

(b) Bladder implantation

Mouse: Three groups of 41, 76 and 80 mice [strain and age unspecified] received a surgical implant of a cholesterol pellet containing 2-nitronaphthalene [dose and purity unspecified] into the urinary bladder (Bryan *et al.*, 1964). Three groups of 72, 82 and 140 controls received implants of cholesterol only. In a further group of 38 mice, a sham operation was performed which consisted of incising the bladder, momentarily inserting a pellet of pure cholesterol, removing the pellet, closing the bladder and suturing the abdomen. Animals were observed for up to 490 days, and only animals surviving longer than 175 days were evaluated. At the end of the experiment, the incidences of carcinomas and benign tumours of the bladder did not differ significantly between the various groups: 2/41, 2/76 and 7/80 in the treated groups; 4/72, 1/82 and 2/140 in the cholesterol controls and none in the sham-operated animals.

¹The Working Group was aware of studies in progress by single subcutaneous injection in mice (IARC, 1988).

3.2 Other relevant data

(a) Experimental systems

(i) Absorption, distribution, excretion and metabolism

2-Naphthylamine (see IARC, 1987) was detected in the urine of male Sprague-Dawley rats administered single intraperitoneal doses of 100 mg/kg bw 2-nitronaphthalene (Johnson & Cornish, 1978).

After male Wistar rats had been administered 2 mmol[350 mg]/kg bw 2-nitronaphthalene orally, the following metabolites (either free or conjugated) were excreted in the urine over the subsequent 32 h: 2-amino-1-naphthyl sulfate, 2-amino-1-naphthol and *N*-hydroxy-2-naphthylamine. All of these products were detected in the urine of rats treated similarly with 2-naphthylamine (Kadlubar *et al.*, 1981). [The Working Group noted that *N*-hydroxy-2-naphthylamine is tumorigenic to experimental animals. See, for example, Garner *et al.* (1984).] The urine of monkeys given oral doses of 2-nitronaphthalene was found to contain 2-amino-1-naphthyl sulfate and 2-acetamido-6-naphthyl glucuronide (Conzelman *et al.*, 1970).

As reported in an abstract, in male Sprague-Dawley rats treated orally with 0.5 mmol[87 mg]/kg bw 2-nitronaphthalene, covalent binding to haemoglobin was detected, although the level was lower than that found with 2-naphthylamine (Suzuki *et al.*, 1987).

Under anaerobic conditions, liver postmitochondrial supernatants and cytosol from male Fischer rats and female Swiss mice catalysed the stoichiometric conversion of 2-nitronaphthalene to 2-naphthylamine (Poirier & Weisburger, 1974).

Incubation of 2-nitronaphthalene with liver postmitochondrial supernatants from Wistar rats pretreated with sodium phenobarbital, 3-methylcholanthrene or Kanechlor-500 resulted in a time-dependent loss of substrate. The rate of metabolism was faster with homogenates from animals pretreated with 3-methylcholanthrene or Kanechlor-500 than with those from animals given phenobarbital (Ohe, 1985).

(ii) Toxic effects

A single intraperitoneal injection of 2-nitronaphthalene (100 mg/kg bw) to male Sprague-Dawley rats did not cause respiratory distress or lung or liver toxicity, in contrast to observations made with 1-nitronaphthalene (see monograph on 1-nitronaphthalene; Johnson *et al.*, 1984).

(iii) Genetic and related effects

The genetic and related effects of nitroarenes and of their metabolites have been reviewed (Rosenkranz & Mermelstein, 1983; Beland *et al.*, 1985; Rosenkranz & Mermelstein, 1985; Tokiwa & Ohnishi, 1986).

2-Nitronaphthalene preferentially inhibited the growth of DNA repair-deficient *Escherichia coli* (Rosenkranz & Poirier, 1979; DeFlora *et al.*, 1984) and *Bacillus subtilis* (at 100 µg/disc; Tokiwa *et al.*, 1987) and induced DNA damage in *Salmonella typhimurium* (lowest effective dose, 6 µg/ml; Nakamura *et al.*, 1987).

2-Nitronaphthalene was mutagenic to *S. typhimurium* strains TA1535, TA1537, TA1538, TA98 and TA100 (McCann *et al.*, 1975; Wang *et al.*, 1978; De Flora, 1979; Rosenkranz & Poirier, 1979; Scribner *et al.*, 1979; Simmon, 1979a; Wang *et al.*, 1980; De Flora, 1981; El-Bayoumy *et al.*, 1981; Ho *et al.*, 1981; Löfroth, 1981; McCoy *et al.*, 1981; De Flora *et al.*, 1984; Morotomi & Watanabe, 1984) but not to strain TA97 (Rosenkranz *et al.*, 1985).

This compound induced recombination in the yeast, *Saccharomyces cerevisiae* D3 (Simmon, 1979b).

In a host-mediated assay in mice, 2-nitronaphthalene (at 125–1300 mg/kg) induced mutation in *S. typhimurium* TA1530 and TA1538 but not recombination in *S. cerevisiae* D3 (Simmon *et al.*, 1979).

2-Nitronaphthalene did not induce unscheduled DNA synthesis in cultured rat or mouse hepatocytes (Mori *et al.*, 1987) but did induce morphological transformation of Syrian hamster embryo cells (Pienta, 1980).

(b) *Humans*

No data were available to the Working Group.

3.3 Epidemiological studies and case reports of carcinogenicity in humans

No data were available to the Working Group.

4. Summary of Data Reported and Evaluation

4.1 Exposure data

2-Nitronaphthalene has been detected in particulate emissions from diesel engines. It has also been found at low concentrations in ambient air.

4.2 Experimental data

2-Nitronaphthalene was tested for carcinogenicity by prolonged oral administration in one monkey, producing papillomas in the urinary bladder. Implantation of cholesterol pellets containing 2-nitronaphthalene into the bladder of mice did not increase the incidence of urinary bladder tumours.

4.3 Human data

No data were available to the Working Group.

4.4 Other relevant data

N-Hydroxy-2-naphthylamine (which has been shown to induce tumours in experimental animals) and 2-naphthylamine (which is causally associated with cancer in humans) have been detected as metabolites of 2-nitronaphthalene in the urine of rats.

2-Nitronaphthalene induced morphological transformation in cultured animal cells but did not induce DNA damage in cultured rodent hepatocytes. It induced DNA damage and mutation in bacteria and recombination in yeast.

4.5 Evaluation¹

There is *inadequate evidence* for the carcinogenicity in experimental animals of 2-nitronaphthalene.

No data were available from studies in humans on the carcinogenicity of 2-nitronaphthalene.

Overall evaluation

2-Nitronaphthalene *is not classifiable as to its carcinogenicity to humans (Group 3)*.

5. References

- Aldrich Chemical Co. (1988) *Aldrich Catalog/ Handbook of Fine Chemicals 1988-1989*, Milwaukee, WI, p. 1118
- Arey, J., Zielinska, B., Atkinson, R. & Winer, A.M. (1987) Polycyclic aromatic hydrocarbon and nitroarene concentrations in ambient air during a wintertime high-NO_x episode in the Los Angeles basin. *Atmos. Environ.*, *21*, 1437-1444
- Beland, F.A., Heflich, R.H., Howard, P.C. & Fu, P.P. (1985) The in vitro metabolic activation of nitro polycyclic aromatic hydrocarbons. In: Harvey, R.G., ed., *Polycyclic Hydrocarbons and Carcinogenesis (ACS Symposium Series No. 283)*, Washington DC, American Chemical Society, pp. 371-396
- Belliardo, J.J., Jacob, J. & Lindsey, A.S. (1988) *The Certification of the Purity of Seven Nitro-polycyclic Aromatic Compounds, CRM Nos 305, 306, 307, 308, 310, 311, 312, Reference Materials, BCR Information (EUR 11254 EN)*, Brussels, Commission of the European Communities, p. III
- Bryan, G.T., Brown, R.R. & Price, J.M. (1964) Mouse bladder carcinogenicity of certain tryptophan metabolites and other aromatic nitrogen compounds suspended in cholesterol. *Cancer Res.*, *24*, 596-602

¹For definitions of the italicized terms, see Preamble, pp. 25-28.

Summary table of genetic and related effects of 2-nitronaphthalene

Nonmammalian systems												Mammalian systems																																		
Proka-ryotes			Lower eukaryotes			Plants			Insects			<i>In vitro</i>						<i>In vivo</i>																												
						Animal cells						Human cells						Animals						Humans																						
D	G		D	R	G	A	D	G	C	R	G	C	A	D	G	S	M	C	A	T	I	D	G	S	M	C	A	T	I	D	G	S	M	C	DL	A	D	S	M	C	A					
+	+				+ ¹									-						+ ¹																										

A, aneuploidy; C, chromosomal aberrations; D, DNA damage; DL, dominant lethal mutation; G, gene mutation; I, inhibition of intercellular communication; M, micronuclei; R, mitotic recombination and gene conversion; S, sister chromatid exchange; T, cell transformation

In completing the table, the following symbols indicate the consensus of the Working Group with regard to the results for each endpoint:

- + considered to be positive for the specific endpoint and level of biological complexity
- +¹ considered to be positive, but only one valid study was available to the Working Group
- considered to be negative

- Butler, M.A., Evans, D.L., Giammarise, A.T., Kiriazides, D.K., Marsh, D., McCoy, E.C., Mermelstein, R., Murphy, C.B. & Rosenkranz, H.S. (1983) Application of *Salmonella* assay to carbon blacks and toners. In: Cooke, M. & Dennis, A.J., eds, *Polynuclear Aromatic Hydrocarbons, 7th International Symposium, Formation, Metabolism and Measurement*, Columbus, OH, Battelle, pp. 225-241
- Conzelman, G.M., Jr, Moulton, J.E. & Flanders, L.E., III (1970) Tumors in the urinary bladder of a monkey: induction with 2-nitronaphthalene. *Gann*, 61, 79-80
- DeFlora, S. (1979) Metabolic activation and deactivation of mutagens and carcinogens. *Ital. J. Biochem.*, 28, 81-103
- De Flora, S. (1981) Study of 106 organic and inorganic compounds in the *Salmonella*/microsome test. *Carcinogenesis*, 2, 283-298
- De Flora, S., Zanicchi, P., Camoirano, A., Bennicelli, C. & Badolati, G.S. (1984) Genotoxic activity and potency of 135 compounds in the Ames reversion test and in a bacterial DNA-repair test. *Mutat. Res.*, 133, 161-198
- Draper, W.M. (1986) Quantitation of nitro- and dinitropolycyclic aromatic hydrocarbons in diesel exhaust particulate matter. *Chemosphere*, 15, 437-447
- El-Bayoumy, K., Lavoie, E.J., Hecht, S.S., Fow, E.A. & Hoffmann, D. (1981) The influence of methyl substitution on the mutagenicity of nitronaphthalenes and nitrobiphenyls. *Mutat. Res.*, 81, 143-153
- Garner, R.C., Martin, C.N. & Clayson, D.B. (1984) Carcinogenic aromatic amines and related compounds. In: Searle, C.E., ed, *Chemical Carcinogens*, 2nd ed., Vol. 1 (*ACS Monograph 182*), Washington DC, American Chemical Society, pp. 175-276
- Gaydos, R.M. (1981) Naphthalene. In: Mark, H.F., Othmer, D.F., Overberger, C.G., Seaborg, G.T. & Grayson, M., eds, *Kirk-Othmer Encyclopedia of Chemical Technology*, 3rd ed., Vol. 15, New York, John Wiley & Sons, pp. 698-719
- Ho, C.H., Clark, B.R., Guerin, M.R., Barkenbus, B.D., Rao, T.K. & Epler, J.L. (1981) Analytical and biological analyses of test materials from the synthetic fuel technologies. IV. Studies of chemical structure-mutagenic activity relationships of aromatic nitrogen compounds relevant to synfuels. *Mutat. Res.*, 85, 335-345
- IARC (1987) *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, Suppl. 7, *Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs Volumes 1 to 42*, Lyon, pp. 261-263
- IARC (1988) *Information Bulletin on the Survey of Chemicals Being Tested for Carcinogenicity*, No. 13, Lyon, p. 20
- Johnson, D.E. & Cornish, H.H. (1978) Metabolic conversion of 1- and 2-nitronaphthalene to 1- and 2-naphthylamine in the rat. *Toxicol. appl. Pharmacol.*, 46, 549-553
- Johnson, D.E., Riley, M.G.I. & Cornish, H.H. (1984) Acute target organ toxicity of 1-nitronaphthalene in the rat. *J. appl. Toxicol.*, 4, 253-257
- Kadlubar, F.F., Unruh, L.E., Flammang, T.J., Sparks, D., Mitchum, R.K. & Mulder, G.J. (1981) Alteration of urinary levels of the carcinogen, *N*-hydroxy-2-naphthylamine, and its *N*-glucuronide in the rat by control of urinary pH, inhibition of metabolic sulfation, and changes in biliary excretion. *Chem.-biol. Interactions*, 33, 129-147
- Kitching, W., Bullpitt, M., Gartshore, D., Adcock, W., Khor, T.C., Doddrell, D. & Rae, I.D. (1977) Carbon-13 nuclear magnetic resonance examination of naphthalene derivatives. Assignments and analysis of substituent chemical shifts. *J. org. Chem.*, 42, 2411-2418

- Löfroth, G. (1981) Comparison of the mutagenic activity in carbon particulate matter and in diesel and gasoline engine exhaust. In: Waters, M.D., Sandhu, S., Lewtas Huising, J., Claxton, L. & Nesnow, S., eds, *Application of Short-term Bioassays in the Analysis of Complex Environmental Mixtures, II*, New York, Plenum, pp. 319–336
- Lucchini, V. & Wells, P.R. (1976) Proton magnetic resonance spectra of monosubstituted naphthalenes. *Org. magn. Resonance*, 8, 137–140
- McCann, J., Choi, E., Yamasaki, E. & Ames, B.N. (1975) Detection of carcinogens as mutagens in the *Salmonella* microsome test: assay of 300 chemicals. *Proc. natl Acad. Sci. USA*, 72, 5135–5139
- McCoy, E.C., Rosenkranz, E.J., Petrullo, L.A., Rosenkranz, H.S. & Mermelstein, R. (1981) Structural basis of the mutagenicity in bacteria of nitrated naphthalene and derivatives. *Environ. Mutagenesis*, 3, 499–511
- Mori, H., Sugie, S., Yoshimi, N., Kinouchi, T. & Ohnishi, Y. (1987) Genotoxicity of a variety of nitroarenes and other nitro compounds in DNA-repair tests with rat and mouse hepatocytes. *Mutat. Res.*, 190, 159–167
- Morotomi, M. & Watanabe, T. (1984) Metabolism of exogenous compounds by the intestinal microflora (Jpn.). *Tikoshikoroji Forumu [Toxicol. Forum]*, 7, 395–408
- Nakamura, S., Oda, Y., Shimada, T., Oki, I. & Sugimoto, K. (1987) SOS-inducing activity of chemical carcinogens and mutagens in *Salmonella typhimurium* TA1535/pSK1002: examination with 151 chemicals. *Mutat. Res.*, 192, 239–246
- Ohe, T. (1985) Studies on comparative decomposition rate by rat liver homogenate and on micronucleus test of nitrated polycyclic aromatic hydrocarbons. *Bull. environ. Contam. Toxicol.*, 34, 715–721
- Pienta, R.J. (1980) Transformation of Syrian hamster embryo cells by diverse chemicals and correlation with their reported carcinogenic and mutagenic activities. In: de Serres, F.J. & Hollaender, A., eds, *Chemical Mutagens, Principles and Methods for Their Detection*, Vol. 6, New York, Plenum, pp. 175–202
- Pitts, J.N., Jr (1987) Nitration of gaseous polycyclic aromatic hydrocarbons in simulated and ambient urban atmospheres: a source of mutagenic nitroarenes. *Atmos. Environ.*, 21, 2531–2547
- Pitts, J.N., Jr, Atkinson, R., Sweetman, J.A. & Zielinska, B. (1985) The gas-phase reactions of naphthalene with N_2O_5 to form nitronaphthalenes. *Atmos. Environ.*, 19, 701–705
- Poirier, L.A. & Weisburger, J.H. (1974) Enzymic reduction of carcinogenic aromatic nitro compounds by rat and mouse liver fractions. *Biochem. Pharmacol.*, 23, 661–669
- Ramdahl, T. & Urdal, K. (1982) Determination of nitrated polycyclic aromatic hydrocarbons by fused silica capillary gas chromatography/negative ion chemical ionization mass spectrometry. *Anal. Chem.*, 54, 2256–2260
- Ramdahl, T., Becher, G. & Bjørseth, A. (1982) Nitrated polycyclic aromatic hydrocarbons in urban air particles. *Environ. Sci. Technol.*, 16, 861–865
- Rosenkranz, H.S. & Mermelstein, R. (1983) Mutagenicity and genotoxicity of nitroarenes: all nitro-containing chemicals were not created equal. *Mutat. Res.*, 114, 217–267
- Rosenkranz, H.S. & Mermelstein, R. (1985) The genotoxicity, metabolism and carcinogenicity of nitrated polycyclic aromatic hydrocarbons. *J. environ. Sci. Health*, C3, 221–272
- Rosenkranz, H.S. & Poirier, L.A. (1979) Evaluation of the mutagenicity and DNA-modifying activity of carcinogens and non-carcinogens in microbial systems. *J. natl Cancer Inst.*, 62, 873–892

- Rosenkranz, H.S., McCoy, E.C., Sanders, D.R., Butler, M., Kiriazides, D.K. & Mermelstein, R. (1980) Nitropyrenes: isolation, identification and reduction of mutagenic impurities in carbon black and toners. *Science*, 209, 1039–1043
- Rosenkranz, H.S., McCoy, E.C., Frierson, M. & Klopman, G. (1985) The role of DNA sequence and structure of the electrophile on the mutagenicity of nitroarenes and arylamine derivatives. *Environ. Mutagenesis*, 7, 645–653
- Sanders, D.R. (1981) Nitropyrenes: the isolation of trace mutagenic impurities from the toluene extract of an aftertreated carbon black. In: Cooke, M. & Dennis, A.J., eds, *Polynuclear Aromatic Hydrocarbons, 5th International Symposium, Chemical Analysis and Biological Fate*, Columbus, OH, Battelle, pp. 145–158
- Schuetzle, D. & Jensen, T.E. (1985) Analysis of nitrated polycyclic aromatic hydrocarbons (nitro-PAH) by mass spectrometry. In: White, C.M., ed., *Nitrated Polycyclic Aromatic Hydrocarbons*, Heidelberg, A. Hüthig Verlag, pp. 121–167
- Scribner, J.D., Fisk, S.R. & Scribner, N.K. (1979) Mechanisms of action of carcinogenic aromatic amines: an investigation using mutagenesis in bacteria. *Chem.-biol. Interactions*, 26, 11–25
- Simmon, V.F. (1979a) In vitro mutagenicity assays of chemical carcinogens and related compounds with *Salmonella typhimurium*. *J. natl Cancer Inst.*, 62, 893–899
- Simmon, V.F. (1979b) In vitro assays for recombinogenic activity of chemical carcinogens and related compounds with *Saccharomyces cerevisiae* D3. *J. natl Cancer Inst.*, 62, 901–909
- Simmon, V.F., Rosenkranz, H.S., Zeiger, E. & Poirier, L.A. (1979) Mutagenic activity of chemical carcinogens and related compounds in the intraperitoneal host-mediated assay. *J. natl Cancer Inst.*, 62, 911–918
- Suzuki, J., Meguro, S. & Suzuki, S. (1987) Comparison of in vivo binding of aromatic nitro and amino compounds to rat hemoglobin (Abstract F 06-Y-13). *J. pharm. Sci.*, 76, S126
- Tokiwa, H. & Ohnishi, Y. (1986) Mutagenicity and carcinogenicity of nitroarenes and their sources in the environment. *CRC crit. Rev. Toxicol.*, 17, 23–69
- Tokiwa, H., Nakagawa, R., Horikawa, K. & Ohkubo, A. (1987) The nature of the mutagenicity and carcinogenicity of nitrated, aromatic compounds in the environment. *Environ. Health Perspect.*, 73, 191–199
- Verschueren, K. (1983) *Handbook of Environmental Data on Organic Chemicals*, 2nd ed., New York, Van Nostrand Reinhold, p. 917
- Wang, Y.Y., Rappaport, S.M., Sawyer, R.F., Talcott, R.E. & Wei, E.T. (1978) Direct-acting mutagens in automobile exhaust. *Cancer Lett.*, 5, 39–47
- Wang, C.Y., Lee, M.-S., King, C.M. & Warner, P.O. (1980) Evidence for nitroaromatics as direct-acting mutagens of airborne particulates. *Chemosphere*, 9, 83–87
- Weast, R.C. (1985) *CRC Handbook of Chemistry and Physics*, 66th ed., Boca Raton, FL, CRC Press, p. C-361