

2.18 *Cancer at other sites*

2.18.1 *Testis (Table 2.87)*

(a) *Parental exposure*

Among two cohort (Robinette *et al.*, 1979; Jensen, 1980) and three case-control studies (Schwartz *et al.*, 1962; Brown *et al.*, 1986; Weir *et al.*, 2000) conducted in the general population, only one case-control study suggested a possible association between testicular cancer in adults and maternal drinking during pregnancy (Brown *et al.*, 1986). The association was of borderline significance for the consumption of more than one drink per week relative to no drinking (odds ratio, 2.3; 95% CI, 1.0–5.2), but no association was observed for one drink (odds ratio, 1.1; 95% CI, 0.6–2.2), and no clear trend was apparent with the amount of alcohol consumed.

Table 2.87 Case-control studies of alcoholic beverage consumption and testicular cancer

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No. of cases	Relative risk (95% CI)	Adjustment factors	Comments
Parental exposure								
Brown <i>et al.</i> (1986), USA, 1979–81	225 mothers (pre- and perinatal cancer); response rate, 88%	213 mothers; response rate, 90%	Standardized telephone questionnaire	Never drinker		1.0	Tobacco smoking	
				1 drink/week		1.1 (0.6–2.2)		
				>1 drink/week		2.3 (1.0–5.2)		
						<i>p</i> -trend=0.14		
Weir <i>et al.</i> (2000), Ontario, Canada, 1987–89	346 case mothers/502 cases, aged 16–59 years; response rate, 80.8%	522 control mothers/ 975 controls; aged 16–59 years; response rate, 67.8%	Self-administered questionnaire	<i>Drinks/ week during pregnancy</i>			Age (5-year age group)	
				0	232	1.0		
				<2	83	1.2 (0.9–1.7)		
				≥2	24	0.8 (0.5–1.3)		
Chen <i>et al.</i> (2005b), USA, 1993–2001	278 incident childhood germ-cell; response rate, 80.8%	422; response rate, 66.6%; 1:2 match	Telephone interview; self-administered questionnaire	<i>Ever drank ≥6 months</i>			Gender of children, age, maternal education, race, family income	
				Never	182	1.0		
				Yes	92	0.9 (0.7–1.2)		
				<i>Ever drank during 1 month before pregnancy to nursing</i>				
				Never	126	1.0		
Yes	148	0.9 (0.7–1.2)						

Table 2.87 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No. of cases	Relative risk (95% CI)	Adjustment factors	Comments
Adult exposure								
Swerdlow <i>et al.</i> (1989), Oxford and West Midlands, United Kingdom 1977–81	259 cases of histologically confirmed testis cancer, aged ≥ 10 years	2 sets of controls: 238 radiotherapy controls treated in the same centres as cases; 251 non-radiotherapy controls who were general surgical, orthopaedic ENT and dental in-patients	Interview	Ever drank <i>Alcohol regularly?</i> Wine No Yes	NR	1.0 1.7 (1.21–2.43)	Social class	There was no dose–response relationship between risk for the tumour in relation to mean or to maximal wine consumption
UK Testicular Cancer Study Group (1994), United Kingdom, 1984–86	794, aged 15–49 years; response rate, 92%	609; 1:2 match (case/controls); response rate, 83.1%	Face-to-face interview	<i>Alcohol (g/week)</i> None <68.8 68.8–124.6 124.6–<211.2 211.2–<364.7 ≥ 364.7	92 150 147 130 135 140	1.0 1.26 (0.86–1.83) 1.23 (0.85–1.79) 0.87 (0.60–1.28) 1.06 (0.72–1.56) 1.13 (0.97–1.66) <i>p</i> -trend=0.41	Cryptorchidism, inguinal hernia at age <15 years	No evidence of an effect of testicular temperature on cancer risk

CI, confidence interval

One additional cohort study conducted among male and female cirrhotics in Denmark found a slightly increased risk for testicular cancer of all histological types (SIR, 2.3; 95% CI, 1.0–4.5) that varied little with type of cirrhosis and disappeared after 10 years of follow-up (Sørensen *et al.*, 1998).

One case–control study investigated the association of childhood germ-cell tumours (seminoma, embryonal carcinoma, yolk-sac tumour, choriocarcinoma, immature teratoma and mixed germ-cell tumours) and parental alcohol drinking (Chen *et al.*, 2005b). Results showed no association between germ-cell cancer overall and alcoholic beverage drinking by either parent before pregnancy, or during pregnancy or nursing; odds ratios were 0.9 (95% CI, 0.7–1.2) and 1.0 (95% CI, 0.8–1.3) for ever drinking, for mothers and fathers, respectively. Additional stratified analyses by sex, histological type and anatomical site did not show any association.

(b) *Adult exposure*

Two case–control studies in the United Kingdom investigated the association between alcoholic beverage drinking and testicular cancer. Swerdlow *et al.* (1989) found no association for regular alcoholic beverage drinking, duration of drinking or consumption of beer, cider or spirits; however, a significant association was found with regular consumption of wine, with an odds ratio of 1.71 (95% CI, 1.21–2.43), but no dose–response relation. The other case–control study found no association with alcohol intake at the time of diagnosis or at age 20 years (UK Testicular Cancer Study Group, 1994).

2.18.2 *Cancer of the brain*

(a) *Parental exposure and childhood brain cancer (Table 2.88)*

Only one cohort study found an association between alcoholic beverage consumption and brain cancer (Robinette *et al.*, 1979). Three additional studies with suboptimal methodology did not provide evidence of an association between increased alcoholic beverage consumption and brain cancer (IARC, 1988). However, a descriptive study based on cancer registries and national mortality data in France (Remontet *et al.*, 2003) showed a large increase in the incidence of and mortality from brain cancer between 1980 and 2000, during which time alcohol consumption decreased markedly.

Five case–control studies have assessed the association between alcoholic beverage consumption of parents and childhood brain cancer. Two of the studies were conducted in the USA and Canada (Bunin *et al.*, 1994; Yang *et al.*, 2000), one in China (Hu *et al.*, 2000), one in Germany (Schüz *et al.*, 2001) and one in the USA (Kramer *et al.*, 1987). Three of the studies examined the association between neuroblastoma and parental alcoholic beverage consumption (Kramer *et al.*, 1987; Yang *et al.*, 2000; Schüz *et al.*, 2001). Kramer *et al.* (1987) found a weak, non-significant association for any maternal alcoholic beverage drinking during pregnancy, with a suggestive increase

Table 2.88 Case-control studies of parental alcoholic beverage consumption and childhood brain tumours

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Kramer <i>et al.</i> (1987), Great Delaware Valley, USA, 1970–79	104 incident from the Great Delaware Valley Pediatric Tumor registry and the Cancer Research Center between 1970 and 1979; response rate, 74.8%	101; selection through RDD; response rate, 57.1%	Telephone interview	<i>Maternal drinking during pregnancy</i>			Not specified	90% CI reported; 1 drink=1 serving of beer, wine or liquor
				Any drinking	36	1.44 (0.94–2.21)		
				≥1 drink/day (frequent)	9	9.0 (2.16–37.56)		
				≥3 drinks/day (binge)	6	6.0 (1.26–28.54)		
Bunin <i>et al.</i> (1994), Canada, USA, 1986–89	322 diagnosed before 6 years of age in 1986–89; identified through the Children’s Cancer Group; response rate, 65%	321; selected through RDD; 1:1 match; response rate, 74%	Telephone interview of the mother or father	≥1 drink/day or ≥3 drinks occasionally	12	12.0 (3.14–45.82)	Income	*Crude odds ratio reported
				<i>Maternal exposure to beer during pregnancy</i>	10	1.4 (0.5–3.7)		
				Astrocytoma	12	4.0 (1.1–22.1)*		
				neurectoderma tumour				

Table 2.88 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Hu <i>et al.</i> (2000), Northeast, Heilongjiang Province, China, 1991–96	82 consecutive incident (43 boys, 39 girls) intracranial primary brain tumours, ≤18 years of age; 100%; residing in Heilongjiang Province at the time of diagnosis; 100% histologically confirmed; participation rate	3 individually matched per case; participation rate, 100%	Structured questionnaire (interview) administered to parents of all study subjects; history of parental liquor drinking obtained	<i>Lifetime paternal liquor consumption (L)</i>			Family income, mother's education, father's education	Similar associations for paternal age when started to drink liquor and numbers of years of drinking liquor; only one mother in the case group and two mothers in the control group reported drinking hard liquor.
				Never	41	1.00		
				≤200	20	3.21 (1.43–7.22)		
			≥201	21	4.43 (1.94–10.14)	<i>p</i> for trend=0.0001		

Table 2.88 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Yang <i>et al.</i> (2000), Canada, USA, 1992–94	538 children newly diagnosed with neuroblastoma in 1992–94, ≤19 years old; 100% histologically confirmed; response rate, 73%	504 mothers selected by RDD; 304 fathers directly interviewed; proxy interviews obtained for 142 (28%); 1:1 match; response rate, 72%	Structured telephone questionnaire to parents	<i>Maternal drinking</i>	253	0.9 (0.7–1.1)	Child's gender, mother's race and education, household income in the birth year	No association for paternal lifetime alcohol consumption, or before mother's pregnancy
				Lifetime	235	1.1 (0.8–1.4)		
				Around pregnancy ^a	205	1.1 (0.8–1.4)		
				1 month before conception	96	1.2 (0.9–1.7)		
				1st trimester	60	1.6 (1.0–2.4)		
				2nd trimester	58	1.4 (0.9–2.1)		
3rd trimester	54	1.0 (0.5–2.0)						
				Breastfeeding				

Table 2.88 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Schüz <i>et al.</i> , (2001), Germany, 1988–94	Pooled analysis of 2 case–control studies (1988–93; 1992–94); total of 192; children; response rate, 83.1%	2537; 2:1 match by gender and date of birth within 1 year; response rate, 71%	Questionnaire and telephone interview; same exposure assessment in both studies	Maternal alcohol consumption			Socioeconomic status, degree of urbanization	Odds ratio from a matched logistic regression on age, gender, birth year
				<i>Overall</i>				
				Never	140	1.0		
				1–7 glasses/week	38	0.84 (0.56–1.26)		
				>7 glasses/week	3	3.04 (0.75–12.2)		
				<i>Stage I/II</i>				
				Never	73	1.0		
				1–7 glasses/week	12	0.90 (0.45–1.80)		
>7 glasses/week	0	–						
<i>Stage III/VI</i>								
Never	39	1.0						
1–7 glasses/week	23	0.88 (0.53–1.45)						
>7 glasses/week	3	5.23 (1.33–20.6)						

CI, confidence interval; RDD, random-digit dialling

^a Exposure category includes drinking 1 month before pregnancy, during pregnancy and during breastfeeding

in risk with amount and frequency. However, these results were based on very small numbers of controls. A case–control study based on the Children’s Cancer Group and Paediatric Oncology Group institutions in the USA and Canada (Yang *et al.*, 2000) found no associations between the risk for neuroblastoma and either maternal or paternal alcoholic beverage consumption, while the combined analysis of two case–control studies used in the German study observed no overall association between maternal alcoholic beverage consumption during pregnancy and neuroblastoma or stage I/II neuroblastoma. However, an association was observed between advanced stage (III/IV) neuroblastoma and high alcoholic beverage consumption either during lifetime or around the time of pregnancy (Schüz *et al.*, 2001).

One study conducted in the USA and Canada found that maternal beer consumption during pregnancy was associated with primitive neuroectodermata tumours, but no association was found between alcoholic beverage consumption and astrocytoma (Bunin *et al.*, 1994), while the Chinese study reported that paternal hard liquor consumption before the pregnancy was associated with brain cancer (Hu *et al.*, 2000). [The Working Group considered that there was a possibility of recall bias in this study.]

(b) *Adult brain cancers (Table 2.89)*

One cohort study (Efird *et al.*, 2004) assessed associations between cigarette smoking and other lifestyle factors, including alcohol, and the occurrence of glioma in adults. There was no association with consumption of alcoholic beverages, beer or wine in the past year, although a slight non-significant association was observed for liquor consumption in the past year.

Nine case–control studies assessed the association between alcoholic beverage consumption and brain cancer in adults (Table 2.89). In studies conducted in Australia (Ryan *et al.*, 1992; Hurley *et al.*, 1996), Germany (Boeing *et al.*, 1993) and the USA (Preston-Martin *et al.*, 1989; Hochberg *et al.*, 1990; Lee *et al.*, 1997), no significant associations or trends were observed with the consumption of alcoholic beverages and the occurrence of glioma or meningioma. However, three studies, one conducted in Canada and two conducted in China, did find an association between the consumption of alcoholic beverages and brain cancer. The Canadian study found an elevated risk for ‘ever use’ of wine, but not of beer or spirits (Burch *et al.*, 1987) and one Chinese study (Hu *et al.*, 1998) found that consumption of liquor was associated with the occurrence of glioma in men with significant trends for the number of years of drinking, lifetime consumption and average consumption. However, no associations were seen for beer in adjusted analyses. In a separate report of the same study (Hu *et al.*, 1999), higher levels of consumption of beer, liquor and total alcohol were all associated with brain cancer, with respective adjusted odds ratios of 2.9 (95% CI, 1.1–7.6), 3.8 (95% CI, 1.6–9.2) and 3.2 (95% CI, 1.5–7.0) in the third tertile of consumption.

Table 2.89 Case-control studies of alcoholic beverage consumption and adult brain cancer

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments	
Choi <i>et al.</i> (1970), Minneapolis-St Paul Metropolitan area, USA, 1963-64	All (157) histologically proven primary tumours diagnosed in 4 hospitals between June and January 1963, and from June 1963 to June 1964; 126 histologically confirmed	157 patients admitted with conditions other than tumour of any site, neurological, psychiatric, ophthalmological or lymphatic disorders; matched on hospital of admission, sex, age, race, geographic area of residence, location of residence	Questionnaire interview	Central nervous system	<i>Verified tumours</i>			Age	Odds ratios and confidence intervals not presented; for subjects <20 years of age, his/her mother was approached for an interview; a proxy was interviewed when a subject could not provide proper responses.	
					Never	39	$p=0.008$			
					Ever	65				
					<i>Gliomas</i>					
					Never	20				
					Ever	35				
					<i>Astrocytoma</i>					
					Never	14				
					Ever	10				
					<i>Glioblastoma</i>					
Never	5									
Ever	23									
<i>Meningioma</i>										
Never	10	$p=0.007$								
Ever	14									

Table 2.89 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Musicco <i>et al.</i> (1982); Milan, Italy, 1979–80	51 patients hospitalized with gliomas, >20 years of age; mean age, 47 years; 15 astrocytomas, grades I and II; 10 oligodendrogliomas; and 26 astrocytomas, grades III and IV, and/or glioblastoma multiforme	201 admitted to the same hospital for meningioma, intervertebral disc prolapse or radiculitis, neuraxitis or multiple sclerosis, epilepsy, cerebrovascular disease, other neurological diseases; mean age 49 years; 2:1 matched for age, sex, place of residence	Interview	Central nervous system	Drinkers	24	1.0 <i>p</i> =1.000		Analyses based on 42 case–control pairs; patients who drank alcoholic beverages daily were considered drinkers; some diseases included in the control group may be linked to alcoholic beverage consumption; CI not reported.

Table 2.89 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Burch <i>et al.</i> (1987), southern Ontario, Canada, 1979–82	247 astrocytomas and glioblastomas (no meningiomas), aged 25–80 years; residents of metropolitan Toronto and southern Ontario; histologically confirmed through medical records; response rate, 75%	228 hospital-based, free of cancer; patients admitted to any hospital in the study area and who had a condition other than cancer at any site; response rate, 56%	Interviewer-administered questionnaire at home	Brain	<i>Beer</i>		1.0	Age, sex, proxy status, residence	Matched pair analysis
					Never		2.68 (1.18–6.07)		
					Low		0.49 (0.23–1.05)		
					Medium		1.47 (0.71–3.03)		
					High				
					<i>Spirits</i>		1.0		
					Never		1.29 (0.74–2.25)		
					Low		1.35 (0.50–3.65)		
					Medium		0.83 (0.41–1.71)		
					High				
					<i>Wine</i>		1.0		
					Never		1.06 (0.46–2.43)		
Low		2.07 (0.91–4.73)							
Medium		2.92 (1.20–7.07)							
High									
Preston-Martin <i>et al.</i> (1989), Los Angeles, USA, 1980–84	277 black and white men residing in Los Angeles County in 1980–1984, aged 25–49 years; first diagnosed with glioma or meningioma; response rate, 74%	272 neighbourhood; response rate, 98.2%	Face-to-face or telephone	Brain	<i>Glioma</i>			No adjustment specified	
					Beer at least once a month	32	0.7 (0.5–1.2)		
					Wine at least once a month	39	0.7 (0.5–1.1)		
					Liquor at least once a month	55	1.3 (0.8–1.9)		
					<i>Meningioma</i>				
					Beer at least once a month	7	0.4 (0.1–0.9)		
					Wine at least once a month	14	0.7 (0.3–1.4)		
Liquor at least once a month	15	0.7 (0.3–1.4)							

Table 2.89 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Hochberg <i>et al.</i> (1990), USA, 1977–81	160 newly diagnosed glioblastoma or astrocytoma identified in collaborating hospitals in Boston, Providence and Baltimore	128 friends of cases, excluding blood relatives; matched for sex, age (± 5 years), place of residence	Self-administered questionnaire, with telephone follow-up	Brain	Regular consumption of beer	67	0.7 (0.4–1.1)	Age, sex, socioeconomic status	Proxy interviews for 20% of cases and 2% of controls
Ryan <i>et al.</i> (1992), Adelaide, Australia, 1987–90	190 incident gliomas or meningiomas in 1987–90, aged 25–74 years; identified through the South Australian Central Cancer Registry; response rate, 90.5%	419 selected from the Australian electoral poll; 2:1 match; response rate, 63.3%	Face-to-face questionnaire at home or at work	Brain (191, 192)	<i>Glioma</i> Non-drinkers All sources 0–6.9 g/day 7–19.9 g/day ≥ 20 g/day <i>Meningioma</i> Non-drinkers All sources 0–6.9 g/day 7–19.9 g/day ≥ 20 g/day		1.0 0.94 (0.57–1.55) 0.86 (0.47–1.60) 0.74 (0.39–1.40) 1.00 (0.53–1.91) 1.0 0.59 (0.33–1.05) 0.63 (0.31–1.30) 0.49 (0.22–1.09) 0.58 (0.22–1.49)	Sex, age	Never drinkers were subjects who never drank at least once a month for a year; similar associations for beer, wine and spirit consumption.

Table 2.89 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Boeing <i>et al.</i> (1993), Southwest Germany, 1987–88	115 gliomas, 81 meningiomas and 30 acoustic neuromas, aged 25–75 years; 100% histopathologically confirmed; participation rate, 97.8%	418 randomly selected from the residential registries of the study area; participation rate, 72%	Standardized interview	Brain (191.0, 192.0, 192.1)	Consumption of alcoholic beverages assessed by lifelong history				No numerical data on alcohol presented; alcohol consumption was assessed by lifelong history; no significant association of risk for glioma or meningioma with lifelong consumption of a single alcoholic beverage or total alcohol.

Table 2.89 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Hurley <i>et al.</i> (1996), Australia (state of Victoria), 1987–91	416 incident (250 men, 166 women) primary gliomas, aged 20–70 years; identified through medical records from 14 Melbourne hospitals; 100% histologically confirmed; participation rate, 66% of eligible and 86% of the contacted cases	Selected from the electoral roll; 422 interviewed (252 men, 170 women); participation rate, 43.5% of those identified as eligible and 64.7% of the contacted controls	Structured questionnaire (interview); subjects sent a section of the questionnaire on details of some other variables	Brain (ICD-0 938–946)	Drank any alcoholic beverages	318	1.00 0.96 (0.67–1.37)	Age, gender, reference date	No increase in risk when average daily alcohol consumption considered
					<i>All</i>				
					Never Ever <i>Men</i> Never Ever				
					<i>Women</i> Never Ever				
Lee <i>et al.</i> (1997), California, USA 1991–1994	494 incident gliomas from 1991 to 1994, aged ≥20 years; identified through hospital records in the San Francisco Bay area; response rate, 82%	462 (random-digit dialling telephone number); frequency matched by age, gender, race/ethnicity; response rate, 63%	Structured questionnaire face-to-face	Brain (glioma) (ICD-0-2 9380–9481)	Mean consumption levels	No levels presented	Age, education, income	Only mean consumption levels of cases and controls presented; no significant differences noted	

Table 2.89 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Hu <i>et al.</i> (1998), China (Northeast, Heilongjiang Province), 1989–95	218 incident primary gliomas (139 astrocytomas and 79 other brain gliomas) identified from the Department of Neurosurgery of 6 major hospitals, aged 20–74 years; 100% histologically confirmed; participation rate, 100%	436 subjects with non-neoplastic, non-neurological diseases; 2:1 matched for sex, age, area of residence; participation rate, 100%.	Structured questionnaire (interview)	Brain	Liquor			Income, education, occupational exposure, consumption of vegetables and fruit; liquor also controlled for number of years of beer drinking, and beer controlled for number of years of liquor consumption	Only subjects directly interviewed included; associations for liquor similar for numbers of years drinking and lifetime liquor consumption; no associations noted for similar measures of beer consumption in the Hu <i>et al.</i> (1998) analysis, but were seen in an expanded analysis (Hu <i>et al.</i> , 1999, see text).
					<i>Age started to drink</i>				
					Never	55	1.00		
					≤20	54	1.98 (1.05–3.72)		
					≥21	31	1.40 (0.70–2.78)		
							<i>p</i> for trend=0.28		
<i>Average oz/day</i>									
Never	55	1.00							
≤2	38	1.54 (0.77–3.06)							
>2	47	1.87 (0.98–3.58)							

CI, confidence interval; ICD, International Classification of Diseases

2.18.3 *Cancer of the thyroid*

The association of alcoholic beverage consumption and thyroid cancer was examined in four cohort (Table 2.90) and six case–control (Tables 2.91) studies.

One cohort study among alcoholics in Sweden reported no significant excess risk for thyroid cancer compared with the general population (Adami *et al.*, 1992a). Two cohort studies conducted in the general population also reported no significant association of increasing alcohol consumption with risk for thyroid cancer (Iribarren *et al.*, 2001; Navarro Silvera *et al.*, 2005).

A pooled analysis of the case–control studies (Table 2.91), based on 1732 cases, found no association with increasing intake of beer and wine (relative risk, 0.9 (95% CI, 0.7–1.1) for more than 14 drinks per week) (Mack *et al.*, 2003). No difference was found for wine or beer separately or between men or women.

No data were available on the effect of duration of alcoholic beverage drinking or cessation of drinking on the risk for thyroid cancer.

2.18.4 *Melanoma*

(a) *Cohort studies (Table 2.92)*

Two cohort studies, one in a group of radiological technologists exposed to ionizing radiation in the USA (Freedman *et al.*, 2003) and one in alcoholic women in Sweden (Sigvardsson *et al.*, 1996), found no significant associations between the risk for melanoma and alcoholic beverage intake.

(b) *Case–control studies (Table 2.93)*

Six of nine case–control studies reported no significant association between alcoholic beverage intake and the risk for melanoma (Østerlind *et al.*, 1988; Bain *et al.*, 1993; Kirkpatrick *et al.*, 1994; Westerdahl *et al.*, 1996; Naldi *et al.*, 2004; Vinceti *et al.*, 2005). These studies were conducted in Australia, Italy, Denmark, Sweden and the USA.

Three case–control studies in the USA reported some increase in risk for melanoma associated with alcoholic beverage intake (Stryker *et al.*, 1990; Millen *et al.*, 2004; Le Marchand *et al.*, 2006). None of these were adjusted for exposure to ultraviolet light and thus the possibility of confounding can not be excluded.

2.18.5 *Other female cancers (vulva and vagina)*

(a) *Cohort studies (Table 2.94)*

Two cohort studies have examined the association between alcoholic beverage intake and risk for other female cancers. These studies were carried out in special populations, namely women being treated for alcohol abuse or alcoholism in Sweden (Sigvardsson *et al.*, 1996; Weiderpass *et al.*, 2001b). One study indicated an elevated

Table 2.90 Cohort studies of alcoholic beverage consumption and thyroid cancer

Reference, location, name of study	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)*	Adjustment factors	Comments
Special populations								
Hakulinen <i>et al.</i> (1974), Finland	Chronic alcoholic men (mean annual number in registry, 4370), aged >30 years, registered in 1967–70 when under custody of alcohol-misuse supervision, or when sent to a labour institute because of the vagrant law		Thyroid	Alcoholics	1 death observed/0.4 expected			No information regarding alcohol consumption, relative risk or CI was reported

Table 2.90 (continued)

Reference, location, name of study	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)*	Adjustment factors	Comments
Adami <i>et al.</i> (1992a), Uppsala, Sweden	9353 patients (8340 men; mean age at entry, 49.8 years; at diagnosis, 68.1 years; 1013 women; mean age at entry, 49.4 years; at diagnosis, 60.0 years) with a hospital discharge diagnosis of alcoholism in 1965–83	Follow-up through to 1984 (average follow-up, 7.7 years; maximum, 19 years)	Thyroid	No data on individual alcohol or tobacco use	Men: 3 Women: 0	SIR <i>Men</i> 1.7 (0.3–4.9) <i>Women</i> 0.0 (0.0–8.0)	Sex	

Table 2.90 (continued)

Reference, location, name of study	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)*	Adjustment factors	Comments
General population								
Iribarren <i>et al.</i> (2001), California, USA, Kaiser-Permanente Medical Care Program Cohort	94 549 men and women, aged 10–89 years, subscribers to the Kaiser Permanente Medical Care Program, northern California, who underwent regular health check-ups in 1964–73; follow-up based on the Cancer Incidence File (San Francisco Bay Area) through to 1997; median follow-up, 19.9 years	Self-administered questionnaire	Thyroid	<i>Alcohol consumption (drinks/day)</i> 0 1–2 3–5 ≥6		0.9 (0.6–1.3) 1.0 1.0 (0.5–1.8) 1.0 (0.3–3.0)	Age, sex, race, education, goitre, treatment to neck with X-rays, family history	Alcohol intake of 1–2 drinks/day = referent category; 73 cases of thyroid cancer in men and 123 cases in women; relative risk by gender not given

Table 2.90 (continued)

Reference, location, name of study	Cohort description	Exposure assessment	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)*	Adjustment factors	Comments
Navarro Silvera <i>et al.</i> (2005), Canada, Canadian National Breast Screening Study Cohort	49 613 women, aged 40–59 years, from the general Canadian population, recruited into the cohort between 1980 and 1985; average follow-up, 15.9 years	Self-administered questionnaire	Thyroid	<i>Alcohol intake (g/day)</i> None Any 1–3 3–10 ≥10	103 total	<i>Hazard ratio</i> 1.0 1.2 (0.7–1.8) 1.2 (0.7–2.0) 0.7 (0.4–1.2) 0.8 (0.5–1.4) <i>p</i> -trend=0.56	Age, education, pack–years of smoking, body mass index	No association for papillary or follicular subtype

CI, confidence interval; ICD, International Classification of Diseases

Table 2.91 Case-control studies of alcoholic beverage consumption and thyroid cancer

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Ron <i>et al.</i> (1987), Connecticut, USA, 1978–80	159 identified via Connecticut Tumor Registry; 100% histologically confirmed; response rate, 80%	285 population (random-digit dialling, Medicare records); 2:1 frequency-matched by sex, age; response rate, 65%	Interviewer-administered questionnaire	<i>Alcohol use</i> Non-user Any beer Any wine Any hard liquor	87 37 56 59	1.0 0.7 (0.4–1.3) 0.8 (0.5–1.3) 0.9 (0.6–1.5)	Age, sex, prior radiotherapy to the head and neck, thyroid nodules, goitre	Non-user: consumer of <1 drink per week
Kolonel <i>et al.</i> (1990), Hawaii, USA, 1980–97	191 (140 women, 51 men), identified through Hawaii Tumor registry, aged ≥ 18 years; 100% histologically confirmed; response rate, 79%	441 from Health Surveillance of the Department of Health; matched by age, sex; response rate, 74%	Self-administered questionnaire plus diet history	Regular alcohol use <i>Men</i> Never Ever <i>Women</i> Never Ever		1.0 0.6 (0.3–1.4) 1.0 1.0 (0.6–1.6)	Age, ethnicity	Number of cases not reported

Table 2.91 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Galanti <i>et al.</i> (1997), Norway/Sweden, 1993–94	Norway: 87 identified through Norwegian Cancer Register, born in Norway and living in the Tromsø Health Care Region, aged 18–75 years; response rate, 75% Sweden: 165 identified through registry, aged 18–75 years; response rate, 86%	Norway: 192 from population register; matched by age, sex; response rate, 56% Sweden: 248 from population register; matched by age, sex, county of residence; response rate, 69%.	Self-administered questionnaire	No. of drinks/month		Odds ratio (univariate analysis)		Not adjusted; results not changed after adjustment for smoking status, education
				<i>Wine (1.5 dL)</i>				
				<1	107	1.0		
				1–3	54	1.1 (0.7–1.7)		
				>3	52	0.7 (0.4–1.1)		
				<i>Light beer (2–5 dL)</i>				
				<1	113	1.0		
				1–4	61	1.0 (0.7–1.6)		
				>4	49	0.8 (0.5–1.2)		
				<i>Strong beer (2–5 dL)</i>				
				<1	181	1.0		
				>1	35	0.9 (0.5–1.6)		
				<i>Mild liquor (0.4 dL)</i>				
				<1	184	1.0		
>1	34	0.8 (0.5–1.2)						
<i>Hard liquor (0.4 dL)</i>								
<1	147	1.0						
>1	71	0.8 (0.5–1.1)						
<i>Ethanol (g/day)</i>								
<1	89	1.0						
1–3.95	80	0.8 (0.6–1.2)						
>3.95	67	0.7 (0.5–1.1)						

Table 2.91 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Chatenoud <i>et al.</i> (1999), Italy, 1983–93	428, aged <75 years; 100% histologically confirmed; refusal rate for interview, <3%	3526 hospital patients (non-malignant); excluded alcohol and tobacco- or dietary-related diseases	Interviewer-administered questionnaire	<i>Alcohol intake 2 years before</i> Lowest Highest		Odds ratio 1.0 1.7 (1.3–2.3)	Age, sex	The main focus of this study was on refined-cereal intake and risk for cancer; the quantity of alcohol consumed was not specified.

Table 2.91 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments		
Rossing <i>et al.</i> (2000), Washington State, USA, 1988–94	410 papillary tumours identified via the Washington State Cancer Surveillance System, aged 18–64 years; response rate, 84%	574 population (random-digit dialling); matched by age, county of residence; response rate, 74%	Interviewer-administered questionnaire	Alcohol intake			Odds ratio	Age	* Never drank ≥ 12 alcoholic drinks within 1 year; cases and controls were only women	
				Never*	126	1.0				
				>10 years ago	28	1.0 (0.5–1.7)				
				6–10 years ago	23	0.8 (0.5–1.5)				
				≤ 5 years ago	33	1.0 (0.6–1.8)				
				Amount (drinks/week)						
				Current drinkers						
				Never*	128	1.0				
				≤ 1	59	0.7 (0.4–1.0)				
				2–3	55	0.6 (0.4–0.9)				
				4–7	44	0.6 (0.4–0.9)				
				>7	42	0.9 (0.5–1.4)				
				Former drinkers						
				Never*	128	1.0				
≤ 1	42	1.2 (0.7–1.9)								
2–3	16	0.9 (0.5–1.9)								
4–7	6	0.3 (0.1–0.8)								
>7	18	1.2 (0.6–2.4)								
Pooled analyses										
Franceschi <i>et al.</i> (1991), 4 hospital-based case-control studies	385, aged <75 years; 100% histologically confirmed; response rate, ~97%	798 hospital patients (non-malignant)	Interviewer-administered questionnaire	Alcohol intake			Odds ratio	Age, sex, education, study centre	CI not reported	
				Low	103	1.0				
				Intermediate	122	1.1				
				High	160	1.3				
						χ^2 (trend), 2.72				

Table 2.91 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	No of cases	Relative risk (95% CI)	Adjustment factors	Comments
Mack <i>et al.</i> (2003), 10 case-control studies	370 men, 1296 women; six studies provided information on wine and beer combined	702 men, 2106 women	Pooled analysis	<i>Weekly drinks of wine and beer</i> None ≤2 >2–7 7–14 >14	787 263 321 146 149	<i>Men</i> 1.0 0.8 (0.6–1.0) 0.8 (0.7–1.0) 1.0 (0.8–1.3) 0.9 (0.7–1.1) <i>p</i> for trend 0.12	Stratification on study, age, sex, ethnicity, current smoking	No difference in cancer risk between men and women

CI, confidence interval

Table 2.92 Cohort studies of alcoholic beverage consumption and melanoma

Reference, location, name of study	Cohort description	Exposure assesment	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment factors	Comments
Sigvardsson <i>et al.</i> (1996), Sweden, Swedish Cancer Registry Study	15 508 alcoholic women individually matched for region and age with one non-alcoholic women; incidence data from the Swedish Cancer Registry	Alcoholic women from the records of the Temperance boards in Sweden	Reference Alcoholic women	28 14	1.0 0.5 (0.3–1.0)		[May be confounded by differences in smoking, dietary habits and/or other factors.]

Table 2.92 (continued)

Reference, location, name of study	Cohort description	Exposure assessment	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment factors	Comments
Freedman <i>et al.</i> (2003), USA, 1926–98 Radiologic Technologists Study	68 588 white cancer-free radiological technologists (54 045 women, 14 543 men); follow-up, 698 028 person-years; cases identified through SEER	Baseline questionnaire 1983–89 on height, weight, smoking, alcohol use, female hormonal factors, work history, other factors; participation rate, 86%; Second questionnaire 1994–98 updated information on risk factors, skin pigmentation, hair and eye colour, family medical history; participation rate, 83%	Alcohol (drinks/week)			Gender, years smoked, skin pigmentation, hair colour, personal history of non-melanoma skin cancer, decade of starting work as a technologist, education, proxy measures for residential childhood and adult exposure to sunlight	
			<i>Women</i>	159			
			Never	23	1.0		
			Ever	136	1.2 (0.8–1.9)		
			<1–6	114	1.2 (0.7–1.9)		
			7–14	19	1.7 (0.9–3.1)		
			>14	3	2.1 (0.6–7.0)		
					<i>p</i> for trend		0.05
			<i>Men</i>	48			
			Never	8	1.0		
			Ever	40	1.5 (0.7–3.3)		
			<1–6	32	1.5 (0.7–3.4)		
			7–14	4	0.9 (0.2–3.0)		
>14	4	2.4 (0.7–8.2)					
<i>All</i>	207						
Never	31	1.0					
Ever	176	1.3 (0.9–1.9)					
<1–6	146	1.2 (0.8–1.8)					
7–14	23	1.4 (0.8–2.5)					
>14	7	2.1 (0.9–4.8)					
		<i>p</i> for trend	0.08				

CI, confidence interval, SEER, Surveillance, Epidemiology and End Result

Table 2.93 Case-control studies of alcoholic beverage consumption and melanoma

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	Number of exposed cases	Odds ratio (95% CI)	Adjustment factors
Østerlind <i>et al.</i> (1988), East Denmark	474 incident, identified in the Danish Cancer Registry, aged 20–79 years; response rate, 92%	926 selected from National Population Register; response rate, 82%	Face-to-face structured questionnaire at home	<i>Alcoholic beverage</i>			Sunbathing, socioeconomic status
				Beer	0.7 (0.5–1.1)		
				Wine	0.7 (0.5–1.1)		
				Fortified wine	0.8 (0.5–1.2)		
				Distilled liquor	0.7 (0.5–1.1)		
				<i>Alcohol (kg/year)</i>			
0–1.1	1.0						
1.2–3.3	0.8 (0.6–1.1)						
3.4–8.4	0.8 (0.5–1.1)						
≥8.5	0.6 (0.4–0.9)						
Stryker <i>et al.</i> (1990), Massachusetts, USA, 1982–85	196 Caucasians; biopsy-confirmed cases older than 18 years; response rate, 92%	232 Caucasians; response rate, 92%	Face-to-face food-frequency questionnaire	Alcoholic bev.			Age, sex, hair colour, ability to tan
				<i>Beer</i>			
				None	1.0		
				<10 g/day	1.1		
				≥10 g/day	1.6		
					<i>p</i> trend=0.2		
				<i>Red wine</i>			
				None	1.0		
				<10 g/day	0.9		
				≥10 g/day	1.1		
	<i>p</i> trend=0.9						
<i>White wine</i>							
None	1.0						
<10 g/day	0.9						
≥10 g/day	0.8						
	<i>p</i> trend=0.9						

Table 2.93 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	Number of exposed cases	Odds ratio (95% CI)	Adjustment factors
Stryker <i>et al.</i> (1990) (contd)				<i>Liquor</i>			
				None		1.0	
				<10 g/day		1.3	
				≥10 g/day		1.2	
						<i>p</i> trend=0.7	
				<i>All types</i>			
None		1.0					
<10 g/day		1.2					
≥10 g/day		1.8 (1.0–3.3)					
		<i>p</i> trend=0.03					
Bain <i>et al.</i> (1993), Brisbane, Queensland, Australia, 1983–85	41 women, aged <80 years; histologically confirmed; [response rate, 63%]	297, aged <80 years; response rate not given	Mailed food-frequency questionnaire plus home interview	<i>Alcohol drinking (g/day)</i>			Age, hair colour, number of painful sunburns, total energy intake, number of years of schooling
			None		1.0		
			0.1–9.9		0.78 (0.32–1.94)		
			10.0–19.9		1.40 (0.46–4.30)		
			≥20.0		2.50 (0.87–7.40)		
Kirkpatrick <i>et al.</i> (1994), Washington State, USA, 1984–87	256 white, aged 25–65 years, identified from SEER cancer registry; response rate, 80%	234 identified by random-digit dialling to approximate age, sex, county of cases; response rate, 73%	Mailed food-frequency questionnaire plus telephone interview	<i>Drinks/month</i>			
				≤1	103	1.0	
				2–10	69	1.55	
				>10	62	1.18 (0.52–2.62)	
				≤1	103	1.0	
				2–10	69	1.31	
		>10	62	1.16 (0.53–2.59)			

Table 2.93 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	Number of exposed cases	Odds ratio (95% CI)	Adjustment factors
Westerdahl <i>et al.</i> (1996), southern Sweden, 1988–90	400 men and women, aged 15–75 years, from Regional Tumour Registry; histopathological diagnosis; response rate, 88.1%	640 population-based, selected by random sampling, matched 2:1 by sex, age, parish; response rate, 70.1%	Mailed comprehensive questionnaire	Any versus none		1.0 (0.7–1.4)	History of sunburn, hair colour, number of raised naevi
				Distilled alcohol >1/month		1.4 (1.0–1.9)	
				<i>Total alcohol intake (g/day)</i>			
				0	84	1.0	
				1–9	160	0.8 (0.6–1.1)	
10–19	37	0.9 (0.5–1.5)					
≥20	25	0.9 (0.5–1.8)	<i>p</i> trend>0.05				
Millen <i>et al.</i> (2004), Philadelphia, California, USA, 1991–92	497 newly diagnosed invasive cutaneous melanoma in two clinics, aged 20–79 years; 100% histologically confirmed; response rate, 84%	561 hospital-based; dermatological or psychiatric problems for clinic visit excluded; response rate, 66%	Food-frequency questionnaire	<i>Alcohol (times/week)</i>			Education, skin response after repeated sun exposure, age, sex, study site, presence of dysplastic nevi
				0	154	1.0	
				0.7	77	1.04 (0.69–1.57)	
				1.4–7.0	160	1.55 (1.09–2.20)	
				7.7–59	106	1.53 (1.03–2.29)	
				<i>p</i> for trend		0.04	

Table 2.93 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	Number of exposed cases	Odds ratio (95% CI)	Adjustment factors
Naldi <i>et al.</i> (2004), 27 centres in Italy, 1992–94	542 (226 men, 316 women), aged 15–87 years; 100% histologically confirmed; participation rate 99%	538 hospital-based (230 men, 308 women), aged 15–92 years; participation rate, 99%	Structured questionnaire, standardized examination	<i>Alcohol (drinks/week)</i> Never <1 1–13 14–27 ≥28	131 89 132 132 58	1.0 0.81 (0.53–1.22) 0.91 (0.62–1.33) 1.26 (0.83–1.91) 0.83 (0.49–1.40)	Age, sex, education, body mass index, history of sunburns, propensity to sunburn, number of naevi, number of freckles, skin, hair and eye colour, tobacco smoking

Table 2.93 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	Number of exposed cases	Odds ratio (95% CI)	Adjustment factors
Vinceti <i>et al.</i> (2005), Modena, Italy, 3 years	59 (28 men, 31 women newly diagnosed cutaneous melanomas attending the Dermatologic Clinic of Modena University Hospital (only centre for diagnosis, therapy and follow-up); 100% histologically confirmed; participation rate, 72%)	59 randomly selected residents of Modena; matched on sex, age	Self-administered questionnaire on diet and lifestyle habits	<i>Alcohol (g)</i> <1.6 ≥1.6–23.3 >23.3		1.0 1.86 (0.64–5.42) 0.97 (0.17–5.50)	Dietary factors, energy intake

Table 2.93 (continued)

Reference, location, period	Characteristics of cases	Characteristics of controls	Exposure assessment	Exposure categories	Number of exposed cases	Odds ratio (95% CI)	Adjustment factors
Le Marchand <i>et al.</i> (2006), Hawaii, USA, 1986–92	278 prevalent and incident (167 men, 111 women) invasive or in situ identified through Hawaii Tumor Registry with four grandparents of pure Caucasian origin; aged 18–79 years 100% histopathologically confirmed; participation rate, 67.5%	278 Caucasians randomly selected from local residential; registries matched to each case on sex, age; participation rate, 60.6%	Standardized interview by trained interviewers, including demographics, sun exposure, vacations, lifetime smoking, alcohol use, quantitative food-frequency questionnaire, skin colour, naevi, hair colour	Alcohol drinking status			Height, education, hair and eye colour, number of blistering sunburns at ages 10–17 years, ability to tan, family history
				<i>Men</i>			
				Never	22	1.0	
				Former	35	1.6 (0.8–3.4)	
				Current	110	1.9 (1.0–3.4)	
				<i>Women</i>			
				Never	35	1.0	
				Former	30	1.3 (0.6–2.6)	
				Current	46	1.5 (0.7–2.9)	
				Lifetime ethanol intake (kg)			
<i>Men</i>							
≤45	47	1.0					
>45–265	52	1.2 (0.6–2.2)					
>265	68	2.3 (1.2–4.4)					
<i>Women</i>							
≥0	35	1.0					
1–48.6	36	1.1 (0.5–2.4)					
>48.6	40	1.7 (0.7–3.8)					

CI, confidence interval; SEER, Surveillance, Epidemiology and End Result

Table 2.94 Cohort studies of alcoholic beverage consumption and other female cancers

Reference, location, name of study	Cohort description	Organ site (ICD code)	Exposure categories	No. of cases/deaths	Relative risk (95% CI)	Adjustment factors	Comments
Sigvardsson <i>et al.</i> (1996), Sweden, Temperance Boards Study	Nested case–control study; 15 508 alcoholic women born in 1870–1961 obtained from Temperance Boards; controls matched for region and day of birth; case ascertainment, Swedish Cancer Registry	Vulva, vagina and other female genital (ICD-7 176)	Alcohol abusers	16	4.0 (1.3–12)	Age, region	Estimate not adjusted for smoking
Weiderpass <i>et al.</i> (2001b), Sweden, National Board of Health and Welfare/Study of Alcoholic Women	36 856 women registered and hospitalized with alcoholism between 1965 and 1994; data from Inpatients Register; linkages to nationwide Registers of Causes of Death and Emigration and national Register of Cancer; mean age, 42.7 years; average follow-up time, 9.4 years	Vulva (ICD-7 176.0) Vagina (ICD-7 176.1)	Total <i>Age at cancer diagnosis</i> <50 years ≥50 years Total <i>Age at cancer diagnosis</i> <50 years ≥50 years	8 0 8 10 1 9	SIR 1.0 (0.4–2.0) – 1.2 (0.5–2.4) 4.6 (2.2–8.5) 2.5 (0.1–14.1) 5.1 (2.3–9.7)		Using expected rates specifically for squamous-cell carcinoma of the vulva, the overall SIR was 1.1 (0.5–2.2)

CI, confidence interval; ICD, International Classification of Diseases; SIR, standardized incidence ratio

risk for vaginal cancer but not for vulvar cancer (Weiderpass *et al.*, 2001b). The other study presented high relative risk estimates for both vulvar and vaginal cancers combined. The cohort studies could not adjust risk estimates for factors that may have confounded the association between alcoholic beverage and vulvar and vaginal cancers, such as HPV infections, number of sexual partners and tobacco smoking. It is possible that women who abuse alcohol have other behavioural patterns that may affect risks for vulvar and vaginal cancer.

(b) *Case-control studies (Table 2.95)*

Three case-control studies investigated the association between alcoholic beverage consumption and risk for vulvar cancer in Italy (Parazzini *et al.*, 1995b) and in the USA (Mabuchi *et al.*, 1985b; Sturgeon *et al.*, 1991). Two of these were hospital-based (Mabuchi *et al.*, 1985b; Parazzini *et al.*, 1995b) and one was population-based (Sturgeon *et al.*, 1991).

Confounding factors were considered in two studies (Sturgeon *et al.*, 1991; Parazzini *et al.*, 1995b), but only one provided risk estimates adjusted for smoking and sexual behaviour (Sturgeon *et al.*, 1991), which are potential confounders.

The three case-control studies reported no association between alcoholic beverage consumption and risk for vulva cancer.

(c) *Evidence of a dose-response*

One case-control study (Parazzini *et al.*, 1995b) and the cross-sectional study (Williams & Horm, 1977) presented information on dose-response for alcoholic beverage consumption and vulvar cancer. Neither study found evidence of a dose-response.

(d) *Types of alcoholic beverage*

Three studies (Williams & Horm, 1977; Mabuchi *et al.*, 1985b; Sturgeon *et al.*, 1991) investigated differences in risk according to the type of beverage and found no evidence of an effect.

Table 2.95 Case-control studies of alcoholic beverage consumption and other female cancers

Reference, study location and period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Williams & Horm (1977), The Third National Cancer Survey (cross-sectional study), USA, 1967-71	3856 cancer patients (all sites); age range not given; response rate, 57%	Randomly selected patients with cancers thought to be unrelated to tobacco and alcohol use	Personal interview	Vulva	Wine	0.63	Age, race, smoking	None of the values were significantly increased ($p > 0.05$) *less/more than one drink per week during a year
					$\leq 51^*$	-		
					>51			
					Beer			
					≤ 51	1.61		
					>51	0.84		
					Hard liquor			
≤ 51	1.67							
>51	0.43							
Total alcohol								
≤ 51	1.20							
>51	0.39							

Table 2.95 (continued)

Reference, study location and period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Mabuchi <i>et al.</i> (1985b), New York, Michigan, Florida, Minnesota, USA, 1972–75	149 patients with vulvar carcinoma from 155 hospitals; patient identification abstracted from hospital records; 100% histologically confirmed; participation rate, 79.7%	149 patients, admitted to the hospital for circulatory, digestive, nervous system, musculoskeletal, respiratory, genitourinary, endocrine, orthopaedic diseases, accidents and others; free of any cancer; matched to cases on hospital, sex, race, age (in 3-year range), marital status	Interview by blinded interviewers, mostly at hospital	Vulva	No association between alcohol consumption or specific alcoholic beverages and risk for vulvar cancer			

Table 2.95 (continued)

Reference, study location and period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Sturgeon <i>et al.</i> (1991), Chicago and Upstate New York, USA, 1985–87	201 incident cancer obtained from 34 hospitals in Chicago and Upstate New York, aged 53.9 years; 100% pathologically confirmed; participation rate, 61%	342 randomly selected using digit dialling techniques for controls <65 years and Health Care Financing Administration for women ≥65 years; mean age, 52.6 years; matched to cases by age in 5-year groups, race, residence; participation rate, 51%	Structured interview and food-frequency questionnaire at home	Vulva	No association between overall ethanol consumption and vulvar cancer; specific types of alcoholic beverage showed no appreciably increased risk with increasing intake.		Age, sexual behaviour, cigarette smoking	

Table 2.95 (continued)

Reference, study location and period	Characteristics of cases	Characteristics of controls	Exposure assessment	Organ site (ICD code)	Exposure categories	Relative risk (95% CI)	Adjustment for potential confounders	Comments
Parazzini <i>et al.</i> (1995b), Milan, Italy, 1987–93	125 admitted to general and teaching hospitals in the greater Milan area, aged 30–80 years; invasive vulvar cancer histologically confirmed	541 patients randomly selected, admitted to the same hospitals for acute conditions, not hormonal, gynaecological or neoplastic, aged 27–79 years; matched by age, interview year	Standard questionnaire; interview during hospital stay	Vulva	<i>Alcohol drinking</i> Never Occasional Regular	1.0 0.7 (0.4–1.2) 1.1 (0.7–1.7) χ^2 trend=0.17 $p=0.68$	Age, education, body mass index	Limited statistical power due to small study sample size; possible information bias

CI, confidence interval; ICD, International Classification of Diseases

References

- Adami HO, Hsing AW, McLaughlin JK *et al.* (1992b). Alcoholism and liver cirrhosis in the etiology of primary liver cancer. *Int J Cancer*, 51: 898–902. doi:10.1002/ijc.2910510611 PMID:1639537
- Adami HO, McLaughlin JK, Hsing AW *et al.* (1992a). Alcoholism and cancer risk: a population-based cohort study. *Cancer Causes Control*, 3: 419–425. doi:10.1007/BF00051354 PMID:1525322
- Adelstein A & White G (1976). Alcoholism and mortality. *Popul Trends*, 6: 7–13.
- Ahrens W, Jöckel KH, Patzak W, Elsner G (1991). Alcohol, smoking, and occupational factors in cancer of the larynx: a case-control study. *Am J Ind Med*, 20: 477–493. doi:10.1002/ajim.4700200404 PMID:1785612
- Akdaş A, Kirkali Z, Bilir N (1990). Epidemiological case-control study on the etiology of bladder cancer in Turkey. *Eur Urol*, 17: 23–26. PMID:2318234
- Akiba S (1994). Analysis of cancer risk related to longitudinal information on smoking habits. *Environ Health Perspect*, 102: Suppl 815–19. PMID:7851325
- Albertsen K & Grønbaek M (2002). Does amount or type of alcohol influence the risk of prostate cancer? *Prostate*, 52: 297–304. doi:10.1002/pros.10120 PMID:12210490
- Althuis MD, Brogan DD, Coates RJ *et al.* (2003). Breast cancers among very young premenopausal women (United States). *Cancer Causes Control*, 14: 151–160. doi:10.1023/A:1023006000760 PMID:12749720
- Altieri A, Bosetti C, Gallus S *et al.* (2004). Wine, beer and spirits and risk of oral and pharyngeal cancer: a case-control study from Italy and Switzerland. *Oral Oncol*, 40: 904–909. doi:10.1016/j.oraloncology.2004.04.005 PMID:15380168
- Altieri A, Bosetti C, Talamini R *et al.* (2002). Cessation of smoking and drinking and the risk of laryngeal cancer. *Br J Cancer*, 87: 1227–1229. doi:10.1038/sj.bjc.6600638 PMID:12439710
- Andersson S-O, Baron J, Bergström R *et al.* (1996). Lifestyle factors and prostate cancer risk: a case-control study in Sweden. *Cancer Epidemiol Biomarkers Prev*, 5: 509–513. PMID:8827354
- Asal NR, Risser DR, Kadamani S *et al.* (1988). Risk factors in renal cell carcinoma: I. Methodology, demographics, tobacco, beverage use, and obesity. *Cancer Detect Prev*, 11: 359–377. PMID:3390857
- Austin DF, Reynolds P (1996) Laryngeal cancer. In: Schottenfeld, D. & Fraumeni, J.F., eds, *Cancer Epidemiology and Prevention*, New York, Oxford University Press, pp. 619–636.
- Austin H, Drews C, Partridge EE (1993). A case-control study of endometrial cancer in relation to cigarette smoking, serum estrogen levels, and alcohol use. *Am J Obstet Gynecol*, 169: 1086–1091. PMID:8238164
- Baena AV, Allam MF, Del Castillo AS *et al.* (2006). Urinary bladder cancer risk factors in men: a Spanish case-control study. *Eur J Cancer Prev*, 15: 498–503. doi:10.1097/01.cej.0000215618.05757.04 PMID:17106329

- Baghurst PA, McMichael AJ, Slavotinek AH *et al.* (1991). A case-control study of diet and cancer of the pancreas. *Am J Epidemiol*, 134: 167–179. PMID:1862800
- Baglietto L, English DR, Gertig DM *et al.* (2005). Does dietary folate intake modify effect of alcohol consumption on breast cancer risk? Prospective cohort study. *BMJ*, 331: 807–900. doi:10.1136/bmj.38551.446470.06 PMID:16087654
- Baglietto L, Severi G, English DR *et al.* (2006). Alcohol consumption and prostate cancer risk: results from the Melbourne collaborative cohort study. *Int J Cancer*, 119: 1501–1504. doi:10.1002/ijc.21983 PMID:16615108
- Bagnardi V, Blangiardo M, La Vecchia C, Corrao G (2001). A meta-analysis of alcohol drinking and cancer risk. *Br J Cancer*, 85: 1700–1705. doi:10.1054/bjoc.2001.2140 PMID:11742491
- Bain C, Green A, Siskind V *et al.* (1993). Diet and melanoma. An exploratory case-control study. *Ann Epidemiol*, 3: 235–238. doi:10.1016/1047-2797(93)90024-X PMID:8275194
- Balaram P, Sridhar H, Rajkumar T *et al.* (2002). Oral cancer in southern India: the influence of smoking, drinking, paan-chewing and oral hygiene. *Int J Cancer*, 98: 440–445. doi:10.1002/ijc.10200 PMID:11920597
- Balder HF, Goldbohm RA, van den Brandt PA (2005). Dietary patterns associated with male lung cancer risk in the Netherlands Cohort Study. *Cancer Epidemiol Biomarkers Prev*, 14: 483–490. doi:10.1158/1055-9965.EPI-04-0353 PMID:15734976
- Band PR, Le ND, MacArthur AC *et al.* (2005). Identification of occupational cancer risks in British Columbia: a population-based case-control study of 1129 cases of bladder cancer. *J Occup Environ Med*, 47: 854–858. PMID:16093936
- Bandera EV, Freudenheim JL, Graham S *et al.* (1992). Alcohol consumption and lung cancer in white males. *Cancer Causes Control*, 3: 361–369. doi:10.1007/BF00146890 PMID:1617124
- Bandera EV, Freudenheim JL, Marshall JR *et al.* (1997). Diet and alcohol consumption and lung cancer risk in the New York State Cohort (United States) *Cancer Causes Control*, 8: 828–840. doi:10.1023/A:1018456127018 PMID:9427425
- Barra S, Barón AE, Franceschi S *et al.* (1991). Cancer and non-cancer controls in studies on the effect of tobacco and alcohol consumption. *Int J Epidemiol*, 20: 845–851. doi:10.1093/ije/20.4.845 PMID:1800421
- Barra S, Franceschi S, Negri E *et al.* (1990). Type of alcoholic beverage and cancer of the oral cavity, pharynx and oesophagus in an Italian area with high wine consumption. *Int J Cancer*, 46: 1017–1020. doi:10.1002/ijc.2910460612 PMID:2249890
- Baumgartner KB, Annegers JF, McPherson RS *et al.* (2002). Is alcohol intake associated with breast cancer in Hispanic women? The New Mexico Women's Health Study. *Ethn Dis*, 12: 460–469. PMID:12477131
- Begg CB, Walker AM, Wessen B, Zelen M (1983). Alcohol consumption and breast cancer. *Lancet*, 1: 293–294. doi:10.1016/S0140-6736(83)91703-8 PMID:6130308
- Benedetti A, Parent M-E, Siemiatycki J (2006). Consumption of alcoholic beverages and risk of lung cancer: results from two case-control studies in Montreal,

- Canada. *Cancer Causes Control*, 17: 469–480. doi:10.1007/s10552-005-0496-y PMID:16596299
- Benhamou S, Lenfant M-H, Ory-Paoletti C, Flamant R (1993). Risk factors for renal-cell carcinoma in a French case-control study. *Int J Cancer*, 55: 32–36. doi:10.1002/ijc.2910550107 PMID:8344750
- Beral V, Bull D, Reeves G Million Women Study Collaborators. (2005). Endometrial cancer and hormone-replacement therapy in the Million Women Study. *Lancet*, 365: 1543–1551. doi:10.1016/S0140-6736(05)66455-0 PMID:15866308
- Besson H, Brennan P, Becker N *et al.* (2006a). Tobacco smoking, alcohol drinking and non-Hodgkin's lymphoma: A European multicenter case-control study (EpiLymph). *Int J Cancer*, 119: 901–908. doi:10.1002/ijc.21913 PMID:16557575
- Besson H, Brennan P, Becker N *et al.* (2006b). Tobacco smoking, alcohol drinking and Hodgkin's lymphoma: a European multi-centre case-control study (EPILYMPH). *Br J Cancer*, 95: 378–384. doi:10.1038/sj.bjc.6603229 PMID:16819547
- Blot WJ, McLaughlin JK, Winn DM *et al.* (1988). Smoking and drinking in relation to oral and pharyngeal cancer. *Cancer Res*, 48: 3282–3287. PMID:3365707
- Boeing H (2002) Alcohol and cancer of the upper gastrointestinal tract: First analysis of the EPIC data. In: Riboli, E. & Lambert, R., eds, *Nutrition and lifestyles: Opportunities for Cancer Prevention* (IARC Scientific Publications No 156), Lyon, International Agency for Research on Cancer, pp. 151–154.
- Boeing H, Frentzel-Beyme R, Berger M *et al.* (1991). Case-control study on stomach cancer in Germany. *Int J Cancer*, 47: 858–864. doi:10.1002/ijc.2910470612 PMID:2010228
- Boeing H, Schlehofer B, Blettner M, Wahrendorf J (1993). Dietary carcinogens and the risk for glioma and meningioma in Germany. *Int J Cancer*, 53: 561–565. doi:10.1002/ijc.2910530406 PMID:8436429
- Boffetta P & Garfinkel L (1990). Alcohol drinking and mortality among men enrolled in an American Cancer Society prospective study. *Epidemiology*, 1: 342–348. doi:10.1097/00001648-199009000-00003 PMID:2078609
- Boffetta P, Stellman SD, Garfinkel L (1989). A case-control study of multiple myeloma nested in the American Cancer Society prospective study. *Int J Cancer*, 43: 554–559. doi:10.1002/ijc.2910430404 PMID:2703267
- Boffetta P, Ye W, Adami H-O *et al.* (2001). Risk of cancers of the lung, head and neck in patients hospitalized for alcoholism in Sweden. *Br J Cancer*, 85: 678–682. doi:10.1054/bjoc.2001.1986 PMID:11531251
- Boice JD Jr, Mandel JS, Doody MM (1995). Breast cancer among radiologic technologists. *JAMA*, 274: 394–401. doi:10.1001/jama.274.5.394 PMID:7616635
- Bosetti C, Franceschi S, Levi F *et al.* (2000). Smoking and drinking cessation and the risk of oesophageal cancer. *Br J Cancer*, 83: 689–691. doi:10.1054/bjoc.2000.1274 PMID:10944613

- Bosetti C, Gallus S, Franceschi S *et al.* (2002). Cancer of the larynx in non-smoking alcohol drinkers and in non-drinking tobacco smokers. *Br J Cancer*, 87: 516–518. doi:10.1038/sj.bjc.6600469 PMID:12189548
- Bosetti C, Garavello W, Gallus S, La Vecchia C (2006). Effects of smoking cessation on the risk of laryngeal cancer: an overview of published studies. *Oral Oncol*, 42: 866–872. doi:10.1016/j.oraloncology.2006.02.008 PMID:16931120
- Bouchardy C, Clavel F, La Vecchia C *et al.* (1990). Alcohol, beer and cancer of the pancreas. *Int J Cancer*, 45: 842–846. doi:10.1002/ijc.2910450509 PMID:2335387
- Boutron M-C, Faivre J, Dop MC *et al.* (1995). Tobacco, alcohol, and colorectal tumors: a multistep process. *Am J Epidemiol*, 141: 1038–1046. PMID:7771440
- Bravi F, Bosetti C, Scotti L *et al.* (2007). Food groups and renal cell carcinoma: a case-control study from Italy. *Int J Cancer*, 120: 681–685. doi:10.1002/ijc.22225 PMID:17058282
- Breslow RA, Graubard BI, Sinha R, Subar AF (2000). Diet and lung cancer mortality: a 1987 National Health Interview Survey cohort study. *Cancer Causes Control*, 11: 419–431. doi:10.1023/A:1008996208313 PMID:10877335
- Breslow RA, Wideroff L, Graubard BI *et al.* First National Health and Nutrition Examination Survey of the United States. (1999). Alcohol and prostate cancer in the NHANES I epidemiologic follow-up study. *Ann Epidemiol*, 9: 254–261. doi:10.1016/S1047-2797(98)00071-4 PMID:10332931
- Briggs NC, Levine RS, Bobo LD *et al.* (2002). Wine drinking and risk of non-Hodgkin's lymphoma among men in the United States: a population-based case-control study. *Am J Epidemiol*, 156: 454–462. doi:10.1093/aje/kwf058 PMID:12196315
- Britton JA, Gammon MD, Schoenberg JB *et al.* (2002). Risk of breast cancer classified by joint estrogen receptor and progesterone receptor status among women 20–44 years of age. *Am J Epidemiol*, 156: 507–516. doi:10.1093/aje/kwf065 PMID:12225998
- Brown LM, Gibson R, Burmeister LF *et al.* (1992). Alcohol consumption and risk of leukemia, non-Hodgkin's lymphoma, and multiple myeloma. *Leuk Res*, 16: 979–984. doi:10.1016/0145-2126(92)90077-K PMID:1405712
- Brown LM, Pottern LM, Hoover RN (1986). Prenatal and perinatal risk factors for testicular cancer. *Cancer Res*, 46: 4812–4816. PMID:3731127
- Brown LM, Pottern LM, Silverman DT *et al.* (1997). Multiple myeloma among Blacks and Whites in the United States: role of cigarettes and alcoholic beverages. *Cancer Causes Control*, 8: 610–614. doi:10.1023/A:1018498414298 PMID:9242477
- Brown LM, Silverman DT, Pottern LM *et al.* (1994). Adenocarcinoma of the esophagus and esophagogastric junction in white men in the United States: alcohol, tobacco, and socioeconomic factors. *Cancer Causes Control*, 5: 333–340. doi:10.1007/BF01804984 PMID:8080945
- Brownson RC (1988). A case-control study of renal cell carcinoma in relation to occupation, smoking, and alcohol consumption. *Arch Environ Health*, 43: 238–241. doi:10.1080/00039896.1988.9934940 PMID:3382249

- Bruemmer B, White E, Vaughan TL, Cheney CL (1997). Fluid intake and the incidence of bladder cancer among middle-aged men and women in a three-county area of western Washington. *Nutr Cancer*, 29: 163–168. doi:10.1080/01635589709514619 PMID:9427981
- Brugère J, Guenel P, Leclerc A, Rodriguez J (1986). Differential effects of tobacco and alcohol in cancer of the larynx, pharynx, and mouth. *Cancer*, 57: 391–395. doi:10.1002/1097-0142(19860115)57:2<391::AID-CNCR2820570235>3.0.CO;2-Q PMID:3942973
- Bueno de Mesquita HB, Maisonneuve P, Moerman CJ *et al.* (1992). Lifetime consumption of alcoholic beverages, tea and coffee and exocrine carcinoma of the pancreas: a population-based case-control study in The Netherlands. *Int J Cancer*, 50: 514–522. doi:10.1002/ijc.2910500403 PMID:1537615
- Bunin GR, Buckley JD, Boesel CP *et al.* (1994). Risk factors for astrocytic glioma and primitive neuroectodermal tumor of the brain in young children: a report from the Children's Cancer Group. *Cancer Epidemiol Biomarkers Prev*, 3: 197–204. PMID:8019366
- Burch JD, Craib KJP, Choi BCK *et al.* (1987). An exploratory case-control study of brain tumors in adults. *J Natl Cancer Inst*, 78: 601–609. PMID:3104645
- Burch JD, Howe GR, Miller AB, Semenciw R (1981). Tobacco, alcohol, asbestos, and nickel in the etiology of cancer of the larynx: a case-control study. *J Natl Cancer Inst*, 67: 1219–1224. PMID:6947107
- Byers T & Funch DP (1982). Alcohol and breast cancer. *Lancet*, 1: 799–800. doi:10.1016/S0140-6736(82)91841-4 PMID:6121249
- Byers T, Marshall J, Graham S *et al.* (1983). A case-control study of dietary and non-dietary factors in ovarian cancer. *J Natl Cancer Inst*, 71: 681–686. PMID:6578362
- Cantor KP, Lynch CF, Hildesheim ME *et al.* (1998). Drinking water source and chlorination byproducts. I. Risk of bladder cancer. *Epidemiology*, 9: 21–28. doi:10.1097/00001648-199801000-00007 PMID:9430264
- Carpenter CL, Morgenstern H, London SJ (1998). Alcoholic beverage consumption and lung cancer risk among residents of Los Angeles County. *J Nutr*, 128: 694–700. PMID:9521630
- Carstensen JM, Bygren LO, Hatschek T (1990). Cancer incidence among Swedish brewery workers. *Int J Cancer*, 45: 393–396. doi:10.1002/ijc.2910450302 PMID:2407667
- Cartwright RA, McKinney PA, O'Brien C *et al.* (1988). Non-Hodgkin's lymphoma: case control epidemiological study in Yorkshire. *Leuk Res*, 12: 81–88. doi:10.1016/S0145-2126(98)80012-X PMID:3357350
- Casagrande JT, Hanisch R, Pike MC *et al.* (1988). A case-control study of male breast cancer. *Cancer Res*, 48: 1326–1330. PMID:3342411
- Castelletto R, Muñoz N, Landoni N *et al.* (1992). Pre-cancerous lesions of the oesophagus in Argentina: prevalence and association with tobacco and alcohol. *Int J Cancer*, 51: 34–37. doi:10.1002/ijc.2910510107 PMID:1563842

- Castellsagué X, Muñoz N, De Stefani E *et al.* (2000). Smoking and drinking cessation and risk of esophageal cancer (Spain). *Cancer Causes Control*, 11: 813–818. doi:10.1023/A:1008984922453 PMID:11075870
- Castellsagué X, Quintana MJ, Martínez MC *et al.* (2004). The role of type of tobacco and type of alcoholic beverage in oral carcinogenesis. *Int J Cancer*, 108: 741–749. doi:10.1002/ijc.11627 PMID:14696101
- Cerhan JR, Torner JC, Lynch CF *et al.* (1997). Association of smoking, body mass, and physical activity with risk of prostate cancer in the Iowa 65+ Rural Health Study (United States). *Cancer Causes Control*, 8: 229–238. doi:10.1023/A:1018428531619 PMID:9134247
- Chang ET, Canchola AJ, Lee VS *et al.* (2007). Wine and other alcohol consumption and risk of ovarian cancer in the California Teachers Study cohort. *Cancer Causes Control*, 18: 91–103. doi:10.1007/s10552-006-0083-x PMID:17186425
- Chang ET, Hedelin M, Adami H-O *et al.* (2005). Alcohol drinking and risk of localized versus advanced and sporadic versus familial prostate cancer in Sweden. *Cancer Causes Control*, 16: 275–284. doi:10.1007/s10552-004-3364-2 PMID:15947879
- Chang ET, Smedby KE, Zhang SM *et al.* (2004). Alcohol intake and risk of non-Hodgkin lymphoma in men and women. *Cancer Causes Control*, 15: 1067–1076. doi:10.1007/s10552-004-2234-2 PMID:15801490
- Chatenoud L, La Vecchia C, Franceschi S *et al.* (1999). Refined-cereal intake and risk of selected cancers in Italy. *Am J Clin Nutr*, 70: 1107–1110. PMID:10584057
- Chen JS, Chen ZC, Chen XC *et al.* (2002b). Dietary and other living habits and the risk of gastric cancer in Changle, a high-risk area in China. *Chin J Nat Med*, 4: 131–134.
- Chen K, Jiang Q, Ma X *et al.* (2005a). Alcohol drinking and colorectal cancer: a population-based prospective cohort study in China. *Eur J Epidemiol*, 20: 149–154. doi:10.1007/s10654-004-2953-4 PMID:15792281
- Chen K-X, Xu W-L, Jia Z-L *et al.* (2003b). Risk factors of lung cancer in Tianjin. *Zhonghua Zhong Liu Za Zhi*, 25: 575–580. PMID:14690566
- Chen M-X, Chen S-D, Wang B-G *et al.* (2003c). Study on the relationship of genetic polymorphisms of human cytochrome P450 2E1 gene and susceptibility to lung cancer. *Sichuan Journal of Cancer Control*, 16: 129–131.
- Chen W, Sun XD, Fan JH *et al.* (2003a). The study for risk factors and changing rule of esophageal cancer in Linxian. *Cancer Res Clin*, 15: 5–7.
- Chen WY, Colditz GA, Rosner B *et al.* (2002a). Use of postmenopausal hormones, alcohol, and risk for invasive breast cancer. *Ann Intern Med*, 137: 798–804. PMID:12435216
- Chen Z, Robison L, Giller R *et al.* (2005b). Risk of childhood germ cell tumors in association with parental smoking and drinking. *Cancer*, 103: 1064–1071. doi:10.1002/cncr.20894 PMID:15685619
- Chen ZY, Zhao KG, Zhou GX *et al.* (2000). The main risk factors of esophageal cancer in Rugao city, Jiangsu: A case-control study. *Cancer Res Prev Treat*, 27: 240–243.

- Chen S, Zhen M, Wang B *et al.* (2004). A case-control study on the impact of CYP2E1 and GST-M1 polymorphisms on the risk of lung cancer. *Tumor Mar*, 24: 99–103.(in Chinese with English abstract).
- Cheng KK, Day NE, Duffy SW *et al.* (1992). Pickled vegetables in the aetiology of oesophageal cancer in Hong Kong Chinese. *Lancet*, 339: 1314–1318. doi:10.1016/0140-6736(92)91960-G PMID:1349991
- Cheng KK, Duffy SW, Day NE *et al.* (1995). Stopping drinking and risk of oesophageal cancer. *BMJ*, 310: 1094–1097. PMID:7742674
- Cheng YJ, Hildesheim A, Hsu MM *et al.* (1999). Cigarette smoking, alcohol consumption and risk of nasopharyngeal carcinoma in Taiwan. *Cancer Causes Control*, 10: 201–207. doi:10.1023/A:1008893109257 PMID:10454065
- Chiaffarino F, Gallus S, Negri E *et al.* (2002). Correspondence re: Weiderpass *et al.*, Alcoholism and risk of cancer of cervix uteri, vagina, and vulva. *Cancer Epidemiol Biomark. Prev.*, 10: 899–901, 2001. *Cancer Epidemiol Biomarkers Prev*, 11: 325–326. PMID:11895888
- Chiu BC-H, Cerhan JR, Gapstur SM *et al.* (1999). Alcohol consumption and non-Hodgkin lymphoma in a cohort of older women. *Br J Cancer*, 80: 1476–1482. doi:10.1038/sj.bjc.6690547 PMID:10424754
- Chiu BC-H, Weisenburger DD, Cantor KP *et al.* (2002). Alcohol consumption, family history of hematolymphoproliferative cancer, and the risk of non-Hodgkin's lymphoma in men. *Ann Epidemiol*, 12: 309–315. doi:10.1016/S1047-2797(01)00259-9 PMID:12062917
- Cho E, Smith-Warner SA, Ritz J *et al.* (2004). Alcohol intake and colorectal cancer: a pooled analysis of 8 cohort studies. *Ann Intern Med*, 140: 603–613. PMID:15096331
- Choi JY, Abel J, Neuhaus T *et al.* (2003). Role of alcohol and genetic polymorphisms of CYP2E1 and ALDH2 in breast cancer development. *Pharmacogenetics*, 13: 67–72. doi:10.1097/00008571-200302000-00002 PMID:12563175
- Choi NW, Schuman LM, Gullen WH (1970). Epidemiology of primary central nervous system neoplasms. II. Case-control study. *Am J Epidemiol*, 91: 467–485. PMID:4314683
- Choi SY & Kahyo H (1991a). Effect of cigarette smoking and alcohol consumption in the aetiology of cancer of the oral cavity, pharynx and larynx. *Int J Epidemiol*, 20: 878–885. doi:10.1093/ije/20.4.878 PMID:1800426
- Choi SY & Kahyo H (1991b). Effect of cigarette smoking and alcohol consumption in the etiology of cancers of the digestive tract. *Int J Cancer*, 49: 381–386. doi:10.1002/ijc.2910490312 PMID:1917136
- Chow WH, Schuman LM, McLaughlin JK *et al.* (1992). A cohort study of tobacco use, diet, occupation, and lung cancer mortality. *Cancer Causes Control*, 3: 247–254. doi:10.1007/BF00124258 PMID:1610971
- Chow W-H, Swanson CA, Lissowska J *et al.* (1999). Risk of stomach cancer in relation to consumption of cigarettes, alcohol, tea and coffee in Warsaw, Poland. *Int J Cancer*, 81:

- 871–876. doi:10.1002/(SICI)1097-0215(19990611)81:6<871::AID-IJC6>3.0.CO;2-# PMID:10362132
- Chyou P-H, Nomura AMY, Stemmermann GN (1993). A prospective study of diet, smoking, and lower urinary tract cancer. *Ann Epidemiol*, 3: 211–216. doi:10.1016/1047-2797(93)90021-U PMID:8275191
- Chyou P-H, Nomura AMY, Stemmermann GN (1995). Diet, alcohol, smoking and cancer of the upper aerodigestive tract: a prospective study among Hawaii Japanese men. *Int J Cancer*, 60: 616–621. doi:10.1002/ijc.2910600508 PMID:7860134
- Chyou P-H, Nomura AMY, Stemmermann GN (1996). A prospective study of colon and rectal cancer among Hawaii Japanese men. *Ann Epidemiol*, 6: 276–282. doi:10.1016/S1047-2797(96)00047-6 PMID:8876837
- Claude J, Kunze E, Frentzel-Beyme R *et al.* (1986). Life-style and occupational risk factors in cancer of the lower urinary tract. *Am J Epidemiol*, 124: 578–589. PMID:3752052
- Colditz GA & Rosner B (2000). Cumulative risk of breast cancer to age 70 years according to risk factor status: data from the Nurses' Health Study. *Am J Epidemiol*, 152: 950–964. doi:10.1093/aje/152.10.950 PMID:11092437
- Colditz GA, Rosner BA, Chen WY *et al.* (2004). Risk factors for breast cancer according to estrogen and progesterone receptor status. *J Natl Cancer Inst*, 96: 218–228. doi:10.1093/jnci/djh025 PMID:14759989
- Corrao G, Bagnardi V, Zambon A, La Vecchia C (2004). A meta-analysis of alcohol consumption and the risk of 15 diseases. *Prev Med*, 38: 613–619. doi:10.1016/j.ypmed.2003.11.027 PMID:15066364
- Cotterchio M, Kreiger N, Theis B *et al.* (2003). Hormonal factors and the risk of breast cancer according to estrogen- and progesterone-receptor subgroup. *Cancer Epidemiol Biomarkers Prev*, 12: 1053–1060. PMID:14578142
- Coughlin SS, Calle EE, Patel AV, Thun MJ (2000). Predictors of pancreatic cancer mortality among a large cohort of United States adults. *Cancer Causes Control*, 11: 915–923. doi:10.1023/A:1026580131793 PMID:11142526
- Crispo A, Talamini R, Gallus S *et al.* (2004). Alcohol and the risk of prostate cancer and benign prostatic hyperplasia. *Urology*, 64: 717–722. doi:10.1016/j.urology.2004.05.002 PMID:15491708
- Cui L, Yuan JC, Yang Y *et al.* (2001a). A contrast study of the risk factors of esophageal cancer in Jiangyan City. *Henan J Oncol*, 14: 406–407.
- Cui L, Yuan J-C, Yang Y *et al.* (2001b). A contrast study of the risky factors of male lung cancer in Jiangyan. *Henan J Oncol*, 8: 251–253.
- Cusimano R, Dardanoni G, Dardanoni L *et al.* (1989a). Risk factors of female cancers in Ragusa population (Sicily). 2. Breast cancer. *Eur J Epidemiol*, 5: 497–506. doi:10.1007/BF00140147 PMID:2606179
- Cusimano R, Dardanoni G, Dardanoni L *et al.* (1989b). Risk factors of female cancers in Ragusa population (Sicily)–1. Endometrium and cervix uteri cancers. *Eur J Epidemiol*, 5: 363–371. doi:10.1007/BF00144839 PMID:2792311

- Cuzick J & Babiker AG (1989). Pancreatic cancer, alcohol, diabetes mellitus and gall-bladder disease. *Int J Cancer*, 43: 415–421. doi:10.1002/ijc.2910430312 PMID:2925272
- D'Avanzo B, La Vecchia C, Franceschi S (1994). Alcohol consumption and the risk of gastric cancer. *Nutr Cancer*, 22: 57–64. doi:10.1080/01635589409514331 PMID:11304910
- Day GL, Blot WJ, McLaughlin JK, Fraumeni JF Jr (1994b). Carcinogenic risk of dark vs. light liquor. *Int J Cancer*, 59: 319–321. doi:10.1002/ijc.2910590305 PMID:7927935
- Day GL, Blot WJ, Shore RE *et al.* (1994a). Second cancers following oral and pharyngeal cancers: role of tobacco and alcohol. *J Natl Cancer Inst*, 86: 131–137. doi:10.1093/jnci/86.2.131 PMID:8271296
- De Stefani E, Boffetta P, Carzoglio J *et al.* (1998a). Tobacco smoking and alcohol drinking as risk factors for stomach cancer: a case-control study in Uruguay. *Cancer Causes Control*, 9: 321–329. doi:10.1023/A:1008829321668 PMID:9684712
- De Stefani E, Boffetta P, Deneo-Pellegrini H *et al.* (2007). The effect of smoking and drinking in oral and pharyngeal cancers: a case-control study in Uruguay. *Cancer Lett*, 246: 282–289. doi:10.1016/j.canlet.2006.03.008 PMID:16624486
- De Stefani E, Brennan P, Boffetta P *et al.* (2004). Comparison between hyperpharyngeal and laryngeal cancers: I-role of tobacco smoking and alcohol drinking. *Cancer Ther*, 2: 99–106.
- De Stefani E, Correa P, Deneo-Pellegrini H *et al.* (2002). Alcohol intake and risk of adenocarcinoma of the lung. A case-control study in Uruguay. *Lung Cancer*, 38: 9–14. doi:10.1016/S0169-5002(02)00153-8 PMID:12367787
- De Stefani E, Correa P, Fierro L *et al.* (1993). The effect of alcohol on the risk of lung cancer in Uruguay. *Cancer Epidemiol Biomarkers Prev*, 2: 21–26. PMID:8380549
- De Stefani E, Correa P, Oreggia F *et al.* (1987). Risk factors for laryngeal cancer. *Cancer*, 60: 3087–3091. doi:10.1002/1097-0142(19871215)60:12<3087::AID-CNCR2820601238>3.0.CO;2-6 PMID:3677031
- De Stefani E, Fierro L, Barrios E, Ronco A (1995). Tobacco, alcohol, diet and risk of prostate cancer. *Tumori*, 81: 315–320. PMID:8804446
- De Stefani E, Fierro L, Barrios E, Ronco A (1998b). Tobacco, alcohol, diet and risk of non-Hodgkin's lymphoma: a case-control study in Uruguay. *Leuk Res*, 22: 445–452. doi:10.1016/S0145-2126(97)00194-X PMID:9652731
- De Stefani E, Muñoz N, Estève J *et al.* (1990). Mate drinking, alcohol, tobacco, diet, and esophageal cancer in Uruguay. *Cancer Res*, 50: 426–431. PMID:2295081
- Dean G, MacLennan R, McLoughlin H, Shelley E (1979). Causes of death of blue-collar workers at a Dublin brewery, 1954--73. *Br J Cancer*, 40: 581–589. PMID:497108
- Dennis LK (2000). Meta-analysis for combining relative risks of alcohol consumption and prostate cancer. *Prostate*, 42: 56–66. doi:10.1002/(SICI)1097-0045(20000101)42:1<56::AID-PROS7>3.0.CO;2-P PMID:10579799

- Dikshit RP, Boffetta P, Bouchardy C *et al.* (2005). Risk factors for the development of second primary tumors among men after laryngeal and hypopharyngeal carcinoma. *Cancer*, 103: 2326–2333. doi:10.1002/cncr.21051 PMID:15852357
- Ding BG, Fan DM, Liu HJ *et al.* (2003). A case–control study on the risk factors of esophageal cancer in countryside. *China Tumor*, 12: 76–78.
- Ding JH, Li SP, Gao CM *et al.* (2001a). A case–control study of social factors and upper digestive tract cancer. *Jiangsu Med J*, 27: 9–11.
- Ding JH, Li SP, Gao CM *et al.* (2001b). On population based case–control study of upper digestive tract cancer. *China Pub Health*, 17: 319–320.
- Djoussé L, Dorgan JF, Zhang Y *et al.* (2002). Alcohol consumption and risk of lung cancer: the Framingham Study. *J Natl Cancer Inst*, 94: 1877–1882. PMID:12488481
- Djoussé L, Schatzkin A, Chibnik LB *et al.* (2004). Alcohol consumption and the risk of bladder cancer in the Framingham Heart Study. *J Natl Cancer Inst*, 96: 1397–1400. doi:10.1093/jnci/djh263 PMID:15367573
- Doll R, Forman D, La Vecchia C, Woutersen RA (1999) Alcoholic beverages and cancers of the digestive tract and larynx. In: Macdonald, I. ed., *Health Issues Related to Alcohol Consumption*, Oxford, ILSI Europe, Blackwell Science, pp. 351–393.
- Doll R, Peto R, Boreham J, Sutherland I (2005). Mortality from cancer in relation to smoking: 50 years observations on British doctors. *Br J Cancer*, 92: 426–429. PMID:15668706
- Doll R, Peto R, Hall E *et al.* (1994). Mortality in relation to consumption of alcohol: 13 years' observations on male British doctors. *BMJ*, 309: 911–918. PMID:7950661
- Donato F, Boffetta P, Fazioli R *et al.* (1997). Bladder cancer, tobacco smoking, coffee and alcohol drinking in Brescia, northern Italy. *Eur J Epidemiol*, 13: 795–800. doi:10.1023/A:1007453322899 PMID:9384269
- Donato F, Tagger A, Gelatti U *et al.* (2002). Alcohol and hepatocellular carcinoma: the effect of lifetime intake and hepatitis virus infections in men and women. *Am J Epidemiol*, 155: 323–331. doi:10.1093/aje/155.4.323 PMID:11836196
- Dosemeci M, Gokmen I, Unsal M *et al.* (1997). Tobacco, alcohol use, and risks of laryngeal and lung cancer by subsite and histologic type in Turkey. *Cancer Causes Control*, 8: 729–737. doi:10.1023/A:1018479304728 PMID:9328195
- Dumeaux V, Lund E, Hjartåker A (2004). Use of oral contraceptives, alcohol, and risk for invasive breast cancer. *Cancer Epidemiol Biomarkers Prev*, 13: 1302–1307. PMID:15298950
- Dupont WD & Page DL (1985). Risk factors for breast cancer in women with proliferative breast disease. *N Engl J Med*, 312: 146–151. doi:10.1056/NEJM198501173120303 PMID:3965932
- Durbec JP, Chevillotte G, Bidart JM *et al.* (1983). Diet, alcohol, tobacco and risk of cancer of the pancreas: a case-control study. *Br J Cancer*, 47: 463–470. PMID:6849792
- Efrid JT, Friedman GD, Sidney S *et al.* (2004). The risk for malignant primary adult-onset glioma in a large, multiethnic, managed-care cohort: cigarette smoking and other life-

- style behaviors. *J Neurooncol*, 68: 57–69. doi:10.1023/B:NEON.0000024746.87666.ed PMID:15174522
- Ellison LF (2000). Tea and other beverage consumption and prostate cancer risk: a Canadian retrospective cohort study. *Eur J Cancer Prev*, 9: 125–130. doi:10.1097/00008469-200004000-00009 PMID:10830580
- Elwood JM, Pearson JC, Skippen DH, Jackson SM (1984). Alcohol, smoking, social and occupational factors in the aetiology of cancer of the oral cavity, pharynx and larynx. *Int J Cancer*, 34: 603–612. doi:10.1002/ijc.2910340504 PMID:6500740
- Enger SM, Ross RK, Paganini-Hill A *et al.* (1999). Alcohol consumption and breast cancer oestrogen and progesterone receptor status. *Br J Cancer*, 79: 1308–1314. doi:10.1038/sj.bjc.6690210 PMID:10098777
- Ewings P & Bowie C (1996). A case-control study of cancer of the prostate in Somerset and east Devon. *Br J Cancer*, 74: 661–666. PMID:8761387
- Falcao JM, Dias JA, Miranda AC *et al.* (1994). Red wine consumption and gastric cancer in Portugal: a case-control study. *Eur J Cancer Prev*, 3: 269–276. doi:10.1097/00008469-199403030-00005 PMID:8061592
- Falk RT, Pickle LW, Brown LM *et al.* (1989). Effect of smoking and alcohol consumption on laryngeal cancer risk in coastal Texas. *Cancer Res*, 49: 4024–4029. PMID:2736543
- Falk RT, Pickle LW, Fontham ET *et al.* (1988). Life-style risk factors for pancreatic cancer in Louisiana: a case-control study. *Am J Epidemiol*, 128: 324–336. PMID:3394699
- Fan ZH, Chu TX, Fan ZL *et al.* (1996). A cohort study on the relationship between alcohol consumption and mortality of digestive tract cancer. *Mod Prev Med*, 23: 20–22.
- Farrow DC & Davis S (1990). Risk of pancreatic cancer in relation to medical history and the use of tobacco, alcohol and coffee. *Int J Cancer*, 45: 816–820. doi:10.1002/ijc.2910450504 PMID:2335385
- Fei SJ & Xiao SD (2004). Diet and gastric cancer: A case-control study in Shanghai urban districts. *Chin J Gastroenterol*, 8: 143–146.
- Feigelson HS, Jonas CR, Robertson AS *et al.* (2003). Alcohol, folate, methionine, and risk of incident breast cancer in the American Cancer Society Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomarkers Prev*, 12: 161–164. PMID:12582027
- Ferraroni M, Negri E, La Vecchia C *et al.* (1989). Socioeconomic indicators, tobacco and alcohol in the aetiology of digestive tract neoplasms. *Int J Epidemiol*, 18: 556–562. doi:10.1093/ije/18.3.556 PMID:2807657
- Fincham SM, Hill GB, Hanson J, Wijayasinghe C (1990). Epidemiology of prostatic cancer: a case-control study. *Prostate*, 17: 189–206. doi:10.1002/pros.2990170303 PMID:2235728
- Fioretti F, Bosetti C, Tavani A *et al.* (1999). Risk factors for oral and pharyngeal cancer in never smokers. *Oral Oncol*, 35: 375–378. doi:10.1016/S1368-8375(98)00125-0 PMID:10645401

- Flanders WD & Rothman KJ (1982). Interaction of alcohol and tobacco in laryngeal cancer. *Am J Epidemiol*, 115: 371–379. PMID:7064973
- Flood A, Caprario L, Chatterjee N *et al.* (2002). Folate, methionine, alcohol, and colorectal cancer in a prospective study of women in the United States. *Cancer Causes Control*, 13: 551–561. doi:10.1023/A:1016330609603 PMID:12195645
- Folsom AR, Demissie Z, Harnack L Iowa Women's Health Study. (2003). Glycemic index, glycemic load, and incidence of endometrial cancer: the Iowa women's health study. *Nutr Cancer*, 46: 119–124. doi:10.1207/S15327914NC4602_03 PMID:14690786
- Franceschi S, Levi F, Dal Maso L *et al.* (2000). Cessation of alcohol drinking and risk of cancer of the oral cavity and pharynx. *Int J Cancer*, 85: 787–790. doi:10.1002/(SICI)1097-0215(20000315)85:6<787::AID-IJC8>3.0.CO;2-6 PMID:10709096
- Franceschi S, Levi F, La Vecchia C *et al.* (1999). Comparison of the effect of smoking and alcohol drinking between oral and pharyngeal cancer. *Int J Cancer*, 83: 1–4. doi:10.1002/(SICI)1097-0215(19990924)83:1<1::AID-IJC1>3.0.CO;2-8 PMID:10449598
- Franceschi S, Levi F, Negri E *et al.* (1991). Diet and thyroid cancer: a pooled analysis of four European case-control studies. *Int J Cancer*, 48: 395–398. doi:10.1002/ijc.2910480315 PMID:2040535
- Franceschi S, Montella M, Polesel J *et al.* (2006). Hepatitis viruses, alcohol, and tobacco in the etiology of hepatocellular carcinoma in Italy. *Cancer Epidemiol Biomarkers Prev*, 15: 683–689. doi:10.1158/1055-9965.EPI-05-0702 PMID:16614109
- Franceschi S, Talamini R, Barra S *et al.* (1990). Smoking and drinking in relation to cancers of the oral cavity, pharynx, larynx, and esophagus in northern Italy. *Cancer Res*, 50: 6502–6507. PMID:2208109
- Freedman DM, Sigurdson A, Doody MM *et al.* (2003). Risk of melanoma in relation to smoking, alcohol intake, and other factors in a large occupational cohort. *Cancer Causes Control*, 14: 847–857. doi:10.1023/B:CACO.0000003839.56954.73 PMID:14682442
- Freudenheim JL, Ambrosone CB, Moysich KB *et al.* (1999). Alcohol dehydrogenase 3 genotype modification of the association of alcohol consumption with breast cancer risk. *Cancer Causes Control*, 10: 369–377. doi:10.1023/A:1008950717205 PMID:10530606
- Freudenheim JL, Graham S, Byers TE *et al.* (1992). Diet, smoking, and alcohol in cancer of the larynx: a case-control study. *Nutr Cancer*, 17: 33–45. doi:10.1080/01635589209514171 PMID:1574443
- Freudenheim JL, Graham S, Marshall JR *et al.* (1990). Lifetime alcohol intake and risk of rectal cancer in western New York. *Nutr Cancer*, 13: 101–109. doi:10.1080/01635589009514050 PMID:2300490
- Freudenheim JL, Graham S, Marshall JR *et al.* (1991). Folate intake and carcinogenesis of the colon and rectum. *Int J Epidemiol*, 20: 368–374. doi:10.1093/ije/20.2.368 PMID:1917236

- Freudenheim JL, Marshall JR, Graham S *et al.* (1995). Lifetime alcohol consumption and risk of breast cancer. *Nutr Cancer*, 23: 1–11. doi:10.1080/01635589509514356 PMID:7739910
- Freudenheim JL, Ram M, Nie J *et al.* (2003). Lung cancer in humans is not associated with lifetime total alcohol consumption or with genetic variation in alcohol dehydrogenase 3 (ADH3). *J Nutr*, 133: 3619–3624. PMID:14608084
- Freudenheim JL, Ritz J, Smith-Warner SA *et al.* (2005). Alcohol consumption and risk of lung cancer: a pooled analysis of cohort studies. *Am J Clin Nutr*, 82: 657–667. PMID:16155281
- Friedenreich CM, Howe GR, Miller AB, Jain MG (1993). A cohort study of alcohol consumption and risk of breast cancer. *Am J Epidemiol*, 137: 512–520. PMID:8465803
- Friedman GD & van den Eeden SK (1993). Risk factors for pancreatic cancer: an exploratory study. *Int J Epidemiol*, 22: 30–37. doi:10.1093/ije/22.1.30 PMID:8449644
- Fuchs CS, Stampfer MJ, Colditz GA *et al.* (1995). Alcohol consumption and mortality among women. *N Engl J Med*, 332: 1245–1250. doi:10.1056/NEJM199505113321901 PMID:7708067
- Gajalakshmi CK & Shanta V (1996). Lifestyle and risk of stomach cancer: a hospital-based case-control study. *Int J Epidemiol*, 25: 1146–1153. doi:10.1093/ije/25.6.1146 PMID:9027518
- Gajalakshmi V, Hung RJ, Mathew A *et al.* (2003). Tobacco smoking and chewing, alcohol drinking and lung cancer risk among men in southern India. *Int J Cancer*, 107: 441–447. doi:10.1002/ijc.11377 PMID:14506745
- Galanti MR, Hansson L, Bergström R *et al.* (1997). Diet and the risk of papillary and follicular thyroid carcinoma: a population-based case-control study in Sweden and Norway. *Cancer Causes Control*, 8: 205–214. doi:10.1023/A:1018424430711 PMID:9134245
- Gallagher RP, Spinelli JJ, Elwood JM, Skippen DH (1983). Allergies and agricultural exposure as risk factors for multiple myeloma. *Br J Cancer*, 48: 853–857. PMID:6652026
- Gallus S, Bosetti C, Franceschi S *et al.* (2001). Oesophageal cancer in women: tobacco, alcohol, nutritional and hormonal factors. *Br J Cancer*, 85: 341–345. doi:10.1054/bjoc.2001.1898 PMID:11487262
- Gammon MD, Hibshoosh H, Terry MB *et al.* (1999). Oral contraceptive use and other risk factors in relation to HER-2/neu overexpression in breast cancer among young women. *Cancer Epidemiol Biomarkers Prev*, 8: 413–419. PMID:10350436
- Gammon MD, Neugut AI, Santella RM *et al.* (2002). The Long Island Breast Cancer Study Project: description of a multi-institutional collaboration to identify environmental risk factors for breast cancer. *Breast Cancer Res Treat*, 74: 235–254. doi:10.1023/A:1016387020854 PMID:12206514
- Gammon MD, Schoenberg JB, Ahsan H *et al.* (1997). Tobacco, alcohol, and socio-economic status and adenocarcinomas of the esophagus and gastric cardia. *J Natl Cancer Inst*, 89: 1277–1284. doi:10.1093/jnci/89.17.1277 PMID:9293918

- Gao CM, Takezaki T, Sugimura H *et al.* (2001). The impact of CYP2E1 Rsa I GSTT1 and GSTM1 polymorphisms on the risk of esophageal cancer. *China Tumor*, 10: 346–349.
- Gao CM, Takezaki T, Wu JZ *et al.* (2002a). Effects of GSTT1 genotypes, lifestyle factors and their interactions on risk of esophageal and stomach cancers. *China Cancer Prev Treat*, 9: 113–117.
- Gao CM, Wu JZ, Ding JH *et al.* (2002b). Polymorphisms of methylenetetrahydrofolate reductase C677T and the risk of stomach cancer *Zhonghua Liu Xing Bing Xue Za Zhi*, 23: 289–292. PMID:12411076
- Gapstur SM, Potter JD, Drinkard C, Folsom AR (1995). Synergistic effect between alcohol and estrogen replacement therapy on risk of breast cancer differs by estrogen/progesterone receptor status in the Iowa Women's Health Study. *Cancer Epidemiol Biomarkers Prev*, 4: 313–318. PMID:7655324
- Gapstur SM, Potter JD, Sellers TA *et al.* (1993). Alcohol consumption and postmenopausal endometrial cancer: results from the Iowa Women's Health Study. *Cancer Causes Control*, 4: 323–329. doi:10.1007/BF00051334 PMID:8347781
- Gapstur SM, Potter JD, Sellers TA, Folsom AR (1992). Increased risk of breast cancer with alcohol consumption in postmenopausal women. *Am J Epidemiol*, 136: 1221–1231. PMID:1476144
- Garavello W, Bosetti C, Gallus S *et al.* (2006). Type of alcoholic beverage and the risk of laryngeal cancer. *Eur J Cancer Prev*, 15: 69–73. doi:10.1097/01.cej.0000186641.19872.04 PMID:16374233
- Garfinkel L, Boffetta P, Stellman SD (1988). Alcohol and breast cancer: a cohort study. *Prev Med*, 17: 686–693. doi:10.1016/0091-7435(88)90086-2 PMID:3244667
- Garland C, Shekelle RB, Barrett-Connor E *et al.* (1985). Dietary vitamin D and calcium and risk of colorectal cancer: a 19-year prospective study in men. *Lancet*, 1: 307–309. doi:10.1016/S0140-6736(85)91082-7 PMID:2857364
- Garland M, Hunter DJ, Colditz GA *et al.* (1999). Alcohol consumption in relation to breast cancer risk in a cohort of United States women 25–42 years of age. *Cancer Epidemiol Biomarkers Prev*, 8: 1017–1021. PMID:10566558
- Garrote LF, Herrero R, Reyes RM *et al.* (2001). Risk factors for cancer of the oral cavity and oro-pharynx in Cuba. *Br J Cancer*, 85: 46–54. doi:10.1054/bjoc.2000.1825 PMID:11437401
- Gelatti U, Covolo L, Talamini R *et al.* (2005). N-Acetyltransferase-2, glutathione S-transferase M1 and T1 genetic polymorphisms, cigarette smoking and hepatocellular carcinoma: a case-control study. *Int J Cancer*, 115: 301–306. doi:10.1002/ijc.20895 PMID:15688397
- Gerhardsson de Verdier M, Romelsjö A, Lundberg M (1993). Alcohol and cancer of the colon and rectum. *Eur J Cancer Prev*, 2: 401–408. doi:10.1097/00008469-199309000-00007 PMID:8401175
- Ghadirian P, Simard A, Baillargeon J (1991). Tobacco, alcohol, and coffee and cancer of the pancreas. A population-based, case-control study in Quebec, Canada.

- Cancer*, 67: 2664–2670. doi:10.1002/1097-0142(19910515)67:10<2664::AID-CNCR2820671043>3.0.CO;2-K PMID:2015568
- Giovannucci E, Rimm EB, Ascherio A *et al.* (1995). Alcohol, low-methionine–low-folate diets, and risk of colon cancer in men. *J Natl Cancer Inst*, 87: 265–273. doi:10.1093/jnci/87.4.265 PMID:7707417
- Giovannucci EL, Colditz GA, Stampfer MJ *et al.* (1991). The assessment of alcohol consumption by a simple self-administered questionnaire. *Am J Epidemiol*, 133: 810–817. PMID:2021148
- Glynn SA, Albanes D, Pietinen P *et al.* (1996). Alcohol consumption and risk of colorectal cancer in a cohort of Finnish men. *Cancer Causes Control*, 7: 214–223. doi:10.1007/BF00051297 PMID:8740734
- Gold EB, Gordis L, Diener MD *et al.* (1985). Diet and other risk factors for cancer of the pancreas. *Cancer*, 55: 460–467. doi:10.1002/1097-0142(19850115)55:2<460::AID-CNCR2820550229>3.0.CO;2-V PMID:3965101
- Goldbohm RA, Van den Brandt PA, Van 't Veer P *et al.* (1994). Prospective study on alcohol consumption and the risk of cancer of the colon and rectum in the Netherlands. *Cancer Causes Control*, 5: 95–104. doi:10.1007/BF01830255 PMID:8167268
- Gomes ALRR, Guimarães MDL, Gomes CC *et al.* (1995). A case-control study of risk factors for breast cancer in Brazil, 1978–1987. *Int J Epidemiol*, 24: 292–299. doi:10.1093/ije/24.2.292 PMID:7635588
- Goodman MT, Cologne JB, Moriwaki H *et al.* (1997a). Risk factors for primary breast cancer in Japan: 8-year follow-up of atomic bomb survivors. *Prev Med*, 26: 144–153. doi:10.1006/pmed.1996.9979 PMID:9010910
- Goodman MT, Hankin JH, Wilkens LR *et al.* (1997b). Diet, body size, physical activity, and the risk of endometrial cancer. *Cancer Res*, 57: 5077–5085. PMID:9371506
- Goodman MT, Moriwaki H, Vaeth M *et al.* (1995). Prospective cohort study of risk factors for primary liver cancer in Hiroshima and Nagasaki, Japan. *Epidemiology*, 6: 36–41. doi:10.1097/00001648-199501000-00008 PMID:7888442
- Goodman MT, Morgenstern H, Wynder EL (1986). A case-control study of factors affecting the development of renal cell cancer. *Am J Epidemiol*, 124: 926–941. PMID:3776975
- Goodman MT & Tung KH (2003). Active and passive tobacco smoking and the risk of borderline and invasive ovarian cancer (United States). *Cancer Causes Control*, 14: 569–577. doi:10.1023/A:1024828309874 PMID:12948288
- Gordon T & Kannel WB (1984). Drinking and mortality. The Framingham Study. *Am J Epidemiol*, 120: 97–107. PMID:6741928
- Gorini G, Stagnaro E, Fontana V *et al.* (2007). Alcohol consumption and risk of leukemia: A multicenter case-control study. *Leuk Res*, 31: 379–386. doi:10.1016/j.leukres.2006.07.002 PMID:16919329
- Graham S, Dayal H, Swanson M *et al.* (1978). Diet in the epidemiology of cancer of the colon and rectum. *J Natl Cancer Inst*, 61: 709–714. PMID:278848

- Greenfield TK & Rogers JD (1999). Who drinks most of the alcohol in the US? The policy implications. *J Stud Alcohol*, 60: 78–89. PMID:10096312
- Grønbaek M, Becker U, Johansen D *et al.* (1998). Population based cohort study of the association between alcohol intake and cancer of the upper digestive tract. *BMJ*, 317: 844–847. PMID:9748175
- Grönberg H, Damber L, Damber J-E (1996). Total food consumption and body mass index in relation to prostate cancer risk: a case-control study in Sweden with prospectively collected exposure data. *J Urol*, 155: 969–974. doi:10.1016/S0022-5347(01)66360-2 PMID:8583620
- Guénel P, Chastang JF, Luce D *et al.* (1988). A study of the interaction of alcohol drinking and tobacco smoking among French cases of laryngeal cancer. *J Epidemiol Community Health*, 42: 350–354. doi:10.1136/jech.42.4.350 PMID:3256577
- Guénel P, Cyr D, Sabroe S *et al.* (2004). Alcohol drinking may increase risk of breast cancer in men: a European population-based case-control study. *Cancer Causes Control*, 15: 571–580. doi:10.1023/B:CACO.0000036154.18162.43 PMID:15280636
- Guess HA, Friedman GD, Sadler MC *et al.* (1997). 5 α -reductase activity and prostate cancer: a case-control study using stored sera. *Cancer Epidemiol Biomarkers Prev*, 6: 21–24. PMID:8993793
- Gullo L, Pezzilli R, Morselli-Labate AM Italian Pancreatic Cancer Study Group. (1995). Coffee and cancer of the pancreas: an Italian multicenter study. *Pancreas*, 11: 223–229. doi:10.1097/00006676-199510000-00002 PMID:8577674
- Guo W, Blot WJ, Li J-Y *et al.* (1994). A nested case-control study of oesophageal and stomach cancers in the Linxian nutrition intervention trial. *Int J Epidemiol*, 23: 444–450. doi:10.1093/ije/23.3.444 PMID:7960367
- Gwinn ML, Webster LA, Lee NC *et al.* Cancer and Steroid Hormone Study Group. (1986). Alcohol consumption and ovarian cancer risk. *Am J Epidemiol*, 123: 759–766. PMID:3962959
- Gyntelberg F (1973). Physical fitness and coronary heart disease in male residents in Copenhagen aged 40–59. *Dan Med Bull*, 20: 1–4. PMID:4694991
- Haile RW, Witte JS, Ursin G *et al.* (1996). A case-control study of reproductive variables, alcohol, and smoking in premenopausal bilateral breast cancer. *Breast Cancer Res Treat*, 37: 49–56. doi:10.1007/BF01806631 PMID:8750527
- Hakulinen T, Lehtimäki L, Lehtonen M, Teppo L (1974). Cancer morbidity among two male cohorts with increased alcohol consumption in Finland. *J Natl Cancer Inst*, 52: 1711–1714. PMID:4834405
- Hamada GS, Kowalski LP, Nishimoto IN *et al.* São Paulo--Japan Cancer Project Gastric Cancer Study Group. (2002). Risk factors for stomach cancer in Brazil (II): a case-control study among Japanese Brazilians in São Paulo. *Jpn J Clin Oncol*, 32: 284–290. doi:10.1093/jjco/hyf061 PMID:12411565
- Hamajima N, Hirose K, Tajima K *et al.* Collaborative Group on Hormonal Factors in Breast Cancer. (2002). Alcohol, tobacco and breast cancer—collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with

- breast cancer and 95,067 women without the disease. *Br J Cancer*, 87: 1234–1245. doi:10.1038/sj.bjc.6600596 PMID:12439712
- Hansson L-E, Baron J, Nyrén O *et al.* (1994). Tobacco, alcohol and the risk of gastric cancer. A population-based case-control study in Sweden. *Int J Cancer*, 57: 26–31. doi:10.1002/ijc.2910570106 PMID:8150537
- Harnack L, Jacobs DR Jr, Nicodemus K *et al.* (2002). Relationship of folate, vitamin B-6, vitamin B-12, and methionine intake to incidence of colorectal cancers. *Nutr Cancer*, 43: 152–158. doi:10.1207/S15327914NC432_5 PMID:12588695
- Harnack LJ, Anderson KE, Zheng W *et al.* (1997). Smoking, alcohol, coffee, and tea intake and incidence of cancer of the exocrine pancreas: the Iowa Women's Health Study. *Cancer Epidemiol Biomarkers Prev*, 6: 1081–1086. PMID:9419407
- Harris RE, Namboodiri KK, Wynder EL (1992). Breast cancer risk: effects of estrogen replacement therapy and body mass. *J Natl Cancer Inst*, 84: 1575–1582. doi:10.1093/jnci/84.20.1575 PMID:1404451
- Harris RE & Wynder EL (1988). Breast cancer and alcohol consumption. A study in weak associations. *JAMA*, 259: 2867–2871. doi:10.1001/jama.259.19.2867 PMID:3367453
- Harris RW, Brinton LA, Coddell RH *et al.* (1980). Characteristics of women with dysplasia or carcinoma in situ of the cervix uteri. *Br J Cancer*, 42: 359–369. PMID:7426342
- Hartge P, Schiffman MH, Hoover R *et al.* (1989). A case-control study of epithelial ovarian cancer. *Am J Obstet Gynecol*, 161: 10–16. PMID:2750791
- Hashibe M, Boffetta P, Janout V *et al.* (2007c). Esophageal cancer in Central and Eastern Europe: tobacco and alcohol. *Int J Cancer*, 120: 1518–1522. doi:10.1002/ijc.22507 PMID:17205526
- Hashibe M, Boffetta P, Zaridze D *et al.* (2007a). Contribution of tobacco and alcohol to the high rates of squamous cell carcinoma of the supraglottis and glottis in Central Europe. *Am J Epidemiol*, 165: 814–820. doi:10.1093/aje/kwk066 PMID:17244634
- Hashibe M, Brennan P, Benhamou S *et al.* (2007b). Alcohol drinking in never users of tobacco, cigarette smoking in never drinkers, and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *J Natl Cancer Inst*, 99: 777–789. doi:10.1093/jnci/djk179 PMID:17505073
- Hayes RB, Bravo-Otero E, Kleinman DV *et al.* (1999). Tobacco and alcohol use and oral cancer in Puerto Rico. *Cancer Causes Control*, 10: 27–33. doi:10.1023/A:1008876115797 PMID:10334639
- Hayes RB, Brown LM, Schoenberg JB *et al.* (1996). Alcohol use and prostate cancer risk in US blacks and whites. *Am J Epidemiol*, 143: 692–697. PMID:8651231
- Hedberg K, Vaughan TL, White E *et al.* (1994). Alcoholism and cancer of the larynx: a case-control study in western Washington (United States). *Cancer Causes Control*, 5: 3–8. doi:10.1007/BF01830720 PMID:8123776

- Hein HO, Suadicani P, Gyntelberg F (1992). Lung cancer risk and social class. The Copenhagen Male Study—17-year follow up. *Dan Med Bull*, 39: 173–176. PMID:1611922
- Herity B, Moriarty M, Daly L *et al.* (1982). The role of tobacco and alcohol in the aetiology of lung and larynx cancer. *Br J Cancer*, 46: 961–964. PMID:7150489
- Herrero R, Brinton LA, Reeves WC *et al.* (1989). Invasive cervical cancer and smoking in Latin America. *J Natl Cancer Inst*, 81: 205–211. doi:10.1093/jnci/81.3.205 PMID:2536087
- Heuch I, Kvåle G, Jacobsen BK, Bjelke E (1983). Use of alcohol, tobacco and coffee, and risk of pancreatic cancer. *Br J Cancer*, 48: 637–643. PMID:6685527
- Hiatt RA, Armstrong MA, Klatsky AL, Sidney S (1994). Alcohol consumption, smoking, and other risk factors and prostate cancer in a large health plan cohort in California (United States). *Cancer Causes Control*, 5: 66–72. doi:10.1007/BF01830728 PMID:7510134
- Hiatt RA, Klatsky AL, Armstrong MA (1988). Pancreatic cancer, blood glucose and beverage consumption. *Int J Cancer*, 41: 794–797. doi:10.1002/ijc.2910410603 PMID:3372055
- Higginson J (1966). Etiological factors in gastrointestinal cancer in man. *J Natl Cancer Inst*, 37: 527–545. PMID:5923503
- Hirayama T (1989). Association between alcohol consumption and cancer of the sigmoid colon: observations from a Japanese cohort study. *Lancet*, 2: 725–727. doi:10.1016/S0140-6736(89)90782-4 PMID:2570969
- Hirayama T (1992). Life-style and cancer: from epidemiological evidence to public behavior change to mortality reduction of target cancers. *J Natl Cancer Inst Monogr*, 12: 65–74. PMID:1616813
- Hirvonen T, Mennen LI, de Bree A *et al.* (2006). Consumption of antioxidant-rich beverages and risk for breast cancer in French women. *Ann Epidemiol*, 16: 503–508. doi:10.1016/j.annepidem.2005.09.011 PMID:16406814
- Ho JW-C, Lam TH, Tse C-W *et al.* (2004). Smoking, drinking and colorectal cancer in Hong Kong Chinese: a case-control study. *Int J Cancer*, 109: 587–597. doi:10.1002/ijc.20018 PMID:14991582
- Hochberg F, Toniolo P, Cole P, Salcman M (1990). Nonoccupational risk indicators of glioblastoma in adults. *J Neurooncol*, 8: 55–60. doi:10.1007/BF00182087 PMID:2319291
- Hodge AM, English DR, McCredie MRE *et al.* (2004). Foods, nutrients and prostate cancer. *Cancer Causes Control*, 15: 11–20. doi:10.1023/B:CACO.0000016568.25127.10 PMID:14970730
- Holmberg L, Baron JA, Byers T *et al.* (1995). Alcohol intake and breast cancer risk: effect of exposure from 15 years of age. *Cancer Epidemiol Biomarkers Prev*, 4: 843–847. PMID:8634655

- Horn-Ross PL, Canchola AJ, West DW *et al.* (2004). Patterns of alcohol consumption and breast cancer risk in the California Teachers Study cohort. *Cancer Epidemiol Biomarkers Prev*, 13: 405–411. PMID:15006916
- Hoshiyama Y & Sasaba T (1992a). A case-control study of single and multiple stomach cancers in Saitama Prefecture, Japan. *Jpn J Cancer Res*, 83: 937–943. PMID:1429203
- Hoshiyama Y & Sasaba T (1992b). A case-control study of stomach cancer and its relation to diet, cigarettes, and alcohol consumption in Saitama Prefecture, Japan. *Cancer Causes Control*, 3: 441–448. doi:10.1007/BF00051357 PMID:1525325
- Hoshiyama Y, Sekine T, Sasaba T (1993). A case-control study of colorectal cancer and its relation to diet, cigarettes, and alcohol consumption in Saitama Prefecture, Japan. *Tohoku J Exp Med*, 171: 153–165. doi:10.1620/tjem.171.153 PMID:8128484
- Howe HL, Wingo PA, Thun MJ *et al.* (2001). Annual report to the nation on the status of cancer (1973 through 1998), featuring cancers with recent increasing trends. *J Natl Cancer Inst*, 93: 824–842. doi:10.1093/jnci/93.11.824 PMID:11390532
- Høyer AP & Engholm G (1992). Serum lipids and breast cancer risk: a cohort study of 5,207 Danish women. *Cancer Causes Control*, 3: 403–408. doi:10.1007/BF00051352 PMID:1525320
- Hsieh C-C, Thanos A, Mitropoulos D *et al.* (1999). Risk factors for prostate cancer: a case-control study in Greece. *Int J Cancer*, 80: 699–703. doi:10.1002/(SICI)1097-0215(19990301)80:5<699::AID-IJC12>3.0.CO;2-7 PMID:10048970
- Hsing AW, McLaughlin JK, Chow WH *et al.* (1998a). Risk factors for colorectal cancer in a prospective study among U.S. white men. *Int J Cancer*, 77: 549–553. doi:10.1002/(SICI)1097-0215(19980812)77:4<549::AID-IJC13>3.0.CO;2-1 PMID:9679757
- Hsing AW, McLaughlin JK, Cocco P *et al.* (1998b). Risk factors for male breast cancer (United States). *Cancer Causes Control*, 9: 269–275. doi:10.1023/A:1008869003012 PMID:9684707
- Hsing AW, McLaughlin JK, Schuman LM *et al.* (1990). Diet, tobacco use, and fatal prostate cancer: results from the Lutheran Brotherhood Cohort Study. *Cancer Res*, 50: 6836–6840. PMID:2208150
- Hsu CC, Chow W-H, Boffetta P *et al.* (2007). Dietary risk factors for kidney cancer in Eastern and Central Europe. *Am J Epidemiol*, 166: 62–70. doi:10.1093/aje/kwm043 PMID:17456477
- Hu J, Johnson KC, Mao Y *et al.* (1998). Risk factors for glioma in adults: a case-control study in northeast China. *Cancer Detect Prev*, 22: 100–108. doi:10.1046/j.1525-1500.1998.CDOA22.x PMID:9544430
- Hu J, La Vecchia C, Negri E *et al.* (1999). Diet and brain cancer in adults: a case-control study in northeast China. *Int J Cancer*, 81: 20–23. doi:10.1002/(SICI)1097-0215(19990331)81:1<20::AID-IJC4>3.0.CO;2-2 PMID:10077146
- Hu J, Mao Y, Dryer D, White K Canadian Cancer Registries Epidemiology Research Group. (2002). Risk factors for lung cancer among Canadian women who have never smoked. *Cancer Detect Prev*, 26: 129–138. doi:10.1016/S0361-090X(02)00038-7 PMID:12102147

- Hu J, Mao Y, Ugnat A-M, Jinfu Hu, Yang Mao, Anne-Marie Ugna. (2000). Parental cigarette smoking, hard liquor consumption and the risk of childhood brain tumors—a case-control study in northeast China. *Acta Oncol*, 39: 979–984. doi:10.1080/02841860050215972 PMID:11207006
- Hu J, Mao Y, White K Canadian Cancer Registries Epidemiology Research Group. (2003). Diet and vitamin or mineral supplements and risk of renal cell carcinoma in Canada. *Cancer Causes Control*, 14: 705–714. doi:10.1023/A:1026310323882 PMID:14674734
- Hu JF, Liu YY, Yu YK *et al.* (1991). Diet and cancer of the colon and rectum: a case-control study in China. *Int J Epidemiol*, 20: 362–367. doi:10.1093/ije/20.2.362 PMID:1917235
- Hu JF, Wang GQ, Jia EM *et al.* (1989). Risk analysis of fuzzy states in data of case-control study for stomach cancer in Heilongjiang Province *Zhonghua Zhong Liu Za Zhi*, 11: 28–30. PMID:2776643
- Huang KF, Li H, Li HQ (2005). Case-control study for risk factors of esophageal cancer in Shandong province. *Chinese J Med Writing*, 12: 385–387.
- Huang WY, Newman B, Millikan RC *et al.* (2000). Hormone-related factors and risk of breast cancer in relation to estrogen receptor and progesterone receptor status. *Am J Epidemiol*, 151: 703–714. PMID:10752798
- Huang WY, Winn DM, Brown LM *et al.* (2003). Alcohol concentration and risk of oral cancer in Puerto Rico. *Am J Epidemiol*, 157: 881–887. doi:10.1093/aje/kwg055 PMID:12746240
- Huang X-H, Chen S-D, Wang G-G *et al.* (2004). Study on the impact of GSTM1 polymorphisms on the risk of histologic types of lung cancer — A case-control study. *J Public Health Prev Med*, 15: 24–26.
- Hurley SF, McNeil JJ, Donnan GA *et al.* (1996). Tobacco smoking and alcohol consumption as risk factors for glioma: a case-control study in Melbourne, Australia. *J Epidemiol Community Health*, 50: 442–446. doi:10.1136/jech.50.4.442 PMID:8882229
- IARC. (1988). Alcohol Drinking. *IARC Monogr Eval Carcinog Risks Hum*, 44: 1–378. PMID:3236394
- IARC. (2004). Tobacco smoke and involuntary smoking. *IARC Monogr Eval Carcinog Risks Hum*, 83: 1–1438. PMID:15285078
- Infante-Rivard C, Krajcinovic M, Labuda D, Sinnett D (2002). Childhood acute lymphoblastic leukemia associated with parental alcohol consumption and polymorphisms of carcinogen-metabolizing genes. *Epidemiology*, 13: 277–281. doi:10.1097/00001648-200205000-00007 PMID:11964928
- Inoue M, Tajima K, Hirose K *et al.* (1994). Life-style and subsite of gastric cancer—joint effect of smoking and drinking habits. *Int J Cancer*, 56: 494–499. doi:10.1002/ijc.2910560407 PMID:8112885
- Inoue M, Tajima K, Takezaki T *et al.* (2003). Epidemiology of pancreatic cancer in Japan: a nested case-control study from the Hospital-based Epidemiologic

- Research Program at Aichi Cancer Center (HERPACC). *Int J Epidemiol*, 32: 257–262. doi:10.1093/ije/dyg062 PMID:12714546
- Iribarren C, Haselkorn T, Tekawa IS, Friedman GD (2001). Cohort study of thyroid cancer in a San Francisco Bay area population. *Int J Cancer*, 93: 745–750. doi:10.1002/ijc.1377 PMID:11477590
- Isaksson B, Jonsson F, Pedersen NL *et al.* (2002). Lifestyle factors and pancreatic cancer risk: a cohort study from the Swedish Twin Registry. *Int J Cancer*, 98: 480–482. doi:10.1002/ijc.10256 PMID:11920604
- Iscovich JM, Iscovich RB, Howe G *et al.* (1989). A case-control study of diet and breast cancer in Argentina. *Int J Cancer*, 44: 770–776. doi:10.1002/ijc.2910440504 PMID:2583858
- Jackson MA, Kovi J, Heshmat MY *et al.* (1981). Factors involved in the high incidence of prostatic cancer among American blacks. *Prog Clin Biol Res*, 53: 111–132. PMID:7465581
- Jain M, Howe GR, St Louis P, Miller AB (1991). Coffee and alcohol as determinants of risk of pancreas cancer: a case-control study from Toronto. *Int J Cancer*, 47: 384–389. doi:10.1002/ijc.2910470313 PMID:1993545
- Jain MG, Ferrenc RG, Rehm JT *et al.* (2000a). Alcohol and breast cancer mortality in a cohort study. *Breast Cancer Res Treat*, 64: 201–209. doi:10.1023/A:1006402323445 PMID:11194456
- Jain MG, Hislop GT, Howe GR *et al.* (1998). Alcohol and other beverage use and prostate cancer risk among Canadian men. *Int J Cancer*, 78: 707–711. doi:10.1002/(SICI)1097-0215(19981209)78:6<707::AID-IJC7>3.0.CO;2-2 PMID:9833763
- Jain MG, Howe GR, Rohan TE (2000c). Nutritional factors and endometrial cancer in Ontario, Canada. *Cancer Control*, 7: 288–296. PMID:10832115
- Jain MG, Rohan TE, Howe GR, Miller AB (2000b). A cohort study of nutritional factors and endometrial cancer. *Eur J Epidemiol*, 16: 899–905. doi:10.1023/A:1011012621990 PMID:11338120
- Jedrychowski W, Boeing H, Wahrendorf J *et al.* (1993). Vodka consumption, tobacco smoking and risk of gastric cancer in Poland. *Int J Epidemiol*, 22: 606–613. doi:10.1093/ije/22.4.606 PMID:8225732
- Jensen OM (1979). Cancer morbidity and causes of death among Danish brewery workers. *Int J Cancer*, 23: 454–463. doi:10.1002/ijc.2910230404 PMID:437924
- Jensen OM (1980) *Cancer morbidity and causes of death among Danish brewery workers*. 1st Edition, International Agency for Research on Cancer (ed), Lyon.
- Ji BT, Chow WH, Dai Q *et al.* (1995). Cigarette smoking and alcohol consumption and the risk of pancreatic cancer: a case-control study in Shanghai, China. *Cancer Causes Control*, 6: 369–376. doi:10.1007/BF00051413 PMID:7548725
- Ji B-T, Chow WH, Yang G *et al.* (1996). The influence of cigarette smoking, alcohol, and green tea consumption on the risk of carcinoma of the cardia and distal stomach in Shanghai, China. *Cancer*, 77: 2449–2457. doi:10.1002/(SICI)1097-0142(19960615)77:12<2449::AID-CNCR6>3.0.CO;2-H PMID:8640692

- Ji B-T, Dai Q, Gao YT *et al.* (2002). Cigarette and alcohol consumption and the risk of colorectal cancer in Shanghai, China. *Eur J Cancer Prev*, 11: 237–244. doi:10.1097/00008469-200206000-00007 PMID:12131657
- Johnson KC, Pan S, Mao Y Canadian Cancer Registries Epidemiology Research Group. (2002). Risk factors for male breast cancer in Canada, 1994–1998. *Eur J Cancer Prev*, 11: 253–263. doi:10.1097/00008469-200206000-00009 PMID:12131659
- Kaaks R, Slimani N, Riboli E (1997). Pilot phase studies on the accuracy of dietary intake measurements in the EPIC project: overall evaluation of results. European Prospective Investigation into Cancer and Nutrition. *Int J Epidemiol*, 26: Suppl 1S26–S36. doi:10.1093/ije/26.suppl_1.S26 PMID:9126531
- Kabat GC, Chang CJ, Wynder EL (1994). The role of tobacco, alcohol use, and body mass index in oral and pharyngeal cancer. *Int J Epidemiol*, 23: 1137–1144. doi:10.1093/ije/23.6.1137 PMID:7721514
- Kabat GC, Howson CP, Wynder EL (1986). Beer consumption and rectal cancer. *Int J Epidemiol*, 15: 494–501. doi:10.1093/ije/15.4.494 PMID:3818156
- Kabat GC, Ng SKC, Wynder EL (1993). Tobacco, alcohol intake, and diet in relation to adenocarcinoma of the esophagus and gastric cardia. *Cancer Causes Control*, 4: 123–132. doi:10.1007/BF00053153 PMID:8481491
- Kabat GC & Wynder EL (1984). Lung cancer in nonsmokers. *Cancer*, 53: 1214–1221. doi:10.1002/1097-0142(19840301)53:5<1214::AID-CNCR2820530532>3.0.CO;2-8 PMID:6692309
- Kadamani S, Asal NR, Nelson RY (1989). Occupational hydrocarbon exposure and risk of renal cell carcinoma. *Am J Ind Med*, 15: 131–141. doi:10.1002/ajim.4700150202 PMID:2729279
- Kalandidi A, Tzonou A, Lipworth L *et al.* (1996). A case-control study of endometrial cancer in relation to reproductive, somatometric, and life-style variables. *Oncology*, 53: 354–359. doi:10.1159/000227587 PMID:8784467
- Kalapothiski V, Tzonou A, Hsieh CC *et al.* (1993). Tobacco, ethanol, coffee, pancreatitis, diabetes mellitus, and cholelithiasis as risk factors for pancreatic carcinoma. *Cancer Causes Control*, 4: 375–382. doi:10.1007/BF00051341 PMID:8347787
- Karlson B-M, Ekblom A, Josefsson S *et al.* (1997). The risk of pancreatic cancer following pancreatitis: an association due to confounding? *Gastroenterology*, 113: 587–592. doi:10.1053/gast.1997.v113.pm9247480 PMID:9247480
- Kato I, Miura S, Kasumi F *et al.* (1992d). A case-control study of breast cancer among Japanese women: with special reference to family history and reproductive and dietary factors. *Breast Cancer Res Treat*, 24: 51–59. doi:10.1007/BF01832358 PMID:1463872
- Kato I, Nomura AMY, Stemmermann GN, Chyou PH (1992c). Prospective study of the association of alcohol with cancer of the upper aerodigestive tract and other sites. *Cancer Causes Control*, 3: 145–151. doi:10.1007/BF00051654 PMID:1562704
- Kato I, Tominaga S, Ito Y *et al.* (1992a). A prospective study of atrophic gastritis and stomach cancer risk. *Jpn J Cancer Res*, 83: 1137–1142. PMID:1483928

- Kato I, Tominaga S, Matsumoto K (1992b). A prospective study of stomach cancer among a rural Japanese population: a 6-year survey. *Jpn J Cancer Res*, 83: 568–575. PMID:1644660
- Kato I, Tominaga S, Terao C (1989). Alcohol consumption and cancers of hormone-related organs in females. *Jpn J Clin Oncol*, 19: 202–207. PMID:2810820
- Kelemen LE, Sellers TA, Vierkant RA *et al.* (2004). Association of folate and alcohol with risk of ovarian cancer in a prospective study of postmenopausal women. *Cancer Causes Control*, 15: 1085–1093. doi:10.1007/s10552-004-1546-6 PMID:15801492
- Keller AZ (1967). Demographic, clinical and survivorship characteristics of males with primary cancer of the breast. *Am J Epidemiol*, 85: 183–199. PMID:6020792
- Kikuchi S, Nakajima T, Kobayashi O *et al.* (2002). U-shaped effect of drinking and linear effect of smoking on risk for stomach cancer in Japan. *Jpn J Cancer Res*, 93: 953–959. PMID:12359047
- Kim D-H, Ahn Y-O, Lee B-H *et al.* (2004). Methylenetetrahydrofolate reductase polymorphism, alcohol intake, and risks of colon and rectal cancers in Korea. *Cancer Lett*, 216: 199–205. doi:10.1016/j.canlet.2004.08.014 PMID:15533596
- Kinjo Y, Cui Y, Akiba S *et al.* (1998). Mortality risks of oesophageal cancer associated with hot tea, alcohol, tobacco and diet in Japan. *J Epidemiol*, 8: 235–243. PMID:9816815
- Kinney AY, Millikan RC, Lin YH *et al.* (2000). Alcohol consumption and breast cancer among black and white women in North Carolina (United States). *Cancer Causes Control*, 11: 345–357. doi:10.1023/A:1008973709917 PMID:10843445
- Kirkpatrick CS, White E, Lee JAH (1994). Case-control study of malignant melanoma in Washington State. II. Diet, alcohol, and obesity. *Am J Epidemiol*, 139: 869–880. PMID:8166137
- Kjaerheim K & Andersen A (1994). Cancer incidence among waitresses in Norway. *Cancer Causes Control*, 5: 31–37. doi:10.1007/BF01830724 PMID:8123777
- Kjaerheim K, Andersen A, Helseth A (1993). Alcohol abstainers: a low-risk group for cancer—a cohort study of Norwegian teetotalers. *Cancer Epidemiol Biomarkers Prev*, 2: 93–97. PMID:8467252
- Kjaerheim K, Gaard M, Andersen A (1998). The role of alcohol, tobacco, and dietary factors in upper aerogastric tract cancers: a prospective study of 10,900 Norwegian men. *Cancer Causes Control*, 9: 99–108. doi:10.1023/A:1008809706062 PMID:9486469
- Klatsky AL, Armstrong MA, Friedman GD, Hiatt RA (1988). The relations of alcoholic beverage use to colon and rectal cancer. *Am J Epidemiol*, 128: 1007–1015. PMID:3189277
- Klatsky AL, Friedman GD, Siegelau AB (1981). Alcohol and mortality. A ten-year Kaiser-Permanente experience. *Ann Intern Med*, 95: 139–145. PMID:7258861
- Kneller RW, McLaughlin JK, Bjelke E *et al.* (1991). A cohort study of stomach cancer in a high-risk American population. *Cancer*, 68: 672–678. doi:10.1002/1097-0142(19910801)68:3<672::AID-CNCR2820680339>3.0.CO;2-T PMID:2065291

- Kolonel LN, Hankin JH, Wilkens LR *et al.* (1990). An epidemiologic study of thyroid cancer in Hawaii. *Cancer Causes Control*, 1: 223–234. doi:10.1007/BF00117474 PMID:2102295
- Kono S, Ikeda M, Tokudome S *et al.* (1985). Alcohol and cancer in male Japanese physicians. *J Cancer Res Clin Oncol*, 109: 82–85. doi:10.1007/BF01884260 PMID:3972887
- Kono S, Ikeda M, Tokudome S *et al.* (1986). Alcohol and mortality: a cohort study of male Japanese physicians. *Int J Epidemiol*, 15: 527–532. doi:10.1093/ije/15.4.527 PMID:3818161
- Kono S, Ikeda M, Tokudome S *et al.* (1987). Cigarette smoking, alcohol and cancer mortality: a cohort study of male Japanese physicians. *Jpn J Cancer Res*, 78: 1323–1328. PMID:3123436
- Koo LC (1988). Dietary habits and lung cancer risk among Chinese females in Hong Kong who never smoked. *Nutr Cancer*, 11: 155–172. doi:10.1080/01635588809513983 PMID:2841651
- Korte JE, Brennan P, Henley SJ, Boffetta P (2002). Dose-specific meta-analysis and sensitivity analysis of the relation between alcohol consumption and lung cancer risk. *Am J Epidemiol*, 155: 496–506. doi:10.1093/aje/155.6.496 PMID:11882523
- Kramer S, Ward E, Meadows AT, Malone KE (1987). Medical and drug risk factors associated with neuroblastoma: a case-control study. *J Natl Cancer Inst*, 78: 797–804. PMID:3471992
- Kreiger N, Marrett LD, Dodds L *et al.* (1993). Risk factors for renal cell carcinoma: results of a population-based case-control study. *Cancer Causes Control*, 4: 101–110. doi:10.1007/BF00053150 PMID:8481488
- Kune S, Kune GA, Watson LF (1987). Case-control study of alcoholic beverages as etiological factors: the Melbourne Colorectal Cancer Study. *Nutr Cancer*, 9: 43–56. doi:10.1080/01635588709513909 PMID:3808969
- Kunze E, Claude J, Frentzel-Beyme R *et al.* (1986). Association of cancer of the lower urinary tract with consumption of alcoholic beverages. A case-control study. *Carcinogenesis*, 7: 163–165. doi:10.1093/carcin/7.1.163 PMID:3943138
- Kuper H, Titus-Ernstoff L, Harlow BL, Cramer DW (2000b). Population based study of coffee, alcohol and tobacco use and risk of ovarian cancer. *Int J Cancer*, 88: 313–318. doi:10.1002/1097-0215(20001015)88:2<313::AID-IJC26>3.0.CO;2-5 PMID:11004686
- Kuper H, Tzonou A, Kaklamani E *et al.* (2000a). Tobacco smoking, alcohol consumption and their interaction in the causation of hepatocellular carcinoma. *Int J Cancer*, 85: 498–502. doi:10.1002/(SICI)1097-0215(20000215)85:4<498::AID-IJC9>3.0.CO;2-F PMID:10699921
- Kuper H, Ye W, Weiderpass E *et al.* (2000c). Alcohol and breast cancer risk: the alcoholism paradox. *Br J Cancer*, 83: 949–951. doi:10.1054/bjoc.2000.1360 PMID:10970699

- Kushi LH, Mink PJ, Folsom AR *et al.* (1999). Prospective study of diet and ovarian cancer. *Am J Epidemiol*, 149: 21–31. PMID:9883790
- Kvåle G, Bjelke E, Gart JJ (1983). Dietary habits and lung cancer risk. *Int J Cancer*, 31: 397–405. doi:10.1002/ijc.2910310402 PMID:6832851
- La Vecchia C, Decarli A, Fasoli M, Gentile A (1986). Nutrition and diet in the etiology of endometrial cancer. *Cancer*, 57: 1248–1253. doi:10.1002/1097-0142(19860315)57:6<1248::AID-CNCR2820570631>3.0.CO;2-V PMID:3002600
- La Vecchia C, Lucchini F, Franceschi S *et al.* (2000). Trends in mortality from primary liver cancer in Europe. *Eur J Cancer*, 36: 909–915. doi:10.1016/S0959-8049(00)00052-6 PMID:10785597
- La Vecchia C, Negri E, Cavalieri d'Oro L, Franceschi S (1998). Liver cirrhosis and the risk of primary liver cancer. *Eur J Cancer Prev*, 7: 315–320. doi:10.1097/00008469-199808000-00007 PMID:9806120
- La Vecchia C, Negri E, Franceschi S *et al.* (1992). Alcohol and epithelial ovarian cancer. *J Clin Epidemiol*, 45: 1025–1030. doi:10.1016/0895-4356(92)90119-8 PMID:1432017
- La Vecchia C, Negri E, Franceschi S, D'Avanzo B (1993). Moderate beer consumption and the risk of colorectal cancer. *Nutr Cancer*, 19: 303–306. doi:10.1080/01635589309514260 PMID:8346078
- Lagergren J, Bergström R, Lindgren A, Nyrén O (2000). The role of tobacco, snuff and alcohol use in the aetiology of cancer of the oesophagus and gastric cardia. *Int J Cancer*, 85: 340–346. doi:10.1002/(SICI)1097-0215(20000201)85:3<340::AID-IJC8>3.0.CO;2-N PMID:10652424
- Lagergren J, Viklund P, Jansson C (2006). Carbonated soft drinks and risk of esophageal adenocarcinoma: a population-based case-control study. *J Natl Cancer Inst*, 98: 1158–1161. doi:10.1093/jnci/djj310 PMID:16912268
- Lagiou P, Ye W, Wedrén S *et al.* (2001). Incidence of ovarian cancer among alcoholic women: a cohort study in Sweden. *Int J Cancer*, 91: 264–266. doi:10.1002/1097-0215(200002)9999:9999<::AID-IJC1027>3.3.CO;2-B PMID:11146456
- Land CE, Hayakawa N, Machado SG *et al.* (1994). A case-control interview study of breast cancer among Japanese A-bomb survivors. II. Interactions with radiation dose. *Cancer Causes Control*, 5: 167–176. doi:10.1007/BF01830263 PMID:8167264
- Larsson SC, Giovannucci E, Wolk A (2007). Alcoholic beverage consumption and gastric cancer risk: a prospective population-based study in women. *Int J Cancer*, 120: 373–377. doi:10.1002/ijc.22204 PMID:17066442
- Le Marchand L, Kolonel LN, Wilkens LR *et al.* (1994). Animal fat consumption and prostate cancer: a prospective study in Hawaii. *Epidemiology*, 5: 276–282. doi:10.1097/00001648-199405000-00004 PMID:8038241
- Le Marchand L, Saltzman BS, Hankin JH *et al.* (2006). Sun exposure, diet, and melanoma in Hawaii Caucasians. *Am J Epidemiol*, 164: 232–245. doi:10.1093/aje/kwj115 PMID:16524953

- Le Marchand L, Wilkens LR, Kolonel LN *et al.* (1997). Associations of sedentary lifestyle, obesity, smoking, alcohol use, and diabetes with the risk of colorectal cancer. *Cancer Res*, 57: 4787–4794. PMID:9354440
- Lee DH, Anderson KE, Harnack LJ *et al.* (2004). Heme iron, zinc, alcohol consumption, and colon cancer: Iowa Women's Health Study. *J Natl Cancer Inst*, 96: 403–407. doi:10.1093/jnci/djh047 PMID:14996862
- Lee H-H, Wu H-Y, Chuang Y-C *et al.* (1990). Epidemiologic characteristics and multiple risk factors of stomach cancer in Taiwan. *Anticancer Res*, 10: 875–881. PMID:2382983
- Lee JE, Giovannucci E, Smith-Warner SA *et al.* (2006). Total fluid intake and use of individual beverages and risk of renal cell cancer in two large cohorts. *Cancer Epidemiol Biomarkers Prev*, 15: 1204–1211. doi:10.1158/1055-9965.EPI-05-0889 PMID:16775182
- Lee JE, Hunter DJ, Spiegelman D *et al.* (2007). Alcohol intake and renal cell cancer in a pooled analysis of 12 prospective studies. *J Natl Cancer Inst*, 99: 801–810. doi:10.1093/jnci/djk181 PMID:17505075
- Lee K-W, Kuo W-R, Tsai S-M *et al.* (2005). Different impact from betel quid, alcohol and cigarette: risk factors for pharyngeal and laryngeal cancer. *Int J Cancer*, 117: 831–836. doi:10.1002/ijc.21237 PMID:15957167
- Lee M, Wrensch M, Miike R (1997). Dietary and tobacco risk factors for adult onset glioma in the San Francisco Bay Area (California, USA) *Cancer Causes Control*, 8: 13–24. doi:10.1023/A:1018470802969 PMID:9051318
- Lenz SK, Goldberg MS, Labrèche F *et al.* (2002). Association between alcohol consumption and postmenopausal breast cancer: results of a case-control study in Montreal, Quebec, Canada. *Cancer Causes Control*, 13: 701–710. doi:10.1023/A:1020296905208 PMID:12420948
- Levi F, Franceschi S, Negri E, La Vecchia C (1993). Dietary factors and the risk of endometrial cancer. *Cancer*, 71: 3575–3581. doi:10.1002/1097-0142(19930601)71:11<3575::AID-CNCR2820711119>3.0.CO;2-0 PMID:8490907
- Levi F, Pasche C, Lucchini F *et al.* (2000). Food groups and esophageal cancer risk in Vaud, Switzerland. *Eur J Cancer Prev*, 9: 257–263. doi:10.1097/00008469-200008000-00005 PMID:10958328
- Li CI, Malone KE, Porter PL *et al.* (2003). The relationship between alcohol use and risk of breast cancer by histology and hormone receptor status among women 65–79 years of age. *Cancer Epidemiol Biomarkers Prev*, 12: 1061–1066. PMID:14578143
- Li K, Yu P, Zhang ZX *et al.* (2001). Food components and risk of esophageal cancer in Chaoshan region of China, a high-risk area of esophageal cancer. *Chinese J Cancer*, 20: 160–163.
- Licciardone JC, Wilkins JR 3rd, Brownson RC, Chang JC (1989). Cigarette smoking and alcohol consumption in the aetiology of uterine cervical cancer. *Int J Epidemiol*, 18: 533–537. doi:10.1093/ije/18.3.533 PMID:2807654

- Lim U, Weinstein S, Albanes D *et al.* (2006). Dietary factors of one-carbon metabolism in relation to non-Hodgkin lymphoma and multiple myeloma in a cohort of male smokers. *Cancer Epidemiol Biomarkers Prev*, 15: 1109–1114. doi:10.1158/1055-9965.EPI-05-0918 PMID:16775167
- Lin Y, Kikuchi S, Tamakoshi K *et al.* Japan Collaborative Group. (2005). Prospective study of alcohol consumption and breast cancer risk in Japanese women. *Int J Cancer*, 116: 779–783. doi:10.1002/ijc.20980 PMID:15838830
- Lin Y, Tamakoshi A, Kawamura T *et al.* JACC Study Group. (2002). Risk of pancreatic cancer in relation to alcohol drinking, coffee consumption and medical history: findings from the Japan collaborative cohort study for evaluation of cancer risk. *Int J Cancer*, 99: 742–746. doi:10.1002/ijc.10402 PMID:12115510
- Lindblad M, Rodríguez LAG, Lagergren J (2005). Body mass, tobacco and alcohol and risk of esophageal, gastric cardia, and gastric non-cardia adenocarcinoma among men and women in a nested case-control study. *Cancer Causes Control*, 16: 285–294. doi:10.1007/s10552-004-3485-7 PMID:15947880
- Lindblad P, Wolk A, Bergström R, Adami H-O (1997). Diet and risk of renal cell cancer: a population-based case-control study. *Cancer Epidemiol Biomarkers Prev*, 6: 215–223. PMID:9107425
- Linet MS, Harlow SD, McLaughlin JK (1987). A case-control study of multiple myeloma in whites: chronic antigenic stimulation, occupation, and drug use. *Cancer Res*, 47: 2978–2981. PMID:3567914
- Liu XM, Wang QS, Zhang YL, Men ZH (2000). Case control study of smoking and drinking associated with male esophageal cancer. *JTMU*, 6: 280–281.
- Llewellyn CD, Johnson NW, Warnakulasuriya KA (2004a). Risk factors for oral cancer in newly diagnosed patients aged 45 years and younger: a case-control study in Southern England. *J Oral Pathol Med*, 33: 525–532. doi:10.1111/j.1600-0714.2004.00222.x PMID:15357672
- Llewellyn CD, Linklater K, Bell J *et al.* (2004b). An analysis of risk factors for oral cancer in young people: a case-control study. *Oral Oncol*, 40: 304–313. doi:10.1016/j.oraloncology.2003.08.015 PMID:14747062
- Loerbroeks A, Schouten LJ, Goldbohm RA, van den Brandt PA (2007). Alcohol consumption, cigarette smoking, and endometrial cancer risk: results from the Netherlands Cohort Study. *Cancer Causes Control*, 18: 551–560. doi:10.1007/s10552-007-0127-x PMID:17437180
- Longnecker MP (1990). A case-control study of alcoholic beverage consumption in relation to risk of cancer of the right colon and rectum in men. *Cancer Causes Control*, 1: 5–14. doi:10.1007/BF00053178 PMID:2102276
- Longnecker MP (1994). Alcoholic beverage consumption in relation to risk of breast cancer: meta-analysis and review. *Cancer Causes Control*, 5: 73–82. doi:10.1007/BF01830729 PMID:8123780

- Longnecker MP, Newcomb PA, Mittendorf R *et al.* (1995). Risk of breast cancer in relation to lifetime alcohol consumption. *J Natl Cancer Inst*, 87: 923–929. doi:10.1093/jnci/87.12.923 PMID:7666482
- Longnecker MP, Orza MJ, Adams ME *et al.* (1990). A meta-analysis of alcoholic beverage consumption in relation to risk of colorectal cancer. *Cancer Causes Control*, 1: 59–68. doi:10.1007/BF00053184 PMID:2151680
- López-Carrillo L, López-Cervantes M, Ramírez-Espitia A *et al.* (1998). Alcohol consumption and gastric cancer in Mexico. *Cad Saude Publica*, 14: Suppl 325–32. doi:10.1590/S0102-311X1998000700004 PMID:9819462
- Lu C-M, Chung M-C, Huang C-H, Ko Y-C (2005). Interaction effect in bladder cancer between N-acetyltransferase 2 genotype and alcohol drinking. *Urol Int*, 75: 360–364. doi:10.1159/000089175 PMID:16327307
- Lu JB, Lian SY, Sun XB *et al.* (2000b). A case-control study on the risk factors of esophageal cancer in Linzhou *Zhonghua Liu Xing Bing Xue Za Zhi*, 21: 434–436. PMID:11860829
- Lu Q-J, Yao S-Y, Huang C-Y *et al.* (2000a). [The cohort study on intake of alcohol and lung cancer risk in the Yunnan Tin Corporation (YTC) miners.] *China Pub Health*, 16: 707–708.
- Lu Y, Xu YC, Shen J *et al.* (2006). Study on the association between the role of polymorphisms of the O⁶-methylguanine-DNA methyltransferase gene and gastric cancer hereditary susceptibility. *Chin J Dis Control Prev*, 10: 222–225.
- Lucas FL, Cauley JA, Stone RA *et al.* Study of Osteoporotic Fractures Research Group. (1998). Bone mineral density and risk of breast cancer: differences by family history of breast cancer. *Am J Epidemiol*, 148: 22–29. PMID:9663400
- Lumey LH, Pittman B, Wynder EL (1998). Alcohol use and prostate cancer in U.S. whites: no association in a confirmatory study. *Prostate*, 36: 250–255. doi:10.1002/(SICI)1097-0045(19980901)36:4<250::AID-PROS6>3.0.CO;2-J PMID:9719025
- Lund Nilssen TI, Johnsen R, Vatten LJ (2000). Socio-economic and lifestyle factors associated with the risk of prostate cancer. *Br J Cancer*, 82: 1358–1363. PMID:10755415
- Luo R-H, Zhao Z-X, Zhou X-Y *et al.* (2005). Risk factors for primary liver carcinoma in Chinese population. *World J Gastroenterol*, 11: 4431–4434. PMID:16038048
- Luo SY (2005). Case-control study on stomach cancer. *China Community Med*, 11: 4
- Lyon JL, Mahoney AW, French TK, Moser R Jr (1992). Coffee consumption and the risk of cancer of the exocrine pancreas: a case-control study in a low-risk population. *Epidemiology*, 3: 164–170. doi:10.1097/00001648-199203000-00015 PMID:1576222
- Ma H, Bernstein L, Ross RK, Ursin G (2006). Hormone-related risk factors for breast cancer in women under age 50 years by estrogen and progesterone receptor status: results from a case-control and a case-case comparison. *Breast Cancer Res*, 8: R39 doi:10.1186/bcr1514 PMID:16846528
- Mabuchi K, Bross DS, Kessler II (1985a). Risk factors for male breast cancer. *J Natl Cancer Inst*, 74: 371–375. PMID:3856050

- Mabuchi K, Bross DS, Kessler II (1985b). Epidemiology of cancer of the vulva. A case-control study. *Cancer*, 55: 1843–1848. doi:10.1002/1097-0142(19850415)55:8<1843::AID-CNCR2820550833>3.0.CO;2-M PMID:3978570
- Mack TM, Yu MC, Hanisch R, Henderson BE (1986). Pancreas cancer and smoking, beverage consumption, and past medical history. *J Natl Cancer Inst*, 76: 49–60. PMID:3455742
- Mack WJ, Preston-Martin S, Dal Maso L *et al.* (2003). A pooled analysis of case-control studies of thyroid cancer: cigarette smoking and consumption of alcohol, coffee, and tea. *Cancer Causes Control*, 14: 773–785. doi:10.1023/A:1026349702909 PMID:14674742
- Maclure M & Willett W (1990). A case-control study of diet and risk of renal adenocarcinoma. *Epidemiology*, 1: 430–440. doi:10.1097/00001648-199011000-00004 PMID:2090280
- MacMahon B, Yen S, Trichopoulos D *et al.* (1981). Coffee and cancer of the pancreas. *N Engl J Med*, 304: 630–633. doi:10.1056/NEJM198103123041102 PMID:7453739
- Mahabir S, Leitzmann MF, Virtanen MJ *et al.* (2005). Prospective study of alcohol drinking and renal cell cancer risk in a cohort of Finnish male smokers. *Cancer Epidemiol Biomarkers Prev*, 14: 170–175. PMID:15668492
- Maier H, Dietz A, Gewelke U *et al.* (1992a). Tobacco and alcohol and the risk of head and neck cancer. *Clin Invest*, 70: 320–327. doi:10.1007/BF00184668 PMID:1521046
- Maier H, Gewelke U, Dietz A, Heller W-D (1992b). Risk factors of cancer of the larynx: results of the Heidelberg case-control study. *Otolaryngol Head Neck Surg*, 107: 577–582. PMID:1437190
- Maier H, Sennewald E, Heller GF, Weidauer H (1994). Chronic alcohol consumption—the key risk factor for pharyngeal cancer. *Otolaryngol Head Neck Surg*, 110: 168–173. PMID:7906410
- Männistö S, Virtanen M, Kataja V *et al.* (2000). Lifetime alcohol consumption and breast cancer: a case-control study in Finland. *Public Health Nutr*, 3: 11–18. doi:10.1017/S1368980000000033 PMID:10786719
- Manousos O, Day NE, Trichopoulos D *et al.* (1983). Diet and colorectal cancer: a case-control study in Greece. *Int J Cancer*, 32: 1–5. doi:10.1002/ijc.2910320102 PMID:6862688
- Manousos O, Trichopoulos D, Koutselinis A *et al.* (1981). Epidemiologic characteristics and trace elements in pancreatic cancer in Greece. *Cancer Detect Prev*, 4: 439–442. PMID:7349806
- Marcus PM, Newman B, Millikan RC *et al.* (2000). The associations of adolescent cigarette smoking, alcoholic beverage consumption, environmental tobacco smoke, and ionizing radiation with subsequent breast cancer risk (United States). *Cancer Causes Control*, 11: 271–278. doi:10.1023/A:1008911902994 PMID:10782661
- Marrero JA, Fontana RJ, Fu S *et al.* (2005). Alcohol, tobacco and obesity are synergistic risk factors for hepatocellular carcinoma. *J Hepatol*, 42: 218–224. doi:10.1016/j.jhep.2004.10.005 PMID:15664247

- Marshall JR, Graham S, Byers T *et al.* (1983). Diet and smoking in the epidemiology of cancer of the cervix. *J Natl Cancer Inst*, 70: 847–851. PMID:6573528
- Marshall JR, Graham S, Haughey BP *et al.* (1992). Smoking, alcohol, dentition and diet in the epidemiology of oral cancer. *Eur J Cancer B Oral Oncol*, 28B: 9–15. doi:10.1016/0964-1955(92)90005-L PMID:1422474
- Martin PMD & Hill GB (1984). Cervical cancer in relation to tobacco and alcohol consumption in Lesotho, southern Africa. *Cancer Detect Prev*, 7: 109–115. PMID:6713445
- Martin-Moreno JM, Boyle P, Gorgojo L *et al.* (1993). Alcoholic beverage consumption and risk of breast cancer in Spain. *Cancer Causes Control*, 4: 345–353. doi:10.1007/BF00051337 PMID:8347784
- Mashberg A, Boffetta P, Winkelman R, Garfinkel L (1993). Tobacco smoking, alcohol drinking, and cancer of the oral cavity and oropharynx among U.S. veterans. *Cancer*, 72: 1369–1375. doi:10.1002/1097-0142(19930815)72:4<1369::AID-CNCR2820720436>3.0.CO;2-L PMID:8339227
- Matsuo K, Hamajima N, Hirose K *et al.* (2001). Alcohol, smoking, and dietary status and susceptibility to malignant lymphoma in Japan: results of a hospital-based case-control study at Aichi Cancer Center. *Jpn J Cancer Res*, 92: 1011–1017. PMID:11676850
- Mattioli S, Truffelli D, Baldasseroni A *et al.* (2002). Occupational risk factors for renal cell cancer: a case–control study in northern Italy. *J Occup Environ Med*, 44: 1028–1036. doi:10.1097/00043764-200211000-00009 PMID:12448354
- Mattisson I, Wirfält E, Wallström P *et al.* (2004). High fat and alcohol intakes are risk factors of postmenopausal breast cancer: a prospective study from the Malmö diet and cancer cohort. *Int J Cancer*, 110: 589–597. doi:10.1002/ijc.20166 PMID:15122593
- Mayans MV, Calvet X, Bruix J *et al.* (1990). Risk factors for hepatocellular carcinoma in Catalonia, Spain. *Int J Cancer*, 46: 378–381. doi:10.1002/ijc.2910460307 PMID:2168342
- Mayne ST, Janerich DT, Greenwald P *et al.* (1994). Dietary beta carotene and lung cancer risk in U.S. nonsmokers. *J Natl Cancer Inst*, 86: 33–38. doi:10.1093/jnci/86.1.33 PMID:8271280
- McCann SE, Freudenheim JL, Marshall JR *et al.* (2000). Diet in the epidemiology of endometrial cancer in western New York (United States). *Cancer Causes Control*, 11: 965–974. doi:10.1023/A:1026551309873 PMID:11142531
- McCann SE, Freudenheim JL, Marshall JR, Graham S (2003). Risk of human ovarian cancer is related to dietary intake of selected nutrients, phytochemicals and food groups. *J Nutr*, 133: 1937–1942. PMID:12771342
- McDonald JA, Mandel MG, Marchbanks PA *et al.* (2004). Alcohol exposure and breast cancer: results of the women’s contraceptive and reproductive experiences study. *Cancer Epidemiol Biomarkers Prev*, 13: 2106–2116. PMID:15598768
- McKinney PA, Cartwright RA, Saiu JMT *et al.* (1987). The inter-regional epidemiological study of childhood cancer (IRESCC): a case control study of aetiological

- factors in leukaemia and lymphoma. *Arch Dis Child*, 62: 279–287. doi:10.1136/adc.62.3.279 PMID:3646026
- McTiernan A, Thomas DB, Johnson LK, Roseman D (1986). Risk factors for estrogen receptor-rich and estrogen receptor-poor breast cancers. *J Natl Cancer Inst*, 77: 849–854. PMID:3463818
- Mellemgaard A, Engholm G, McLaughlin JK, Olsen JH (1994). Risk factors for renal cell carcinoma in Denmark. I. Role of socioeconomic status, tobacco use, beverages, and family history. *Cancer Causes Control*, 5: 105–113. doi:10.1007/BF01830256 PMID:8167257
- Menegaux F, Steffen C, Bellec S *et al.* (2005). Maternal coffee and alcohol consumption during pregnancy, parental smoking and risk of childhood acute leukaemia. *Cancer Detect Prev*, 29: 487–493. doi:10.1016/j.cdp.2005.06.008 PMID:16289502
- Menvielle G, Luce D, Goldberg P *et al.* (2004). Smoking, alcohol drinking and cancer risk for various sites of the larynx and hypopharynx. A case-control study in France. *Eur J Cancer Prev*, 13: 165–172. doi:10.1097/01.cej.0000130017.93310.76 PMID:15167214
- Merletti F, Boffetta P, Ciccone G *et al.* (1989). Role of tobacco and alcoholic beverages in the etiology of cancer of the oral cavity/oropharynx in Torino, Italy. *Cancer Res*, 49: 4919–4924. PMID:2758421
- Mettlin C (1989). Milk drinking, other beverage habits, and lung cancer risk. *Int J Cancer*, 43: 608–612. doi:10.1002/ijc.2910430412 PMID:2703270
- Mettlin C, Selenskas S, Natarajan N, Huben R (1989). Beta-carotene and animal fats and their relationship to prostate cancer risk. A case-control study. *Cancer*, 64: 605–612. doi:10.1002/1097-0142(19890801)64:3<605::AID-CNCR2820640307>3.0.CO;2-I PMID:2743255
- Meyer F & White E (1993). Alcohol and nutrients in relation to colon cancer in middle-aged adults. *Am J Epidemiol*, 138: 225–236. PMID:8395140
- Michaud DS, Giovannucci E, Willett WC *et al.* (2001). Coffee and alcohol consumption and the risk of pancreatic cancer in two prospective United States cohorts. *Cancer Epidemiol Biomarkers Prev*, 10: 429–437. PMID:11352851
- Millen AE, Tucker MA, Hartge P *et al.* (2004). Diet and melanoma in a case-control study. *Cancer Epidemiol Biomarkers Prev*, 13: 1042–1051. PMID:15184262
- Miller AB, Howe GR, Jain M *et al.* (1983). Food items and food groups as risk factors in a case-control study of diet and colo-rectal cancer. *Int J Cancer*, 32: 155–161. doi:10.1002/ijc.2910320204 PMID:6307893
- Mills PK, Beeson WL, Phillips RL, Fraser GE (1989). Cohort study of diet, lifestyle, and prostate cancer in Adventist men. *Cancer*, 64: 598–604. doi:10.1002/1097-0142(19890801)64:3<598::AID-CNCR2820640306>3.0.CO;2-6 PMID:2743254
- Mills PK, Beeson WL, Phillips RL, Fraser GE (1991). Bladder cancer in a low risk population: results from the Adventist Health Study. *Am J Epidemiol*, 133: 230–239. PMID:2000840

- Mills PK, Beeson WL, Phillips RL, Fraser GE (1994). Cancer incidence among California Seventh-Day Adventists, 1976–1982. *Am J Clin Nutr*, 59: Suppl1136S–1142S. PMID:8172114
- Mishina T, Watanabe H, Araki H *et al.* (1981). High risk group for prostatic cancer by matched pair analysis (author's transl) *Nippon Hinyokika Gakkai Zasshi*, 72: 1256–1279. PMID:7328963
- Mizuno S, Watanabe S, Nakamura K *et al.* (1992). A multi-institute case-control study on the risk factors of developing pancreatic cancer. *Jpn J Clin Oncol*, 22: 286–291. PMID:1434027
- Modugno F, Ness RB, Allen GO (2003). Alcohol consumption and the risk of mucinous and nonmucinous epithelial ovarian cancer. *Obstet Gynecol*, 102: 1336–1343. doi:10.1016/j.obstetgynecol.2003.08.008 PMID:14662224
- Mohamed AE, Kew MC, Groeneveld HT (1992). Alcohol consumption as a risk factor for hepatocellular carcinoma in urban southern African blacks. *Int J Cancer*, 51: 537–541. doi:10.1002/ijc.2910510406 PMID:1318267
- Momas I, Daurès J-P, Festy B *et al.* (1994). Relative importance of risk factors in bladder carcinogenesis: some new results about Mediterranean habits. *Cancer Causes Control*, 5: 326–332. doi:10.1007/BF01804983 PMID:8080944
- Mommsen S, Aagaard J, Sell A (1983). An epidemiological study of bladder cancer in a predominantly rural district. *Scand J Urol Nephrol*, 17: 307–312. doi:10.3109/00365598309182137 PMID:6689086
- Monson RR & Lyon JL (1975). Proportional mortality among alcoholics. *Cancer*, 36: 1077–1079. doi:10.1002/1097-0142(197509)36:3<1077::AID-CNCR2820360335>3.0.CO;2-E PMID:1182660
- Mori M, Harabuchi I, Miyake H *et al.* (1988). Reproductive, genetic, and dietary risk factors for ovarian cancer. *Am J Epidemiol*, 128: 771–777. PMID:3421242
- Morton LM, Holford TR, Leaderer B *et al.* (2003). Alcohol use and risk of non-Hodgkin's lymphoma among Connecticut women (United States). *Cancer Causes Control*, 14: 687–694. doi:10.1023/A:1025626208861 PMID:14575367
- Morton LM, Zheng T, Holford TR *et al.* InterLymph Consortium. (2005). Alcohol consumption and risk of non-Hodgkin lymphoma: a pooled analysis. *Lancet Oncol*, 6: 469–476. doi:10.1016/S1470-2045(05)70214-X PMID:15992695
- Moskal A, Norat T, Ferrari P, Riboli E (2007). Alcohol intake and colorectal cancer risk: a dose-response meta-analysis of published cohort studies. *Int J Cancer*, 120: 664–671. doi:10.1002/ijc.22299 PMID:17096321
- Mu LN, Zhou XF, Ding BG *et al.* (2003). A case-control study on drinking green tea and decreasing risk of cancers in the alimentary canal among cigarette smokers and alcohol drinkers *Zhonghua Liu Xing Bing Xue Za Zhi*, 24: 192–195. PMID:12816709
- Muñoz N, Plummer M, Vivas J *et al.* (2001). A case-control study of gastric cancer in Venezuela. *Int J Cancer*, 93: 417–423. doi:10.1002/ijc.1333 PMID:11433408

- Muñoz SE, Ferraroni M, La Vecchia C, Decarli A (1997). Gastric cancer risk factors in subjects with family history. *Cancer Epidemiol Biomarkers Prev*, 6: 137–140. PMID:9037565
- Muñoz SE, Navarro A, Lantieri MJ *et al.* (1998). Alcohol, methylxanthine-containing beverages, and colorectal cancer in Córdoba, Argentina. *Eur J Cancer Prev*, 7: 207–213. doi:10.1097/00008469-199806000-00005 PMID:9696929
- Murata M, Takayama K, Choi BCK, Pak AWP (1996). A nested case-control study on alcohol drinking, tobacco smoking, and cancer. *Cancer Detect Prev*, 20: 557–565. PMID:8939341
- Murtaugh MA, Ma KN, Caan BJ, Slattery ML (2004). Association of fluids from beverages with risk of rectal cancer. *Nutr Cancer*, 49: 25–31. doi:10.1207/s15327914nc4901_4 PMID:15456632
- Muscat JE, Hoffmann D, Wynder EL (1995). The epidemiology of renal cell carcinoma. A second look. *Cancer*, 75: 2552–2557. doi:10.1002/1097-0142(19950515)75:10<2552::AID-CNCR2820751023>3.0.CO;2-1 PMID:7736400
- Muscat JE & Wynder EL (1992). Tobacco, alcohol, asbestos, and occupational risk factors for laryngeal cancer. *Cancer*, 69: 2244–2251. doi:10.1002/1097-0142(19920501)69:9<2244::AID-CNCR2820690906>3.0.CO;2-O PMID:1562970
- Musicco M, Filippini G, Bordo BM *et al.* (1982). Gliomas and occupational exposure to carcinogens: case-control study. *Am J Epidemiol*, 116: 782–790. PMID:7148804
- Nakata S, Imai K, Yamanaka H (1993). Study of risk factors for prostatic cancer *Hinyokika Kyo*, 39: 1017–1024, discussion 1024–1025. PMID:8266869
- Nakata S, Sato J, Ohtake N *et al.* (1995). Epidemiological study of risk factors for bladder cancer *Hinyokika Kyo*, 41: 969–977. PMID:8578986
- Naldi L, Gallus S, Tavani A *et al.* Oncology Study Group of the Italian Group for Epidemiologic Research in Dermatology. (2004). Risk of melanoma and vitamin A, coffee and alcohol: a case-control study from Italy. *Eur J Cancer Prev*, 13: 503–508. doi:10.1097/00008469-200412000-00007 PMID:15548944
- Nam J-M, McLaughlin JK, Blot WJ (1992). Cigarette smoking, alcohol, and nasopharyngeal carcinoma: a case-control study among U.S. whites. *J Natl Cancer Inst*, 84: 619–622. doi:10.1093/jnci/84.8.619 PMID:1556772
- Nandakumar A, Anantha N, Dhar M *et al.* (1995). A case-control investigation on cancer of the ovary in Bangalore, India. *Int J Cancer*, 63: 361–365. doi:10.1002/ijc.2910630310 PMID:7591232
- Nasca PC, Baptiste MS, Field NA *et al.* (1990). An epidemiological case-control study of breast cancer and alcohol consumption. *Int J Epidemiol*, 19: 532–538. doi:10.1093/ije/19.3.532 PMID:2262245
- Nasca PC, Liu S, Baptiste MS *et al.* (1994). Alcohol consumption and breast cancer: estrogen receptor status and histology. *Am J Epidemiol*, 140: 980–988. PMID:7985660

- Navarro Silvera SA, Miller AB, Rohan TE (2005). Risk factors for thyroid cancer: a prospective cohort study. *Int J Cancer*, 116: 433–438. doi:10.1002/ijc.21079 PMID:15818623
- Negri E, La Vecchia C, Franceschi S *et al.* (1992). Attributable risks for oesophageal cancer in northern Italy. *Eur J Cancer*, 28A: 1167–1171. doi:10.1016/0959-8049(92)90479-L PMID:1627389
- Nelson RA, Levine AM, Marks G, Bernstein L (1997). Alcohol, tobacco and recreational drug use and the risk of non-Hodgkin's lymphoma. *Br J Cancer*, 76: 1532–1537. PMID:9400954
- Newcomb PA, Storer BE, Marcus PM (1993). Cancer of the large bowel in women in relation to alcohol consumption: a case-control study in Wisconsin (United States). *Cancer Causes Control*, 4: 405–411. doi:10.1007/BF00050858 PMID:8218871
- Newcomb PA, Trentham-Dietz A, Storer BE (1997). Alcohol consumption in relation to endometrial cancer risk. *Cancer Epidemiol Biomarkers Prev*, 6: 775–778. PMID:9332758
- Newton R, Ziegler J, Casabonne D *et al.* Uganda Kaposi's Sarcoma Study Group. (2007). A case-control study of cancer of the uterine cervix in Uganda. *Eur J Cancer Prev*, 16: 555–558. doi:10.1097/01.cej.0000243863.22137.b7 PMID:18090129
- Ng SK, Kabat GC, Wynder EL (1993). Oral cavity cancer in non-users of tobacco. *J Natl Cancer Inst*, 85: 743–745. doi:10.1093/jnci/85.9.743 PMID:8478961
- Nicholls P, Edwards G, Kyle E (1974). Alcoholics admitted to four hospitals in England. II. General and cause-specific mortality. *Q J Stud Alcohol*, 35: 841–855. PMID:4411913
- Nicodemus KK, Sweeney C, Folsom AR (2004). Evaluation of dietary, medical and lifestyle risk factors for incident kidney cancer in postmenopausal women. *Int J Cancer*, 108: 115–121. doi:10.1002/ijc.11532 PMID:14618625
- Nieters A, Deeg E, Becker N (2006). Tobacco and alcohol consumption and risk of lymphoma: results of a population-based case-control study in Germany. *Int J Cancer*, 118: 422–430. doi:10.1002/ijc.21306 PMID:16080191
- Niiijima T & Koiso K (1980). Incidence of prostatic cancer in Japan and Asia. *Scand J Urol Nephrol Suppl*, 55: 17–21. PMID:6938022
- Nishimoto IN, Hamada GS, Kowalski LP *et al.* São Paulo--Japan Cancer Project Gastric Cancer Study Group. (2002). Risk factors for stomach cancer in Brazil (I): a case-control study among non-Japanese Brazilians in São Paulo. *Jpn J Clin Oncol*, 32: 277–283. doi:10.1093/jjco/hyf060 PMID:12411564
- Nishino Y, Wakai K, Kondo T *et al.* JACC Study Group. (2006). Alcohol consumption and lung cancer mortality in Japanese men: results from Japan collaborative cohort (JACC) study. *J Epidemiol*, 16: 49–56. doi:10.2188/jea.16.49 PMID:16537984
- Nomura A, Grove JS, Stemmermann GN, Severson RK (1990). A prospective study of stomach cancer and its relation to diet, cigarettes, and alcohol consumption. *Cancer Res*, 50: 627–631. PMID:2297702

- Nomura A, Kolonel LN, Yoshizawa CN (1989). Smoking, alcohol, occupation, and hair dye use in cancer of the lower urinary tract. *Am J Epidemiol*, 130: 1159–1163. PMID:2589309
- Nomura AMY, Stemmermann GN, Chyou PH (1995). Gastric cancer among the Japanese in Hawaii. *Jpn J Cancer Res*, 86: 916–923. PMID:7493909
- Norell SE, Ahlbom A, Erwald R *et al.* (1986). Diet and pancreatic cancer: a case-control study. *Am J Epidemiol*, 124: 894–902. PMID:3776972
- O’Connell DL, Hulka BS, Chambless LE *et al.* (1987). Cigarette smoking, alcohol consumption, and breast cancer risk. *J Natl Cancer Inst*, 78: 229–234. PMID:3468286
- Olsen GW, Mandel JS, Gibson RW *et al.* (1989). A case-control study of pancreatic cancer and cigarettes, alcohol, coffee and diet. *Am J Public Health*, 79: 1016–1019. doi:10.2105/AJPH.79.8.1016 PMID:2751016
- Olsen J & Kronborg O (1993). Coffee, tobacco and alcohol as risk factors for cancer and adenoma of the large intestine. *Int J Epidemiol*, 22: 398–402. doi:10.1093/ije/22.3.398 PMID:8359954
- Olsen J, Sabreo S, Fasting U (1985). Interaction of alcohol and tobacco as risk factors in cancer of the laryngeal region. *J Epidemiol Community Health*, 39: 165–168. doi:10.1136/jech.39.2.165 PMID:4009100
- Olsson H & Ranstam J (1988). Head trauma and exposure to prolactin-elevating drugs as risk factors for male breast cancer. *J Natl Cancer Inst*, 80: 679–683. doi:10.1093/jnci/80.9.679 PMID:3373557
- Omenn GS, Goodman GE, Thornquist MD *et al.* (1996). Risk factors for lung cancer and for intervention effects in CARET, the Beta-Carotene and Retinol Efficacy Trial. *J Natl Cancer Inst*, 88: 1550–1559. doi:10.1093/jnci/88.21.1550 PMID:8901853
- Østerlind A, Tucker MA, Stone BJ, Jensen OM (1988). The Danish case-control study of cutaneous malignant melanoma. IV. No association with nutritional factors, alcohol, smoking or hair dyes. *Int J Cancer*, 42: 825–828. PMID:3192325
- Otani T, Iwasaki M, Yamamoto S *et al.* Japan Public Health Center-based Prospective Study Group. (2003). Alcohol consumption, smoking, and subsequent risk of colorectal cancer in middle-aged and elderly Japanese men and women: Japan Public Health Center-based prospective study. *Cancer Epidemiol Biomarkers Prev*, 12: 1492–1500. PMID:14693743
- Pacella-Norman R, Urban MI, Sitas F *et al.* (2002). Risk factors for oesophageal, lung, oral and laryngeal cancers in black South Africans. *Br J Cancer*, 86: 1751–1756. doi:10.1038/sj.bjc.6600338 PMID:12087462
- Parazzini F, La Vecchia C, D’Avanzo B *et al.* (1995a). Alcohol and endometrial cancer risk: findings from an Italian case-control study. *Nutr Cancer*, 23: 55–62. doi:10.1080/01635589509514361 PMID:7739915
- Parazzini F, La Vecchia C, Negri E (1992). Use of intrauterine device and risk of invasive cervical cancer. *Int J Epidemiol*, 21: 1030–1031. doi:10.1093/ije/21.5.1030 PMID:1468840

- Parazzini F, La Vecchia C, Negri E *et al.* (1997). Case-control study of oestrogen replacement therapy and risk of cervical cancer. *BMJ*, 315: 85–88. PMID:9240046
- Parazzini F, Moroni S, Negri E *et al.* (1995b). Selected food intake and risk of vulvar cancer. *Cancer*, 76: 2291–2296. doi:10.1002/1097-0142(19951201)76:11<2291::AID-CNCR2820761117>3.0.CO;2-W PMID:8635034
- Parker AS, Cerhan JR, Lynch CF *et al.* (2002). Gender, alcohol consumption, and renal cell carcinoma. *Am J Epidemiol*, 155: 455–462. doi:10.1093/aje/155.5.455 PMID:11867357
- Parkin DM (2006). The global health burden of infection-associated cancers in the year 2002. *Int J Cancer*, 118: 3030–3044. doi:10.1002/ijc.21731 PMID:16404738
- Parkin DM, Bray F, Ferlay J, Pisani P (2005). Global cancer statistics, 2002. *CA Cancer J Clin*, 55: 74–108. doi:10.3322/canclin.55.2.74 PMID:15761078
- Parkin DM, Vizzaino AP, Skinner ME, Ndhlovu A (1994). Cancer patterns and risk factors in the African population of southwestern Zimbabwe, 1963–1977. *Cancer Epidemiol Biomarkers Prev*, 3: 537–547. PMID:7827583
- Partanen TJ, Vainio HU, Ojajärvi IA, Kauppinen TP (1997). Pancreas cancer, tobacco smoking and consumption of alcoholic beverages: a case-control study. *Cancer Lett*, 116: 27–32. doi:10.1016/S0304-3835(97)04744-7 PMID:9177454
- Pawlega J (1992). Breast cancer and smoking, vodka drinking and dietary habits. A case-control study. *Acta Oncol*, 31: 387–392. doi:10.3109/02841869209088276 PMID:1632971
- Pedersen A, Johansen C, Grønbaek M (2003). Relations between amount and type of alcohol and colon and rectal cancer in a Danish population based cohort study. *Gut*, 52: 861–867. doi:10.1136/gut.52.6.861 PMID:12740343
- Pell S & D'Alonzo CA (1973). A five-year mortality study of alcoholics. *J Occup Med*, 15: 120–125. PMID:4685423
- Pelucchi C, La Vecchia C, Negri E *et al.* (2002b). Alcohol drinking and renal cell carcinoma in women and men. *Eur J Cancer Prev*, 11: 543–545. doi:10.1097/00008469-200212000-00006 PMID:12457106
- Pelucchi C, Mereghetti M, Talamini R *et al.* (2005). Dietary folate, alcohol consumption, and risk of ovarian cancer in an Italian case-control study. *Cancer Epidemiol Biomarkers Prev*, 14: 2056–2058. doi:10.1158/1055-9965.EPI-05-0192 PMID:16103462
- Pelucchi C, Negri E, Franceschi S *et al.* (2002a). Alcohol drinking and bladder cancer. *J Clin Epidemiol*, 55: 637–641. doi:10.1016/S0895-4356(02)00397-9 PMID:12160910
- Pernu J (1960). An epidemiological study of cancer of the digestive organs and respiratory system. A study based on 7078 cases. *Ann Med Intern Fenn Suppl*, 49: Suppl 331–117. PMID:14431928
- Peters RK, Garabrant DH, Yu MC, Mack TM (1989). A case-control study of occupational and dietary factors in colorectal cancer in young men by subsite. *Cancer Res*, 49: 5459–5468. PMID:2766308

- Peterson NB, Trentham-Dietz A, Newcomb PA *et al.* (2006). Alcohol consumption and ovarian cancer risk in a population-based case-control study. *Int J Cancer*, 119: 2423–2427. doi:10.1002/ijc.22137 PMID:16921486
- Petri AL, Tjønneland A, Gamborg M *et al.* (2004). Alcohol intake, type of beverage, and risk of breast cancer in pre- and postmenopausal women. *Alcohol Clin Exp Res*, 28: 1084–1090. doi:10.1097/01.ALC.0000130812.85638.E1 PMID:15252295
- Petridou E, Giokas G, Kuper H *et al.* (2000). Endocrine correlates of male breast cancer risk: a case-control study in Athens, Greece. *Br J Cancer*, 83: 1234–1237. doi:10.1054/bjoc.2000.1467 PMID:11027439
- Petridou E, Koukoulomatis P, Dessypris N *et al.* (2002). Why is endometrial cancer less common in Greece than in other European Union countries? *Eur J Cancer Prev*, 11: 427–432. doi:10.1097/00008469-200210000-00004 PMID:12394239
- Pickle LW, Greene MH, Ziegler RG *et al.* (1984). Colorectal cancer in rural Nebraska. *Cancer Res*, 44: 363–369. PMID:6690049
- Pierce RJ, Kune GA, Kune S *et al.* (1989). Dietary and alcohol intake, smoking pattern, occupational risk, and family history in lung cancer patients: results of a case-control study in males. *Nutr Cancer*, 12: 237–248. doi:10.1080/01635588909514023 PMID:2771801
- Platz EA, Leitzmann MF, Rimm EB *et al.* (2004). Alcohol intake, drinking patterns, and risk of prostate cancer in a large prospective cohort study. *Am J Epidemiol*, 159: 444–453. doi:10.1093/aje/kwh062 PMID:14977640
- Pogoda JM, Nichols PW, Preston-Martin S (2004). Alcohol consumption and risk of adult-onset acute myeloid leukemia: results from a Los Angeles County case-control study. *Leuk Res*, 28: 927–931. doi:10.1016/j.leukres.2004.01.007 PMID:15234569
- Pohlabein H, Jöckel K-H, Bolm-Audorff U (1999). Non-occupational risk factors for cancer of the lower urinary tract in Germany. *Eur J Epidemiol*, 15: 411–419. doi:10.1023/A:1007595809278 PMID:10442466
- Polesel J, Dal Maso L, Bagnardi V *et al.* (2005). Estimating dose-response relationship between ethanol and risk of cancer using regression spline models. *Int J Cancer*, 114: 836–841. doi:10.1002/ijc.20756 PMID:15609308
- Pollack ES, Nomura AM, Heilbrun LK *et al.* (1984). Prospective study of alcohol consumption and cancer. *N Engl J Med*, 310: 617–621. doi:10.1056/NEJM198403083101003 PMID:6694673
- Polychronopoulou A, Tzonou A, Hsieh C-C *et al.* (1993). Reproductive variables, tobacco, ethanol, coffee and somatometry as risk factors for ovarian cancer. *Int J Cancer*, 55: 402–407. doi:10.1002/ijc.2910550312 PMID:8375923
- Potter JD, Cerhan JR, Sellers TA *et al.* (1995). Progesterone and estrogen receptors and mammary neoplasia in the Iowa Women's Health Study: how many kinds of breast cancer are there? *Cancer Epidemiol Biomarkers Prev*, 4: 319–326. PMID:7655325
- Potter JD & McMichael AJ (1986). Diet and cancer of the colon and rectum: a case-control study. *J Natl Cancer Inst*, 76: 557–569. PMID:3007842

- Potter JD, Sellers TA, Folsom AR, McGovern PG (1992). Alcohol, beer, and lung cancer in postmenopausal women. The Iowa Women's Health Study. *Ann Epidemiol*, 2: 587–595. doi:10.1016/1047-2797(92)90003-9 PMID:1342310
- Prescott E, Grønbaek M, Becker U, Sørensen TIA (1999). Alcohol intake and the risk of lung cancer: influence of type of alcoholic beverage. *Am J Epidemiol*, 149: 463–470. PMID:10067906
- Preston-Martin S, Mack W, Henderson BE (1989). Risk factors for gliomas and meningiomas in males in Los Angeles County. *Cancer Res*, 49: 6137–6143. PMID:2790826
- Prior P (1988). Long-term cancer risk in alcoholism. *Alcohol Alcohol*, 23: 163–171. PMID:3390240
- Probert JL, Persad RA, Greenwood RP *et al.* (1998). Epidemiology of transitional cell carcinoma of the bladder: profile of an urban population in the south-west of England. *Br J Urol*, 82: 660–666. PMID:9839580
- Putnam SD, Cerhan JR, Parker AS *et al.* (2000). Lifestyle and anthropometric risk factors for prostate cancer in a cohort of Iowa men. *Ann Epidemiol*, 10: 361–369. doi:10.1016/S1047-2797(00)00057-0 PMID:10964002
- Qiu XQ, Qin YM, Zhang ZY, Peng RK (1999). A case-control study on risk factors of gastric cancer in Guangxi Province. *Guangxi J Prev Med.*, 5: 203–206.
- Rachtan J (2002). Alcoholic beverages consumption and lung cancer cell types among women in Poland. *Lung Cancer*, 35: 119–127. doi:10.1016/S0169-5002(01)00331-2 PMID:11804683
- Rachtan J & Sokolowski A (1997). Risk factors for lung cancer among women in Poland. *Lung Cancer*, 18: 137–145. doi:10.1016/S0169-5002(97)00062-7 PMID:9316005
- Rao DN & Desai PB (1998). Risk assessment of tobacco, alcohol and diet in cancers of base tongue and oral tongue—a case control study. *Indian J Cancer*, 35: 65–72. PMID:9849026
- Rao DN, Desai PB, Ganesh B (1999). Alcohol as an additional risk factor in laryngopharyngeal cancer in Mumbai—a case-control study. *Cancer Detect Prev*, 23: 37–44. doi:10.1046/j.1525-1500.1999.09906.x PMID:9892989
- Rashidkhani B, Åkesson A, Lindblad P, Wolk A (2005). Alcohol consumption and risk of renal cell carcinoma: a prospective study of Swedish women. *Int J Cancer*, 117: 848–853. doi:10.1002/ijc.21231 PMID:15957170
- Rauscher GH, Shore D, Sandler DP (2004). Alcohol intake and incidence of de novo adult acute leukemia. *Leuk Res*, 28: 1263–1265. doi:10.1016/j.leukres.2004.04.004 PMID:15475066
- Remontet L, Estève J, Bouvier A-M *et al.* (2003). Cancer incidence and mortality in France over the period 1978–2000. *Rev Epidemiol Sante Publique*, 51: 3–30. PMID:12684578
- Riboli E, Cornée J, Macquart-Moulin G *et al.* (1991). Cancer and polyps of the colorectum and lifetime consumption of beer and other alcoholic beverages. *Am J Epidemiol*, 134: 157–166. PMID:1862799

- Riman T, Dickman PW, Nilsson S *et al.* (2004). Some life-style factors and the risk of invasive epithelial ovarian cancer in Swedish women. *Eur J Epidemiol*, 19: 1011–1019. doi:10.1007/s10654-004-1633-8 PMID:15648594
- Robinette CD, Hrubec Z, Fraumeni JF Jr (1979). Chronic alcoholism and subsequent mortality in World War II veterans. *Am J Epidemiol*, 109: 687–700. PMID:453188
- Rodriguez T, Altieri A, Chatenoud L *et al.* (2004). Risk factors for oral and pharyngeal cancer in young adults. *Oral Oncol*, 40: 207–213. doi:10.1016/j.oraloncology.2003.08.014 PMID:14693246
- Rohan TE, Jain M, Howe GR, Miller AB (2000). Alcohol consumption and risk of breast cancer: a cohort study. *Cancer Causes Control*, 11: 239–247. doi:10.1023/A:1008933824645 PMID:10782658
- Rohrmann S, Linseisen J, Boshuizen HC *et al.* (2006). Ethanol intake and risk of lung cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Am J Epidemiol*, 164: 1103–1114. doi:10.1093/aje/kwj326 PMID:16987924
- Ron E, Kleinerman RA, Boice JD Jr *et al.* (1987). A population-based case-control study of thyroid cancer. *J Natl Cancer Inst*, 79: 1–12. PMID:3474436
- Rosenberg L, Slone D, Shapiro S *et al.* (1982). Breast cancer and alcoholic-beverage consumption. *Lancet*, 1: 267–270. doi:10.1016/S0140-6736(82)90987-4 PMID:6120284
- Rosenblatt KA, Thomas DB, Jimenez LM *et al.* (1999). The relationship between diet and breast cancer in men (United States). *Cancer Causes Control*, 10: 107–113. doi:10.1023/A:1008808925665 PMID:10231158
- Rosner B 1995. *Fundamentals of biostatistics*. (4th ed.). Belmont, CA: Duxbury Press.
- Ross RK, Shimizu H, Paganini-Hill A *et al.* (1987). Case-control studies of prostate cancer in blacks and whites in southern California. *J Natl Cancer Inst*, 78: 869–874. PMID:3471995
- Rossing MA, Cushing KL, Voigt LF *et al.* (2000). Risk of papillary thyroid cancer in women in relation to smoking and alcohol consumption. *Epidemiology*, 11: 49–54. doi:10.1097/00001648-200001000-00011 PMID:10615843
- Rothman KS, Greenland S 1998. *Modern Epidemiology*. 2nd Ed. Philadelphia: Lippincott-Raven.
- Royo-Bordonada MA, Martín-Moreno JM, Guallar E *et al.* (1997). Alcohol intake and risk of breast cancer: the euramic study. *Neoplasma*, 44: 150–156. PMID:9372855
- Ruano-Ravina A, Figueiras A, Barros-Dios JM (2004). Type of wine and risk of lung cancer: a case-control study in Spain. *Thorax*, 59: 981–985. doi:10.1136/thx.2003.018861 PMID:15516476
- Ryan P, Lee MW, North B, McMichael AJ (1992). Risk factors for tumors of the brain and meninges: results from the Adelaide Adult Brain Tumor Study. *Int J Cancer*, 51: 20–27. doi:10.1002/ijc.2910510105 PMID:1563840
- Sakata K, Hoshiyama Y, Morioka S *et al.* JACC Study Group. (2005). Smoking, alcohol drinking and esophageal cancer: findings from the JACC Study. *J Epidemiol*, 15: Suppl 2S212–S219. doi:10.2188/jea.15.S212 PMID:16127236

- Sanderson RJ, de Boer MF, Damhuis RAM *et al.* (1997). The influence of alcohol and smoking on the incidence of oral and oropharyngeal cancer in women. *Clin Otolaryngol Allied Sci*, 22: 444–448. doi:10.1046/j.1365-2273.1997.00049.x PMID:9372256
- Sanjoaquin MA, Appleby PN, Thorogood M *et al.* (2004). Nutrition, lifestyle and colorectal cancer incidence: a prospective investigation of 10998 vegetarians and non-vegetarians in the United Kingdom. *Br J Cancer*, 90: 118–121. doi:10.1038/sj.bjc.6601441 PMID:14710217
- Sankaranarayanan R, Duffy SW, Nair MK *et al.* (1990). Tobacco and alcohol as risk factors in cancer of the larynx in Kerala, India. *Int J Cancer*, 45: 879–882. doi:10.1002/ijc.2910450517 PMID:2335391
- Sasazuki S, Sasaki S, Tsugane S Japan Public Health Center Study Group. (2002). Cigarette smoking, alcohol consumption and subsequent gastric cancer risk by subsite and histologic type. *Int J Cancer*, 101: 560–566. doi:10.1002/ijc.10649 PMID:12237898
- Schatzkin A, Jones DY, Hoover RN *et al.* (1987). Alcohol consumption and breast cancer in the epidemiologic follow-up study of the first National Health and Nutrition Examination Survey. *N Engl J Med*, 316: 1169–1173. doi:10.1056/NEJM198705073161901 PMID:3574367
- Schildt E-B, Eriksson M, Hardell L, Magnuson A (1998). Oral snuff, smoking habits and alcohol consumption in relation to oral cancer in a Swedish case-control study. *Int J Cancer*, 77: 341–346. doi:10.1002/(SICI)1097-0215(19980729)77:3<341::AID-IJC6>3.0.CO;2-O PMID:9663593
- Schlecht NF, Franco EL, Pintos J *et al.* (1999). Interaction between tobacco and alcohol consumption and the risk of cancers of the upper aero-digestive tract in Brazil. *Am J Epidemiol*, 150: 1129–1137. PMID:10588073
- Schlecht NF, Pintos J, Kowalski LP, Franco EL (2001). Effect of type of alcoholic beverage on the risks of upper aerodigestive tract cancers in Brazil. *Cancer Causes Control*, 12: 579–587. doi:10.1023/A:1011226520220 PMID:11552705
- Schmidt W & Popham RE (1981). The role of drinking and smoking in mortality from cancer and other causes in male alcoholics. *Cancer*, 47: 1031–1041. doi:10.1002/1097-0142(19810301)47:5<1031::AID-CNCR2820470534>3.0.CO;2-C PMID:7226036
- Schoonen WM, Salinas CA, Kiemeny LALM, Stanford JL (2005). Alcohol consumption and risk of prostate cancer in middle-aged men. *Int J Cancer*, 113: 133–140. doi:10.1002/ijc.20528 PMID:15386436
- Schouten LJ, Zeegers MPA, Goldbohm RA, van den Brandt PA (2004). Alcohol and ovarian cancer risk: results from the Netherlands Cohort Study. *Cancer Causes Control*, 15: 201–209. doi:10.1023/B:CACO.0000019512.71560.2b PMID:15017133
- Schuman LM, Mandel J, Blackard C *et al.* (1977). Epidemiologic study of prostatic cancer: preliminary report. *Cancer Treat Rep*, 61: 181–186. PMID:194689
- Schuurman AG, Goldbohm RA, van den Brandt PA (1999). A prospective cohort study on consumption of alcoholic beverages in relation to prostate cancer incidence (The

- Netherlands). *Cancer Causes Control*, 10: 597–605. doi:10.1023/A:1008925103542 PMID:10616828
- Schüz J, Kaletsch U, Meinert R *et al.* (2001). Risk factors for neuroblastoma at different stages of disease. Results from a population-based case-control study in Germany. *J Clin Epidemiol*, 54: 702–709. doi:10.1016/S0895-4356(00)00339-5 PMID:11438411
- Schwartz D, Lellouch J, Flamant R, Denoix PF (1962). Alcohol and cancer. Results of a retrospective investigation. *Rev Fr Etud Clin Biol*, 7: 590–604. PMID:13987358
- Schwartz SM, Doody DR, Fitzgibbons ED *et al.* (2001). Oral squamous cell cancer risk in relation to alcohol consumption and alcohol dehydrogenase-3 genotypes. *Cancer Epidemiol Biomarkers Prev*, 10: 1137–1144. PMID:11700261
- Sellers TA, Vierkant RA, Cerhan JR *et al.* (2002). Interaction of dietary folate intake, alcohol, and risk of hormone receptor-defined breast cancer in a prospective study of postmenopausal women. *Cancer Epidemiol Biomarkers Prev*, 11: 1104–1107. PMID:12376515
- Sesso HD, Paffenbarger RS Jr, Lee IM (2001). Alcohol consumption and risk of prostate cancer: The Harvard Alumni Health Study. *Int J Epidemiol*, 30: 749–755. doi:10.1093/ije/30.4.749 PMID:11511598
- Severson RK, Buckley JD, Woods WG *et al.* (1993). Cigarette smoking and alcohol consumption by parents of children with acute myeloid leukemia: an analysis within morphological subgroups—a report from the Childrens Cancer Group. *Cancer Epidemiol Biomarkers Prev*, 2: 433–439. PMID:8220087
- Severson RK, Nomura AM, Grove JS, Stemmermann GN (1989). A prospective study of demographics, diet, and prostate cancer among men of Japanese ancestry in Hawaii. *Cancer Res*, 49: 1857–1860. PMID:2924323
- Sharpe CR & Siemiatycki J (2001). Case-control study of alcohol consumption and prostate cancer risk in Montréal, Canada. *Cancer Causes Control*, 12: 589–598. doi:10.1023/A:1011289108040 PMID:11552706
- Sharpe CR, Siemiatycki J, Rachet B (2002). Effects of alcohol consumption on the risk of colorectal cancer among men by anatomical subsite (Canada). *Cancer Causes Control*, 13: 483–491. doi:10.1023/A:1015700415808 PMID:12146853
- Shen J, Wang R-T, Wang L-W *et al.* (2004). A novel genetic polymorphism of inducible nitric oxide synthase is associated with an increased risk of gastric cancer. *World J Gastroenterol*, 10: 3278–3283. PMID:15484300
- Shen J, Wang RT, Xing HX *et al.* (2001). Comparison of risk factors for stomach cancer in high and low incidence areas of Yangzhong County, China. *Chin J Prev Control Chron Non-commun Dis*, 9: 114–116.
- Shibata A, Mack TM, Paganini-Hill A *et al.* (1994). A prospective study of pancreatic cancer in the elderly. *Int J Cancer*, 58: 46–49. doi:10.1002/ijc.2910580109 PMID:8014014
- Shimizu N, Nagata C, Shimizu H *et al.* (2003). Height, weight, and alcohol consumption in relation to the risk of colorectal cancer in Japan: a prospective study. *Br J Cancer*, 88: 1038–1043. doi:10.1038/sj.bjc.6600845 PMID:12671701

- Shiu M-N & Chen TH-H (2004). Impact of betel quid, tobacco and alcohol on three-stage disease natural history of oral leukoplakia and cancer: implication for prevention of oral cancer. *Eur J Cancer Prev*, 13: 39–45. doi:10.1097/00008469-200402000-00007 PMID:15075787
- Shu X-O, Brinton LA, Zheng W *et al.* (1991). A population-based case-control study of endometrial cancer in Shanghai, China. *Int J Cancer*, 49: 38–43. doi:10.1002/ijc.2910490108 PMID:1874568
- Shu X-O, Ross JA, Pendergrass TW *et al.* (1996). Parental alcohol consumption, cigarette smoking, and risk of infant leukemia: a Childrens Cancer Group study. *J Natl Cancer Inst*, 88: 24–31. doi:10.1093/jnci/88.1.24 PMID:8847721
- Sigvardsson S, Hardell L, Przybeck TR, Cloninger R (1996). Increased cancer risk among Swedish female alcoholics. *Epidemiology*, 7: 140–143. doi:10.1097/00001648-199603000-00006 PMID:8834552
- Silverman DT (2001). Risk factors for pancreatic cancer: a case-control study based on direct interviews. *Teratog Carcinog Mutagen*, 21: 7–25. doi:10.1002/1520-6866(2001)21:1<7::AID-TCM3>3.0.CO;2-A PMID:11135318
- Silverman DT, Brown LM, Hoover RN *et al.* (1995). Alcohol and pancreatic cancer in blacks and whites in the United States. *Cancer Res*, 55: 4899–4905. PMID:7585527
- Simon MS, Carman W, Wolfe R, Schottenfeld D (1991). Alcohol consumption and the risk of breast cancer: a report from the Tecumseh Community Health Study. *J Clin Epidemiol*, 44: 755–761. doi:10.1016/0895-4356(91)90127-U PMID:1941026
- Singh PN & Fraser GE (1998). Dietary risk factors for colon cancer in a low-risk population. *Am J Epidemiol*, 148: 761–774. PMID:9786231
- Sjödahl K, Lu Y, Nilsen TIL *et al.* (2007). Smoking and alcohol drinking in relation to risk of gastric cancer: a population-based, prospective cohort study. *Int J Cancer*, 120: 128–132. doi:10.1002/ijc.22157 PMID:17036324
- Slattery ML & West DW (1993). Smoking, alcohol, coffee, tea, caffeine, and theobromine: risk of prostate cancer in Utah (United States). *Cancer Causes Control*, 4: 559–563. doi:10.1007/BF00052432 PMID:8280834
- Slattery ML, West DW, Robison LM (1988). Fluid intake and bladder cancer in Utah. *Int J Cancer*, 42: 17–22. doi:10.1002/ijc.2910420105 PMID:3391705
- Slattery ML, West DW, Robison LM *et al.* (1990). Tobacco, alcohol, coffee, and caffeine as risk factors for colon cancer in a low-risk population. *Epidemiology*, 1: 141–145. doi:10.1097/00001648-199003000-00010 PMID:2073501
- Smith-Warner SA, Spiegelman D, Yaun SS *et al.* (1998). Alcohol and breast cancer in women: a pooled analysis of cohort studies. *JAMA*, 279: 535–540. doi:10.1001/jama.279.7.535 PMID:9480365
- Soler M, Chatenoud L, La Vecchia C *et al.* (1998). Diet, alcohol, coffee and pancreatic cancer: final results from an Italian study. *Eur J Cancer Prev*, 7: 455–460. doi:10.1097/00008469-199812000-00005 PMID:9926293

- Sørensen HT, Friis S, Olsen JH *et al.* (1998). Risk of liver and other types of cancer in patients with cirrhosis: a nationwide cohort study in Denmark. *Hepatology*, 28: 921–925. doi:10.1002/hep.510280404 PMID:9755226
- Spalajkovic M (1976). [Alcoholism and cancer of the larynx and hypopharynx.] *J Fr Oto-rhinolaryngol*, 25: 49–50.
- Spleissl B, Beahrs OH, Hermanek P *et al.*, editors (1990) *TNM Atlas Illustrated Guide to the TNM/p TNM Classification of Malignant Tumors*, New York, Springer-Verlag.
- Stemmermann GN, Nomura AMY, Chyou P-H, Yoshizawa C (1990). Prospective study of alcohol intake and large bowel cancer. *Dig Dis Sci*, 35: 1414–1420. doi:10.1007/BF01536750 PMID:2226103
- Stocks P (1957) Cancer incidence in North Wales and Liverpool region in relation to habits and environment. In: *British Empire Cancer Campaign 35th Annual Report*, Part II (Suppl.), London.
- Stolzenberg-Solomon RZ, Chang SC, Leitzmann MF *et al.* (2006). Folate intake, alcohol use, and postmenopausal breast cancer risk in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Am J Clin Nutr*, 83: 895–904. PMID:16600944
- Stolzenberg-Solomon RZ, Pietinen P, Barrett MJ *et al.* (2001). Dietary and other methyl-group availability factors and pancreatic cancer risk in a cohort of male smokers. *Am J Epidemiol*, 153: 680–687. doi:10.1093/aje/153.7.680 PMID:11282796
- Stryker WS, Stampfer MJ, Stein EA *et al.* (1990). Diet, plasma levels of beta-carotene and alpha-tocopherol, and risk of malignant melanoma. *Am J Epidemiol*, 131: 597–611. PMID:2316493
- Sturgeon SR, Ziegler RG, Brinton LA *et al.* (1991). Diet and the risk of vulvar cancer. *Ann Epidemiol*, 1: 427–437. doi:10.1016/1047-2797(91)90012-2 PMID:1669523
- Su LJ & Arab L (2004). Alcohol consumption and risk of colon cancer: evidence from the national health and nutrition examination survey I epidemiologic follow-up study. *Nutr Cancer*, 50: 111–119. doi:10.1207/s15327914nc5002_1 PMID:15623458
- Suadicani P, Hein HO, Gyntelberg F (1993). Height, weight, and risk of colorectal cancer. An 18-year follow-up in a cohort of 5249 men. *Scand J Gastroenterol*, 28: 285–288. doi:10.3109/00365529309096087 PMID:8446855
- Sun XW, Dai XD, Lin YJ, Shi YB (1999). The relationship between gastric cancer and the unhealthy habits and the gastric diseases. *Chin J Prev Control Chron Non-commun Dis*, 7: 220–222.
- Sun XW, Jiang JS, Dai XD *et al.* (2000). The risk factors of stomach cancer—a case-control study. *Chin J Prev Control Chron Non-commun Dis*, 5: 259–261.
- Sundby P (1967) *Alcoholism and Mortality*, Oslo, Universitetsforlaget.
- Suzuki R, Ye W, Rylander-Rudqvist T *et al.* (2005). Alcohol and postmenopausal breast cancer risk defined by estrogen and progesterone receptor status: a prospective cohort study. *J Natl Cancer Inst*, 97: 1601–1608. doi:10.1093/jnci/dji341 PMID:16264180

- Swanson CA, Wilbanks GD, Twiggs LB *et al.* (1993). Moderate alcohol consumption and the risk of endometrial cancer. *Epidemiology*, 4: 530–536. doi:10.1097/00001648-199311000-00009 PMID:8268282
- Swerdlow AJ, Huttly SRA, Smith PG (1989). Testis cancer: post-natal hormonal factors, sexual behaviour and fertility. *Int J Cancer*, 43: 549–553. doi:10.1002/ijc.2910430403 PMID:2539327
- Tajima K & Tominaga S (1985). Dietary habits and gastro-intestinal cancers: a comparative case-control study of stomach and large intestinal cancers in Nagoya, Japan. *Jpn J Cancer Res*, 76: 705–716. PMID:3930448
- Talamini G, Bassi C, Falconi M *et al.* (1999). Early detection of pancreatic cancer following the diagnosis of chronic pancreatitis. *Digestion*, 60: 554–561. doi:10.1159/000007706 PMID:10545726
- Talamini R, Barón AE, Barra S *et al.* (1990b). A case-control study of risk factor for renal cell cancer in northern Italy. *Cancer Causes Control*, 1: 125–131. doi:10.1007/BF00053163 PMID:2102282
- Talamini R, Bosetti C, La Vecchia C *et al.* (2002). Combined effect of tobacco and alcohol on laryngeal cancer risk: a case-control study. *Cancer Causes Control*, 13: 957–964. doi:10.1023/A:1021944123914 PMID:12588092
- Talamini R, Franceschi S, Barra S, La Vecchia C (1990a). The role of alcohol in oral and pharyngeal cancer in non-smokers, and of tobacco in non-drinkers. *Int J Cancer*, 46: 391–393. doi:10.1002/ijc.2910460310 PMID:2394506
- Talamini R, La Vecchia C, Decarli A *et al.* (1986). Nutrition, social factors and prostatic cancer in a Northern Italian population. *Br J Cancer*, 53: 817–821. PMID:3718835
- Talamini R, La Vecchia C, Levi F *et al.* (1998). Cancer of the oral cavity and pharynx in nonsmokers who drink alcohol and in nondrinkers who smoke tobacco. *J Natl Cancer Inst*, 90: 1901–1903. doi:10.1093/jnci/90.24.1901 PMID:9862628
- Tanaka K, Hirohata T, Takeshita S *et al.* (1992). Hepatitis B virus, cigarette smoking and alcohol consumption in the development of hepatocellular carcinoma: a case-control study in Fukuoka, Japan. *Int J Cancer*, 51: 509–514. doi:10.1002/ijc.2910510402 PMID:1318264
- Tavani A, Ferraroni M, Mezzetti M *et al.* (1998). Alcohol intake and risk of cancers of the colon and rectum. *Nutr Cancer*, 30: 213–219. doi:10.1080/01635589809514666 PMID:9631493
- Tavani A, Gallus S, Dal Maso L *et al.* (2001a). Coffee and alcohol intake and risk of ovarian cancer: an Italian case-control study. *Nutr Cancer*, 39: 29–34. doi:10.1207/S15327914nc391_4 PMID:11588899
- Tavani A, Gallus S, La Vecchia C, Franceschi S (2001b). Alcohol drinking and risk of non-Hodgkin's lymphoma. *Eur J Clin Nutr*, 55: 824–826. doi:10.1038/sj.ejcn.1601245 PMID:11593342
- Tavani A, Negri E, Franceschi S *et al.* (1994b). Alcohol consumption and risk of prostate cancer. *Nutr Cancer*, 21: 24–31. doi:10.1080/01635589409514301 PMID:8183720

- Tavani A, Pregenolato A, Negri E, La Vecchia C (1997). Alcohol consumption and risk of pancreatic cancer. *Nutr Cancer*, 27: 157–161. doi:10.1080/01635589709514518 PMID:9121943
- Terry MB, Zhang FF, Kabat G *et al.* (2006). Lifetime alcohol intake and breast cancer risk. *Ann Epidemiol*, 16: 230–240. doi:10.1016/j.annepidem.2005.06.048 PMID:16230024
- Terry P, Baron JA, Weiderpass E *et al.* (1999). Lifestyle and endometrial cancer risk: a cohort study from the Swedish Twin Registry. *Int J Cancer*, 82: 38–42. doi:10.1002/(SICI)1097-0215(19990702)82:1<38::AID-IJC8>3.0.CO;2-Q PMID:10360818
- Terry P, Nyrén O, Yuen J (1998). Protective effect of fruits and vegetables on stomach cancer in a cohort of Swedish twins. *Int J Cancer*, 76: 35–37. doi:10.1002/(SICI)1097-0215(19980330)76:1<35::AID-IJC7>3.0.CO;2-Z PMID:9533759
- Thomas DB, Jimenez LM, McTiernan A *et al.* (1992). Breast cancer in men: risk factors with hormonal implications. *Am J Epidemiol*, 135: 734–748. PMID:1350708
- Thomas DB, Qin Q, Kuypers J *et al.* (2001b). Human papillomaviruses and cervical cancer in Bangkok. II. Risk factors for in situ and invasive squamous cell cervical carcinomas. *Am J Epidemiol*, 153: 732–739. doi:10.1093/aje/153.8.732 PMID:11296144
- Thomas DB, Ray RM, Koetsawang A *et al.* (2001a). Human papillomaviruses and cervical cancer in Bangkok. I. Risk factors for invasive cervical carcinomas with human papillomavirus types 16 and 18 DNA. *Am J Epidemiol*, 153: 723–731. doi:10.1093/aje/153.8.723 PMID:11296143
- Thomas DB, Uhl CN, Hartge P (1983). Bladder cancer and alcoholic beverage consumption. *Am J Epidemiol*, 118: 720–727. PMID:6637998
- Thun MJ, Peto R, Lopez AD *et al.* (1997). Alcohol consumption and mortality among middle-aged and elderly U.S. adults. *N Engl J Med*, 337: 1705–1714. doi:10.1056/NEJM199712113372401 PMID:9392695
- Thygesen LC, Albertsen K, Johansen C, Grønbaek M (2005). Cancer incidence among Danish brewery workers. *Int J Cancer*, 116: 774–778. doi:10.1002/ijc.21076 PMID:15838831
- Tjønneland A, Christensen J, Olsen A *et al.* (2007). Alcohol intake and breast cancer risk: the European Prospective Investigation into Cancer and Nutrition (EPIC). *Cancer Causes Control*, 18: 361–373. doi:10.1007/s10552-006-0112-9 PMID:17364225
- Tjønneland A, Christensen J, Thomsen BL *et al.* (2004). Lifetime alcohol consumption and postmenopausal breast cancer rate in Denmark: a prospective cohort study. *J Nutr*, 134: 173–178. PMID:14704313
- Tjønneland A, Thomsen BL, Stripp C *et al.* (2003). Alcohol intake, drinking patterns and risk of postmenopausal breast cancer in Denmark: a prospective cohort study. *Cancer Causes Control*, 14: 277–284. doi:10.1023/A:1023640720385 PMID:12814207
- Tong WJ, Li J, Yue GQ, Liu XW (2001). Case-control study on risk factors of 120 cases of esophageal cancer and gastric cancer. *China Pub Health*, 17: 1093–1094.

- Toniolo P, Riboli E, Protta F *et al.* (1989). Breast cancer and alcohol consumption: a case-control study in northern Italy. *Cancer Res*, 49: 5203–5206. PMID:2766288
- Tønnesen H, Møller H, Andersen JR *et al.* (1994). Cancer morbidity in alcohol abusers. *Br J Cancer*, 69: 327–332. PMID:8297729
- Tran GD, Sun X-D, Abnet CC *et al.* (2005). Prospective study of risk factors for esophageal and gastric cancers in the Linxian general population trial cohort in China. *Int J Cancer*, 113: 456–463. doi:10.1002/ijc.20616 PMID:15455378
- Tung HT, Tsukuma H, Tanaka H *et al.* (1999). Risk factors for breast cancer in Japan, with special attention to anthropometric measurements and reproductive history. *Jpn J Clin Oncol*, 29: 137–146. doi:10.1093/jjco/29.3.137 PMID:10225696
- Tuyns AJ (1982) Incidence trends of laryngeal cancer in relation to national alcohol and tobacco consumption. In: Magnus, K. ed., *Trends in Cancer Incidence*, Washington DC, Hemisphere, pp. 199–214.
- Tuyns AJ, Estève J, Raymond L *et al.* (1988). Cancer of the larynx/hypopharynx, tobacco and alcohol: IARC international case-control study in Turin and Varese (Italy), Zaragoza and Navarra (Spain), Geneva (Switzerland) and Calvados (France). *Int J Cancer*, 41: 483–491. doi:10.1002/ijc.2910410403 PMID:3356483
- Tuyns AJ, Péquignot G, Gignoux M, Valla A (1982). Cancers of the digestive tract, alcohol and tobacco. *Int J Cancer*, 30: 9–11. doi:10.1002/ijc.2910300103 PMID:7118300
- Tzonou A, Day NE, Trichopoulos D *et al.* (1984). The epidemiology of ovarian cancer in Greece: a case-control study. *Eur J Cancer Clin Oncol*, 20: 1045–1052. doi:10.1016/0277-5379(84)90107-X PMID:6540687
- UK Testicular Cancer Study Group. (1994). Social, behavioural and medical factors in the aetiology of testicular cancer: results from the UK study. *Br J Cancer*, 70: 513–520. PMID:8080739
- Vachon CM, Cerhan JR, Vierkant RA, Sellers TA (2001). Investigation of an interaction of alcohol intake and family history on breast cancer risk in the Minnesota Breast Cancer Family Study. *Cancer*, 92: 240–248. doi:10.1002/1097-0142(20010715)92:2<240::AID-CNCR1315>3.0.CO;2-I PMID:11466675
- van der Gulden JWJ, Verbeek ALM, Kolk JJ (1994). Smoking and drinking habits in relation to prostate cancer. *Br J Urol*, 73: 382–389. doi:10.1111/j.1464-410X.1994.tb07601.x PMID:8199826
- van Dijk BAC, van Houwelingen KP, Witjes JA *et al.* (2001). Alcohol dehydrogenase type 3 (ADH3) and the risk of bladder cancer. *Eur Urol*, 40: 509–514. doi:10.1159/000049827 PMID:11752857
- van Duijn CM, van Steensel-Moll HA, Coebergh JW, van Zanen GE (1994). Risk factors for childhood acute non-lymphocytic leukemia: an association with maternal alcohol consumption during pregnancy? *Cancer Epidemiol Biomarkers Prev*, 3: 457–460. PMID:8000294
- van't Veer P, Kok FJ, Hermus RJ, Sturmans F (1989). Alcohol dose, frequency and age at first exposure in relation to the risk of breast cancer. *Int J Epidemiol*, 18: 511–517. doi:10.1093/ije/18.3.511 PMID:2807651

- Vaughan TL, Davis S, Kristal A, Thomas DB (1995). Obesity, alcohol, and tobacco as risk factors for cancers of the esophagus and gastric cardia: adenocarcinoma versus squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev*, 4: 85–92. PMID:7742727
- Viel J-F, Perarnau J-M, Challier B, Faivre-Nappez I (1997). Alcoholic calories, red wine consumption and breast cancer among premenopausal women. *Eur J Epidemiol*, 13: 639–643. doi:10.1023/A:1007368115200 PMID:9324209
- Villeneuve PJ, Johnson KC, Hanley AJ, Mao Y Canadian Cancer Registries Epidemiology Research Group. (2000). Alcohol, tobacco and coffee consumption and the risk of pancreatic cancer: results from the Canadian Enhanced Surveillance System case-control project. *Eur J Cancer Prev*, 9: 49–58. doi:10.1097/00008469-200002000-00007 PMID:10777010
- Vincent RG & Marchetta F (1963). The relationship of the use of tobacco and alcohol to cancer of the oral cavity, pharynx or larynx. *Am J Surg*, 106: 501–505. doi:10.1016/0002-9610(63)90137-5 PMID:14062955
- Vinceti M, Pellacani G, Malagoli C *et al.* (2005). A population-based case-control study of diet and melanoma risk in northern Italy. *Public Health Nutr*, 8: 1307–1314. doi:10.1079/PHN2005754 PMID:16372927
- Voirol M, Infante F, Raymond L *et al.* (1987). Nutritional profile of patients with cancer of the pancreas *Schweiz Med Wochenschr*, 117: 1101–1104. PMID:3672063
- Wakabayashi I, Sakamoto K, Masui H *et al.* (1994). A case-control study on risk factors for leukemia in a district of Japan. *Intern Med*, 33: 198–203. doi:10.2169/internalmedicine.33.198 PMID:8069013
- Wakai K, Kojima M, Tamakoshi K *et al.* JACC Study Group. (2005). Alcohol consumption and colorectal cancer risk: findings from the JACC Study. *J Epidemiol*, 15: Suppl 2S173–S179. doi:10.2188/jea.15.S173 PMID:16127230
- Wakai K, Ohno Y, Watanabe S *et al.* (1994). Risk factors for breast cancer among Japanese women in Tokyo: A case-control study. *J Epidemiol*, 4: 65–71.
- Walker ARP, Walker BF, Tsoetsi NG *et al.* (1992). Case-control study of prostate cancer in black patients in Soweto, South Africa. *Br J Cancer*, 65: 438–441. PMID:1558801
- Wang B, Xu DZ, Zhang Y *et al.* (2003a). Meta analysis of the relationship between tobacco consumption, drinking and esophageal cancer among males in Xi'an. *J Xi'an Jiaotong Univ Medical Science*, 24: 280–284.
- Wang J, Gao YT, Wang XL *et al.* (2005a). Prospective male cohort study on alcohol consumption and mortality in Shanghai. *Chin J Publ Health*, 21: 299–302.
- Wang J, Zhang X, Li D, Zhang YH (2005b). Analysis on heritability and risk factors of eastern resident in Inner Mongolia. *Chin J Publ Health*, 21: 788–789.
- Wang L-Y, You S-L, Lu S-N *et al.* (2003b). Risk of hepatocellular carcinoma and habits of alcohol drinking, betel quid chewing and cigarette smoking: a cohort of 2416 HBsAg-seropositive and 9421 HBsAg-seronegative male residents in

- Taiwan. *Cancer Causes Control*, 14: 241–250. doi:10.1023/A:1023636619477 PMID:12814203
- Wang W, Shi RH, Zhao ZQ (2004). Impact of CYP2E1 polymorphisms on the risk of esophageal cancer. [Nat Sci]*Acta Univ Med Nanjing*, 24: 344–347.
- Webb PM, Purdie DM, Bain CJ, Green AC (2004). Alcohol, wine, and risk of epithelial ovarian cancer. *Cancer Epidemiol Biomarkers Prev*, 13: 592–599. PMID:15066924
- Webster LA & Weiss NS Cancer and Steroid Hormone Study Group. (1989). Alcoholic beverage consumption and the risk of endometrial cancer. *Int J Epidemiol*, 18: 786–791. doi:10.1093/ije/18.4.786 PMID:2695474
- Wei EK, Giovannucci E, Wu K *et al.* (2004). Comparison of risk factors for colon and rectal cancer. *Int J Cancer*, 108: 433–442. doi:10.1002/ijc.11540 PMID:14648711
- Wei Q, Tang X, Yang Y *et al.* (1994). Risk factors of prostate cancer—a matched case-control study *Hua Xi Yi Ke Da Xue Xue Bao*, 25: 87–90. PMID:8070782
- Weiderpass E & Baron JA (2001). Cigarette smoking, alcohol consumption, and endometrial cancer risk: a population-based study in Sweden. *Cancer Causes Control*, 12: 239–247. doi:10.1023/A:1011201911664 PMID:11405329
- Weiderpass E, Ye W, Adami H-O *et al.* (2001c). Breast cancer risk in male alcoholics in Sweden. *Cancer Causes Control*, 12: 661–664. doi:10.1023/A:1011216502678 PMID:11552714
- Weiderpass E, Ye W, Mucci LA *et al.* (2001a). Alcoholism and risk for endometrial cancer. *Int J Cancer*, 93: 299–301. doi:10.1002/ijc.1334 PMID:11410881
- Weiderpass E, Ye W, Tamimi R *et al.* (2001b). Alcoholism and risk for cancer of the cervix uteri, vagina, and vulva. *Cancer Epidemiol Biomarkers Prev*, 10: 899–901. PMID:11489758
- Weir HK, Marrett LD, Kreiger N *et al.* (2000). Pre-natal and peri-natal exposures and risk of testicular germ-cell cancer. *Int J Cancer*, 87: 438–443. doi:10.1002/1097-0215(20000801)87:3<438::AID-IJC20>3.0.CO;2-1 PMID:10897052
- WestRO(1966).Epidemiologicstudyofmalignanciesoftheovaries. *Cancer*, 19: 1001–1007. doi:10.1002/1097-0142(196607)19:7<1001::AID-CNCR2820190714>3.0.CO;2-S PMID:5939299
- Westerdahl J, Olsson H, Måsbäck A *et al.* (1996). Risk of malignant melanoma in relation to drug intake, alcohol, smoking and hormonal factors. *Br J Cancer*, 73: 1126–1131. PMID:8624275
- Whittemore AS, Paffenbarger RS Jr, Anderson K, Lee JE (1985). Early precursors of site-specific cancers in college men and women. *J Natl Cancer Inst*, 74: 43–51. PMID:3855486
- Whittemore AS, Wu ML, Paffenbarger RS Jr *et al.* (1988). Personal and environmental characteristics related to epithelial ovarian cancer. II. Exposures to talcum powder, tobacco, alcohol, and coffee. *Am J Epidemiol*, 128: 1228–1240. PMID:3195564
- Willett EV, Smith AG, Dovey GJ *et al.* (2004). Tobacco and alcohol consumption and the risk of non-Hodgkin lymphoma. *Cancer Causes Control*, 15: 771–780. doi:10.1023/B:CACO.0000043427.77739.60 PMID:15456990

- Willett W (1998) *Nutritional Epidemiology*, Oxford, Oxford University Press.
- Willett WC, Reynolds RD, Cottrell-Hoehner S *et al.* (1987a). Validation of a semi-quantitative food frequency questionnaire: comparison with a 1-year diet record. *J Am Diet Assoc*, 87: 43–47. PMID:3794132
- Willett WC, Stampfer MJ, Colditz GA *et al.* (1987b). Moderate alcohol consumption and the risk of breast cancer. *N Engl J Med*, 316: 1174–1180. doi:10.1056/NEJM198705073161902 PMID:3574368
- Williams RR & Horm JW (1977). Association of cancer sites with tobacco and alcohol consumption and socioeconomic status of patients: interview study from the Third National Cancer Survey. *J Natl Cancer Inst*, 58: 525–547. PMID:557114
- Wolk A, Gridley G, Niwa S *et al.* (1996). International renal cell cancer study. VII. Role of diet. *Int J Cancer*, 65: 67–73. doi:10.1002/(SICI)1097-0215(19960103)65:1<67::AID-IJC12>3.0.CO;2-F PMID:8543399
- Woodson K, Albanes D, Tangrea JA *et al.* (1999). Association between alcohol and lung cancer in the alpha-tocopherol, beta-carotene cancer prevention study in Finland. *Cancer Causes Control*, 10: 219–226. doi:10.1023/A:1008911624785 PMID:10454067
- Wrensch M, Chew T, Farren G *et al.* (2003). Risk factors for breast cancer in a population with high incidence rates. *Breast Cancer Res*, 5: R88–R102. doi:10.1186/bcr605 PMID:12817999
- Wu AH, Paganini-Hill A, Ross RK, Henderson BE (1987). Alcohol, physical activity and other risk factors for colorectal cancer: a prospective study. *Br J Cancer*, 55: 687–694. PMID:3620314
- Wu AH, Wan P, Bernstein L (2001). A multiethnic population-based study of smoking, alcohol and body size and risk of adenocarcinomas of the stomach and esophagus (United States). *Cancer Causes Control*, 12: 721–732. doi:10.1023/A:1011290704728 PMID:11562112
- Wu C-L, Chen S-D, Lin G-P *et al.* (2003). A case–control study on relationship between polymorphisms of CYP1A1, GSTM1 and lung cancer. *S China J Prev Med*, 29: 13–16.
- Wu IC, Lu CY, Kuo FC *et al.* (2006a). Interaction between cigarette, alcohol and betel nut use on esophageal cancer risk in Taiwan. *Eur J Clin Invest*, 36: 236–241. doi:10.1111/j.1365-2362.2006.01621.x PMID:16620285
- Wu M, Zhao JK, Hu XS *et al.* (2006b). Association of smoking, alcohol drinking and dietary factors with esophageal cancer in high- and low-risk areas of Jiangsu Province, China. *World J Gastroenterol*, 12: 1686–1693. PMID:16586535
- Wu SQ & Yao FY (1994). A case–control study on the risk factors of stomach cancer in Shanxi province. *Chin J Prev Control Chron Non-commun Dis*, 2: 147–149.
- Wynder EL (1952). Some practical aspects of cancer prevention. *N Engl J Med*, 246: 492–502. doi:10.1056/NEJM195203272461305 PMID:14910846

- Wynder EL, Bross IJ, Day E (1956). A study of environmental factors in cancer of the larynx. *Cancer*, 9: 86–110. doi:10.1002/1097-0142(195601/02)9:1<86::AID-CNCR2820090108>3.0.CO;2-6 PMID:13284704
- Wynder EL, Covey LS, Mabuchi K, Mushinski M (1976). Environmental factors in cancer of the larynx: a second look. *Cancer*, 38: 1591–1601. doi:10.1002/1097-0142(197610)38:4<1591::AID-CNCR2820380425>3.0.CO;2-R PMID:991080
- Wynder EL, Hall NE, Polansky M (1983). Epidemiology of coffee and pancreatic cancer. *Cancer Res*, 43: 3900–3906. PMID:6861152
- Wynder EL, Kajitani T, Ishikawa S *et al.* (1969). Environmental factors of cancer of the colon and rectum. II. Japanese epidemiological data. *Cancer*, 23: 1210–1220. doi:10.1002/1097-0142(196905)23:5<1210::AID-CNCR2820230530>3.0.CO;2-M PMID:5778239
- Wynder EL, Mabuchi K, Whitmore WF Jr (1971). Epidemiology of cancer of the prostate. *Cancer*, 28: 344–360. doi:10.1002/1097-0142(197108)28:2<344::AID-CNCR2820280214>3.0.CO;2-# PMID:5109447
- Wynder EL & Shigematsu T (1967). Environmental factors of cancer of the colon and rectum. *Cancer*, 20: 1520–1561. doi:10.1002/1097-0142(196709)20:9<1520::AID-CNCR2820200920>3.0.CO;2-3 PMID:6038396
- Yamada K, Araki S, Tamura M *et al.* (1997). Case-control study of colorectal carcinoma in situ and cancer in relation to cigarette smoking and alcohol use (Japan). *Cancer Causes Control*, 8: 780–785. doi:10.1023/A:1018491607454 PMID:9328201
- Yan ZR, Wang XQ, Yan G *et al.* (2004). Relationship between the expression of Bcl-2 gene in esophageal cancer and the risk factors of esophageal cancer. *J Lanzhou Med Coll*, 30: 7–9.
- Yang C-X, Matsuo K, Ito H *et al.* (2005). Esophageal cancer risk by ALDH2 and ADH2 polymorphisms and alcohol consumption: exploration of gene-environment and gene-gene interactions. *Asian Pac J Cancer Prev*, 6: 256–262. PMID:16235983
- Yang J, Shen HB, Niu JY *et al.* (2004). Relationship between polymorphisms interleukin-1 B-31 and interleukin-1receptor antagonist gene and susceptibility to gastric cancer. *Acta Univ Med Nanjing*, 24: 193–197.
- Yang Q, Olshan AF, Bondy ML *et al.* (2000). Parental smoking and alcohol consumption and risk of neuroblastoma. *Cancer Epidemiol Biomarkers Prev*, 9: 967–972. PMID:11008916
- Ye W, Ekström AM, Hansson LE *et al.* (1999). Tobacco, alcohol and the risk of gastric cancer by sub-site and histologic type. *Int J Cancer*, 83: 223–229. doi:10.1002/(SICI)1097-0215(19991008)83:2<223::AID-IJC13>3.0.CO;2-M PMID:10471531
- Ye W, Lagergren J, Weiderpass E *et al.* (2002). Alcohol abuse and the risk of pancreatic cancer. *Gut*, 51: 236–239. doi:10.1136/gut.51.2.236 PMID:12117886
- Ye WM, Yi YN, Luo RX (1998). A case-control study on diet and gastric cancer *Zhonghua Yu Fang Yi Xue Za Zhi*, 32: 100–102. PMID:10322809

- Yen M-L, Yen BL, Bai C-H, Lin RS (2003). Risk factors for ovarian cancer in Taiwan: a case-control study in a low-incidence population. *Gynecol Oncol*, 89: 318–324. doi:10.1016/S0090-8258(03)00088-X PMID:12713998
- Yokoyama A, Kato H, Yokoyama T *et al.* (2006). Esophageal squamous cell carcinoma and aldehyde dehydrogenase-2 genotypes in Japanese females. *Alcohol Clin Exp Res*, 30: 491–500. doi:10.1111/j.1530-0277.2006.00053.x PMID:16499490
- Yong L-C, Brown CC, Schatzkin A *et al.* (1997). Intake of vitamins E, C, and A and risk of lung cancer. The NHANES I epidemiologic followup study. First National Health and Nutrition Examination Survey. *Am J Epidemiol*, 146: 231–243. PMID:9247007
- Yoo KY, Tajima K, Miura S *et al.* (1997). Breast cancer risk factors according to combined estrogen and progesterone receptor status: a case-control analysis. *Am J Epidemiol*, 146: 307–314. PMID:9270409
- Young TB (1989). A case-control study of breast cancer and alcohol consumption habits. *Cancer*, 64: 552–558. doi:10.1002/1097-0142(19890715)64:2<552::AID-CNCR2820640233>3.0.CO;2-Y PMID:2736501
- Yu H, Harris RE, Wynder EL (1988). Case-control study of prostate cancer and socioeconomic factors. *Prostate*, 13: 317–325. doi:10.1002/pros.2990130407 PMID:3217278
- Yu MC, Mack TM, Hanisch R *et al.* (1986). Cigarette smoking, obesity, diuretic use, and coffee consumption as risk factors for renal cell carcinoma. *J Natl Cancer Inst*, 77: 351–356. PMID:3461197
- Yuan J-M, Govindarajan S, Arakawa K, Yu MC (2004). Synergism of alcohol, diabetes, and viral hepatitis on the risk of hepatocellular carcinoma in blacks and whites in the U.S. *Cancer*, 101: 1009–1017. doi:10.1002/cncr.20427 PMID:15329910
- Yuan JM, Ross RK, Gao YT *et al.* (1997). Follow up study of moderate alcohol intake and mortality among middle aged men in Shanghai, China. *BMJ*, 314: 18–23. PMID:9001474
- Zagraniski RT, Kelsey JL, Walter SD (1986). Occupational risk factors for laryngeal carcinoma: Connecticut, 1975–1980. *Am J Epidemiol*, 124: 67–76. PMID:3717141
- Zang EA & Wynder EL (2001). Reevaluation of the confounding effect of cigarette smoking on the relationship between alcohol use and lung cancer risk, with larynx cancer used as a positive control. *Prev Med*, 32: 359–370. doi:10.1006/pmed.2000.0818 PMID:11304097
- Zaridze D, Borisova E, Maximovitch D, Chkhikvadze V (2000). Alcohol consumption, smoking and risk of gastric cancer: case-control study from Moscow, Russia. *Cancer Causes Control*, 11: 363–371. doi:10.1023/A:1008907924938 PMID:10843447
- Zaridze D, Lifanova Y, Maximovitch D *et al.* (1991). Diet, alcohol consumption and reproductive factors in a case-control study of breast cancer in Moscow. *Int J Cancer*, 48: 493–501. doi:10.1002/ijc.2910480404 PMID:2045197
- Zatonski W, Becher H, Lissowska J, Wahrendorf J (1991). Tobacco, alcohol, and diet in the etiology of laryngeal cancer: a population-based case-control study. *Cancer Causes Control*, 2: 3–10. doi:10.1007/BF00052355 PMID:1873431

- Zatonski WA, Boyle P, Przewozniak K *et al.* (1993). Cigarette smoking, alcohol, tea and coffee consumption and pancreas cancer risk: a case-control study from Opole, Poland. *Int J Cancer*, 53: 601–607. doi:10.1002/ijc.2910530413 PMID:8436433
- Zeegers MPA, Volovics A, Dorant E *et al.* (2001). Alcohol consumption and bladder cancer risk: results from The Netherlands Cohort Study. *Am J Epidemiol*, 153: 38–41. doi:10.1093/aje/153.1.38 PMID:11159145
- Zhang C, Shen L, Wang Y *et al.* (2002). Psychosocial factors and lung cancer development. *Chin J Lung Cancer*, 2002: 92–94.
- Zhang GS, He YT, Hou J (2000). A case control study on risk factor of esophageal cancer in Cixian county. *Sichuan J Cancer Control*, 13: 65–67.
- Zhang H, Guan S, Zhuang Y (1989). A stepwise regression analysis of the risk of lung cancer. *Journal of Jinzhou Medical College*, 10: 148–150.
- Zhang L, Wang D-S, Liu F-L *et al.* (1992). An analysis on the risk factors of lung cancer in Lanzhou city. *Gansu Environmental Study and Monitoring*, 5: 35–37.
- Zhang SW, Ma K, Zhang AH (1998). Prospective cohort study of smoking and drinking associated with death of esophageal cancer. *Chin J Publ Health*, 14: 327–329.
- Zhang SW, Ma K, Zhang AH *et al.* (1997). A prospective cohort study of smoking and drinking associated with death of lung cancer. *Chin J Publ Health*, 16: 206–208.
- Zhang Y, Kreger BE, Dorgan JF *et al.* (1999). Alcohol consumption and risk of breast cancer: the Framingham Study revisited. *Am J Epidemiol*, 149: 93–101. PMID:9921953
- Zhang Y-X, Zhou Y, Pei F-X *et al.* (1990). Study on logistic model analysis on six diseases of main death causes: The role of smoking and drinking. *Chin J Publ Health*, 9: 77–82.
- Zhang ZF, Kurtz RC, Sun M *et al.* (1996). Adenocarcinomas of the esophagus and gastric cardia: medical conditions, tobacco, alcohol, and socioeconomic factors. *Cancer Epidemiol Biomarkers Prev*, 5: 761–768. PMID:8896886
- Zhao DL, Yang YD, Chen MH *et al.* (2003). Study on risk factors of esophageal cancer in Feicheng city. *China J Cancer Prev Treat*, 10: 27–30.
- Zhao JK, Wu M, Wang XS *et al.* (2005). Risk factors of esophageal cancer in a low incidence area of Jiangsu province, China. *China Tumor*, 14: 229–231.
- Zheng T, Holford T, Chen Y *et al.* (1997). Risk of tongue cancer associated with tobacco smoking and alcohol consumption: a case-control study. *Oral Oncol*, 33: 82–85. doi:10.1016/S0964-1955(96)00056-5 PMID:9231164
- Zheng TZ, Boyle P, Hu HF *et al.* (1990). Tobacco smoking, alcohol consumption, and risk of oral cancer: a case-control study in Beijing, People's Republic of China. *Cancer Causes Control*, 1: 173–179. doi:10.1007/BF00053170 PMID:2102288
- Zheng W, Blot WJ, Shu X-O *et al.* (1992). Diet and other risk factors for laryngeal cancer in Shanghai, China. *Am J Epidemiol*, 136: 178–191. PMID:1415140
- Zheng W, McLaughlin JK, Gridley G *et al.* (1993). A cohort study of smoking, alcohol consumption, and dietary factors for pancreatic cancer (United States). *Cancer Causes Control*, 4: 477–482. doi:10.1007/BF00050867 PMID:8218880

- Zheng ZL, Huang CX, Cai L (2001). A case-control study on gastric cancer in Fujian. *Strait J Prev Med*, 7: 1-4.
- Znaor A, Brennan P, Gajalakshmi V *et al.* (2003). Independent and combined effects of tobacco smoking, chewing and alcohol drinking on the risk of oral, pharyngeal and esophageal cancers in Indian men. *Int J Cancer*, 105: 681-686. doi:10.1002/ijc.11114 PMID:12740918
- Zou H, Luo S, Yang C-Y (2005). The association between the risk of lung cancer and environmental low concentration exposure to asbestos: A nested case-control study. *Chin J Dis Control Prev*, 9: 100-103.